

Steering Bell-diagonal states

Quan Quan¹, Huangjun Zhu^{2,3*}, Si-Yuan Liu^{1‡}, Shao-Ming Fei^{4,5}, Heng Fan⁶, and Wen-Li Yang^{1,7}

¹Institute of Modern Physics, Northwest University, Xi'an 710069, China

²Institute for Theoretical Physics, University of Cologne, Cologne 50937, Germany

³Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada

⁴School of Mathematical Sciences, Capital Normal University, Beijing 100048, China

⁵Max-Planck-Institute for Mathematics in the Sciences, Leipzig 04103, Germany

⁶Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

⁷Center for Mathematics and Information Interdisciplinary Sciences, Beijing, 100048, China

*Corresponding author: hzhu1@uni-koeln.de, hzhu@pitp.ca

‡lsy5227@163.com

ABSTRACT

We investigate the steerability of two-qubit Bell-diagonal states under projective measurements by the steering party. In the simplest nontrivial scenario of two projective measurements, we solve this problem completely by virtue of the connection between the steering problem and the joint-measurement problem. A necessary and sufficient criterion is derived together with a simple geometrical interpretation. Our study shows that a Bell-diagonal state is steerable by two projective measurements iff it violates the Clauser-Horne-Shimony-Holt (CHSH) inequality, in sharp contrast with the strict hierarchy expected between steering and Bell nonlocality. We also introduce a steering measure and clarify its connections with concurrence and the volume of the steering ellipsoid. In particular, we determine the maximal concurrence and ellipsoid volume of Bell-diagonal states that are not steerable by two projective measurements. Furthermore, we explore the steerability of Bell-diagonal states under three projective measurements. A simple sufficient criterion is derived, which can detect the steerability of many states that are not steerable by two projective measurements. Finally, we generalize some of our results to arbitrary two-qubit states. Our study offers valuable insight on steering of Bell-diagonal states as well as the connections between entanglement, steering, and Bell nonlocality.

Introduction

Einstein-Podolsky-Rosen (EPR) steering,¹ as noticed by Schrödinger,² is an intermediate type of nonlocal correlation between entanglement and Bell nonlocality. In the framework of modern quantum information theory, this “spooky action” can be described as a task of entanglement verification with an untrusted party, as explained by Wiseman *et al.*^{3,4} It hinges on the question of whether Alice can convince Bob that they share an entangled state, despite the fact that Bob does not trust Alice. In order to achieve this task, Alice needs to change Bob’s state remotely in a way that would be impossible if they shared classical correlations only. Contrary to entanglement and Bell nonlocality, steering features a fundamental asymmetry because the two observers play different roles in the steering test.³⁻⁵ Recently, growing attention has been directed to steering because of its potential applications in quantum information processing, such as quantum key distribution (QKD),⁶ secure teleportation,⁷ and entanglement assisted subchannel discrimination.⁸

Two basic questions concerning steering are its detection and quantification. Recently, various steering inequalities have been introduced for detecting steering.⁹⁻¹⁴ The first such inequality was derived by Reid in 1989,⁹ which is applicable to continuous variable systems, as considered in EPR’s original argument. General theory of experimental steering criteria were developed in Ref.,¹⁰ followed by many other works.¹¹⁻¹⁴ In line with theoretical development, a loophole-free steering experiment was reported in Ref.,¹⁵ and one-way steering was demonstrated in Ref.¹⁶ Meanwhile, quantification of steering has also received increasing attention in the past few years,^{8,17,18} which leads to several useful steering measures, such as steerable weight¹⁷ and steering robustness.⁸

Despite these fruitful achievements, steering detection and quantification have remained challenging tasks, and many basic questions are poorly understood. For example, no conclusive criterion is known for determining the steerability of generic two-qubit states except for Werner states.^{3,4} Even for Bell-diagonal states, only a few partial results are known concerning their steerability, including several necessary criteria and several sufficient criteria,¹⁹⁻²¹ further progresses are thus highly desirable. In addition, many results in the literature rely heavily on numerical calculation and lack intuitive pictures. Analytical results

are quite rare since difficult optimization problems are often involved in solving steering problems.

In this work, we investigate the steerability of two-qubit Bell-diagonal states under projective measurements by the steering party. These states are appealing to both theoretical and experimental studies since they have a relatively simple structure and are particularly suitable for illustrating ideas and cultivating intuition. In addition, generic two-qubit states can be turned into Bell-diagonal states by invertible stochastic local operation and classical communication (SLOCC),²² so any progress on Bell-diagonal states may potentially help understand two-qubit states in general.

We first consider the steerability of Bell-diagonal states under the simplest nontrivial measurement setting on the steering party, that is, two projective measurements. We solve this problem completely by virtue of the connection between the steering problem and the joint-measurement problem.^{14,23–25} In particular, we derive a necessary and sufficient steering criterion analytically and providing a simple geometrical interpretation. Such analytical results are valuable but quite rare in the literature on steering. Our study leads to a measure of steering, which turns out to equal the maximal violation of the Clauser-Horne-Shimony-Holt (CHSH) inequality.^{26,27} As an implication, a Bell-diagonal state is steerable by two projective measurements iff it violates the CHSH inequality. This conclusion presents a sharp contrast with the observation that steering is necessary but usually not sufficient for Bell nonlocality.^{3,4,28} The relations between our steering measure and concurrence as well as the volume of the steering ellipsoid are then clarified. Quite surprisingly, the steering measure and the volume of the steering ellipsoid seem to display opposite behaviors for states with given concurrence.

Furthermore, we explore the steerability of Bell-diagonal states under three projective measurements. Although such problems are generally very difficult to address, we derive a nontrivial sufficient criterion, which also has a simple geometrical interpretation. This criterion can detect the steerability of many states that are not steerable by two projective measurements. The relation between entanglement and steering in this scenario is also clarified.

Finally, we discuss briefly the generalization of the above results to arbitrary two-qubit states. In particular, we derive sufficient criteria on steerability of arbitrary two-qubit states by two and three projective measurements.

Setting up the stage

Consider two remote parties, Alice and Bob, who share a bipartite quantum state ρ with reduced states ρ_A and ρ_B for the two parties, respectively. Alice can perform a collection of local measurements as characterized by a collection of positive-operator-valued measures (POVMs) $\{A_{a|x}\}_{a,x}$, where x labels the POVM and a labels the outcome in each POVM. Recall that a POVM $\{A_{a|x}\}_a$ is composed of a set of positive operators that sum up to the identity, that is, $\sum_a A_{a|x} = I$. The whole collection of POVMs $\{A_{a|x}\}_{a,x}$ is called a *measurement assemblage*. If Alice performs the measurement x and obtains the outcome a , then Bob's subnormalized reduced state is given by $\rho_{a|x} = \text{tr}_A[(A_{a|x} \otimes I)\rho]$. Note that $\sum_a \rho_{a|x} = \rho_B$ is independent of the measurement chosen by Alice, as required by the no signaling principle. The set of subnormalized states $\{\rho_{a|x}\}_a$ for a given measurement x is an *ensemble* for ρ_B , and the whole collection of ensembles $\{\rho_{a|x}\}_{a,x}$ is a *state assemblage*.¹²

The state assemblage $\{\rho_{a|x}\}_{a,x}$ is *unsteerable* if there exists a local hidden state (LHS) model:^{3,4,14,23–25}

$$\rho_{a|x} = \sum_{\lambda} p_{\rho}(a|x, \lambda) \sigma_{\lambda}, \quad (1)$$

where $p_{\rho}(a|x, \lambda) \geq 0$, $\sum_a p_{\rho}(a|x, \lambda) = 1$, and σ_{λ} are a collection of subnormalized states that sum up to ρ_B and thus form an ensemble for ρ_B . This means that Bob can interpret his conditional states $\rho_{a|x}$ as coming from the preexisting states σ_{λ} , where only the probabilities are changed due to the knowledge of Alice's measurements and outcomes.

The steering problem is closely related to the joint-measurement problem. A measurement assemblage $\{A_{a|x}\}_{a,x}$ is *compatible* or *jointly measurable*^{23–25,29,30} if there exist a POVM $\{G_{\lambda}\}$ and probabilities $p_A(a|x, \lambda)$ with $\sum_a p_A(a|x, \lambda) = 1$ such that

$$A_{a|x} = \sum_{\lambda} p_A(a|x, \lambda) G_{\lambda}. \quad (2)$$

Physically, this means that all the measurements in the assemblage can be measured jointly by performing the measurement $\{G_{\lambda}\}_{\lambda}$ and post processing the measurement data. According to the above discussion, determining the compatibility of a measurement assemblage is mathematically equivalent to determining the unsteerability of a state assemblage. Therefore, many compatibility problems can be translated into steering problems, and vice versa.^{14,23–25} This observation will play an important role in the present study.

When ρ_B is of full rank, the state assemblage $\{\rho_{a|x}\}_{a,x}$ for Bob can be turned into a measurement assemblage as follows,^{14,25}

$$B_{a|x} = \rho_B^{-1/2} \rho_{a|x} \rho_B^{-1/2}. \quad (3)$$

Note that the set of operators $\{B_{a|x}\}_a$ for a given x forms a POVM, which is referred to as Bob's *steering-equivalent observable* (or POVM).²⁵ The measurement assemblage $\{B_{a|x}\}_{a,x}$ is compatible iff the state assemblage $\{\rho_{a|x}\}_{a,x}$ is unsteerable. For example, if $\rho_{a|x} = \sum_\lambda p(a|x, \lambda) \sigma_\lambda$, then $B_{a|x} = \sum_\lambda p(a|x, \lambda) G_\lambda$ with $G_\lambda = \rho_B^{-1/2} \sigma_\lambda \rho_B^{-1/2}$; the converse follows from the same reasoning. This observation suggests a fruitful approach for understanding steering via steering-equivalent observables.

Results

Steer Bell-diagonal states by projective measurements

Any two-qubit state can be written in the following form

$$\rho = \frac{1}{4}(I \otimes I + \mathbf{a} \cdot \boldsymbol{\sigma} \otimes I + I \otimes \mathbf{b} \cdot \boldsymbol{\sigma} + \sum_{i,j=1}^3 t_{ij} \sigma_i \otimes \sigma_j), \quad (4)$$

where σ_j for $j = 1, 2, 3$ are three Pauli matrices, $\boldsymbol{\sigma}$ is the vector composed of these Pauli matrices, \mathbf{a} and \mathbf{b} are the Bloch vectors associated with the reduced states of Alice and Bob, respectively, and $T = (t_{ij})$ is the correlation matrix. The two-qubit state is a Bell-diagonal state iff $\mathbf{a} = \mathbf{b} = \mathbf{0}$,³¹ in which case we have

$$\rho = \frac{1}{4}(I \otimes I + \sum_{i,j=1}^3 t_{ij} \sigma_i \otimes \sigma_j), \quad (5)$$

with the two reduced states being completely mixed, that is, $\rho_A = \rho_B = I/2$. Bell-diagonal states are of special interest because they have a simple structure and are thus a good starting point for understanding states with more complex structure. In addition, all two-qubit states except for a set of measure zero can be turned into Bell-diagonal states by invertible SLOCC.²²

With a suitable local unitary transformation, the correlation matrix T in (5) can be turned into diagonal form, so that

$$\rho = \frac{1}{4}(I \otimes I + \sum_{j=1}^3 t_j \sigma_j \otimes \sigma_j). \quad (6)$$

As an implication of this observation, a Bell-diagonal state is steerable by one party iff it is steerable by the other party, so there is no one-way steering⁵ for Bell-diagonal states. It does not matter which party serves as the steering party in the present study.

In the case of a qubit, any projective measurement $\{A_{\pm|x}\}$ with two outcomes \pm is uniquely determined by a unit vector \mathbf{e}_x on the Bloch sphere as $A_{\pm|x} = (I \pm \mathbf{e}_x \cdot \boldsymbol{\sigma})/2$. If Alice and Bob share the Bell-diagonal state (5) and Alice performs the projective measurement determined by \mathbf{e}_x , then the two outcomes will occur with the same probability of 1/2, and the subnormalized reduced states of Bob are given by $\rho_{\pm|x} = [I \pm (T^T \mathbf{e}_x) \cdot \boldsymbol{\sigma}]/4$. Accordingly, Bob's steering-equivalent observable takes on the form

$$B_{\pm|x} = \frac{1}{2}(I \pm \mathbf{r}_x \cdot \boldsymbol{\sigma}), \quad \mathbf{r}_x = T^T \mathbf{e}_x. \quad (7)$$

Note that this observable is uniquely characterized by the subnormalized vector \mathbf{r}_x , which determines an unbiased noisy (or unsharp) von Neumann observable. Here "unbiased" means that $\text{tr}(B_{+|x}) = \text{tr}(B_{-|x}) = 1$. In this way, the correlation matrix T induces a map from projective measurements of Alice to noisy projective measurements of Bob. Alice can steer Bob's system using the measurement assemblage $\{A_{\pm|x}\}_x$ iff the set of noisy projective measurements $\{B_{\pm|x}\}_x$ is incompatible.

To see the geometric meaning of the map induced by T , note that the end point of \mathbf{r}_x lies on an ellipsoid \mathcal{E} centered at origin and characterized by the symmetric matrix $T^T T$: the three eigenvalues of $T^T T$ are the squares of the three semiaxes (some of which may vanish), and the eigenvectors determine the orientation of these semiaxes; see Fig 1. This ellipsoid encodes the set of potential noisy projective measurements of Bob induced by projective measurements of Alice. Geometrically, this ellipsoid is identical to the steering ellipsoid introduced in Refs.,^{20,32,33} which encodes the set of states to which Alice can steer Bob's system. It is also referred to as the steering ellipsoid here although the meaning is slightly different from that in Ref.^{20,32,33} Since its discovery, the steering ellipsoid has played an important role in understanding various features pertinent to entanglement and steering.^{20,21,32-34} To appreciate its significance in the current context, note that the steerability of a Bell-diagonal state by the measurement assemblage $\{A_{\pm|x}\}_x$ is completely determined by the set of vectors \mathbf{r}_x on the steering ellipsoid. Moreover, in several cases of primary interest to us, the steerability can be determined by purely geometrical means, as we shall see shortly.




Figure 1. The steering ellipsoids of three Bell-diagonal states. The ellipsoid of a Bell state (left) coincides with the Bloch sphere; the ellipsoid of a rank-2 Bell-diagonal state or an edge state (middle) is rotationally symmetric with the largest semi-axis equal to the radius of the Bloch sphere; the ellipsoid of a Werner state (right) is a sphere contained in the Bloch sphere.

Steering by two projective measurements

In this section we derive a necessary and sufficient criterion on the steerability of a Bell-diagonal state under two projective measurements. We also introduce a steering measure and illustrate its geometrical meaning. Our study shows that a Bell-diagonal state is steerable by two projective measurements iff it violates the CHSH inequality. Furthermore, we clarify the relations between entanglement, steering, and Bell nonlocality by deriving tight inequalities between the following three measures: the concurrence, the steering measure, and the volume of the steering ellipsoid.

Theorem 1. *A Bell-diagonal state with correlation matrix T is steerable by two projective measurements iff $\lambda_1 + \lambda_2 > 1$, where λ_1, λ_2 are the two larger eigenvalues of TT^T .*

Proof. Suppose Alice performs two projective measurements $\{A_{\pm|x}\} = \{(I \pm \mathbf{e}_x \cdot \boldsymbol{\sigma})/2\}$ for $x = 1, 2$. Then Bob's steering equivalent observables are given by $\{B_{\pm|x}\} = \{(I \pm \mathbf{r}_x \cdot \boldsymbol{\sigma})/2\}$, where $\mathbf{r}_x = T^T \mathbf{e}_x$ for $x = 1, 2$, as specified in (7). According to Ref.³⁵ (see also Refs.^{30,36–38}), the two observables are compatible iff

$$|\mathbf{r}_1 + \mathbf{r}_2| + |\mathbf{r}_1 - \mathbf{r}_2| \leq 2. \quad (8)$$

Note that \mathbf{r}_1 and \mathbf{r}_2 are two vectors on the steering ellipsoid, and the left hand side of the inequality is half of the perimeter of a parallelogram inscribed on the steering ellipsoid, with the plane spanned by the parallelogram passing the centre of the ellipsoid. So the Bell-diagonal state is steerable iff the maximal perimeter of such parallelograms is larger than 4. Interestingly, the maximum can be derived with a similar method used for deriving the maximal violation of the CHSH inequality,^{27,39}

$$\begin{aligned} \max_{\mathbf{e}_1, \mathbf{e}_2} \{|\mathbf{r}_1 + \mathbf{r}_2| + |\mathbf{r}_1 - \mathbf{r}_2|\} &= \max_{\mathbf{e}_1, \mathbf{e}_2} \{|T^T(\mathbf{e}_1 + \mathbf{e}_2)| + |T^T(\mathbf{e}_1 - \mathbf{e}_2)|\} = \max_{\chi, \mathbf{c}, \mathbf{c}^\perp} \{2 \cos \chi |T^T \mathbf{c}| + 2 \sin \chi |T^T \mathbf{c}^\perp|\} \\ &= 2 \max_{\mathbf{c}, \mathbf{c}^\perp} \sqrt{|T^T \mathbf{c}|^2 + |T^T \mathbf{c}^\perp|^2} = 2 \max_{\mathbf{c}, \mathbf{c}^\perp} \sqrt{\mathbf{c}^T T T^T \mathbf{c} + \mathbf{c}^{\perp T} T T^T \mathbf{c}^\perp} = 2 \sqrt{\lambda_1 + \lambda_2}, \end{aligned} \quad (9)$$

where 2χ is the angle spanned by \mathbf{e}_1 and \mathbf{e}_2 ; \mathbf{c} and \mathbf{c}^\perp are the direction vectors of $(\mathbf{e}_1 + \mathbf{e}_2)$ and $(\mathbf{e}_1 - \mathbf{e}_2)$, respectively, which are always orthogonal. Here the maximum in the last step is attained when \mathbf{c} and \mathbf{c}^\perp span the same space as that spanned by the two eigenvectors associated with the two larger eigenvalues of TT^T . The maximum over \mathbf{e}_1 and \mathbf{e}_2 can be attained when the two vectors are eigenvectors corresponding to the two larger eigenvalues of TT^T . The Bell-diagonal state is steerable by two projective measurements iff $2\sqrt{\lambda_1 + \lambda_2} > 2$, that is, $\lambda_1 + \lambda_2 > 1$. \square

The choices of \mathbf{c} and \mathbf{c}^\perp that maximize (9) are highly not unique. Therefore, the optimal projective measurements that Alice needs to perform are also not unique. Although the optimal measurements can always be chosen to be mutually unbiased as shown in the above proof, it is usually not necessary to do so. As an example, consider the Bell-diagonal state characterized by the correlation matrix $T = \text{diag}(t_1, t_2, t_3)$ with $t_1 \geq t_2 \geq |t_3|$. One choice of \mathbf{c} and \mathbf{c}^\perp reads $\mathbf{c} = (1, 0, 0)$ and $\mathbf{c}^\perp = (0, 1, 0)$, which leads to the optimal measurement directions $\mathbf{e}_1 = (t_1, t_2, 0)/\sqrt{t_1^2 + t_2^2}$ and $\mathbf{e}_2 = (t_1, -t_2, 0)/\sqrt{t_1^2 + t_2^2}$. Note that \mathbf{e}_1 and \mathbf{e}_2 are not orthogonal in general, so the corresponding projective measurements are not mutually unbiased.

The proof of Theorem 1 also suggests a steering measure of a Bell-diagonal state under two projective measurements, namely, $S := 2\sqrt{\lambda_1 + \lambda_2}$. This measure has a simple geometrical meaning: $(S/2)^2$ is equal to the sum of squares of the two larger semi-axes of the steering ellipsoid. A Bell-diagonal state is steerable in this scenario iff $S > 2$. The maximum $2\sqrt{2}$ of S is attained when $\lambda_1 = \lambda_2 = 1$, which corresponds to a Bell state. To obtain a normalized measure of steering, we may

opt for $\max\{0, (S-2)/(2\sqrt{2}-2)\}$. According to Ref.,²⁷ the maximal violation of the CHSH inequality by the Bell-diagonal state is equal to $2\sqrt{\lambda_1 + \lambda_2}$, which coincides with the steering measure S introduced here. This observation has an important implication for the relation between steering and Bell nonlocality.

Corollary 1. *A Bell-diagonal state is steerable by two projective measurements iff it violates the CHSH inequality.*

To clarify the geometric meaning of Theorem 1 and the steering measure S , it is convenient to choose a concrete Bell basis. Here we shall adopt the following choice,⁴⁰

$$|\beta_{\mu\nu}\rangle = \frac{1}{\sqrt{2}}(|0, \nu\rangle + (-1)^\mu |1, 1 \oplus \nu\rangle), \quad \mu, \nu = 0, 1. \quad (10)$$

Note that $|\beta_{11}\rangle$ is the singlet. Thanks to the choice of the Bell basis, the correlation matrices of the four Bell states are diagonal as given by $\text{diag}((-1)^\mu, -(-1)^{\mu+\nu}, (-1)^\nu)$. Up to a local unitary transformation, any Bell-diagonal state is a mixture of the four Bell states. Without loss of generality, we can focus on Bell-diagonal states of this form, whose correlation matrices are also diagonal, as in (6).




Figure 2. Geometric illustration of Bell-diagonal states steerable by two projective measurements. (left) The regular tetrahedron represents the set of Bell-diagonal states. The octahedron in green represents the set of separable states. The blue regions represent those states that are steerable by two projective measurements. (right) A face of the regular tetrahedron which represents the set of rank-3 Bell-diagonal states.

Geometrically, the set of Bell-diagonal states forms a regular tetrahedron, whose vertices correspond to the four Bell states.^{31,40} The set of separable Bell-diagonal states forms an octahedron inside the tetrahedron.^{31,40} The tetrahedron can be embedded into a cube whose sides are aligned with the three axes labelled by t_1, t_2, t_3 , as shown in Fig. 2. In this way, a Bell-diagonal state is uniquely specified by its three coordinates (t_1, t_2, t_3) . The half steering measure $S/2$ of this Bell-diagonal state is equal to the maximum over $\sqrt{t_1^2 + t_2^2}$, $\sqrt{t_2^2 + t_3^2}$, and $\sqrt{t_3^2 + t_1^2}$, which is equal to the maximal length of the three projections of (t_1, t_2, t_3) onto the three coordinate planes. Note that S is convex in t_1, t_2, t_3 and defines a norm in the three-dimensional vector space accommodating Bell-diagonal states. Each level surface of this norm is determined by three orthogonal cylinders of equal radius. In particular, the set of unsteerable Bell-diagonal states (determined by the level surface with $S = 2$) is contained in the intersection of the three solid cylinders specified by the following three inequalities, respectively,

$$t_1^2 + t_2^2 \leq 1, \quad t_2^2 + t_3^2 \leq 1, \quad t_3^2 + t_1^2 \leq 1. \quad (11)$$

In the rest of this section we clarify the relations between the following three measures: the concurrence, the steering measure S , and the volume of the steering ellipsoid. Since S is equal to the maximal violation of the CHSH inequality, our discussion is also of interest to studying Bell nonlocality.

Recall that a two-qubit state is entangled iff it has nonzero concurrence and that the concurrence of a Bell-diagonal state is given by $C = \max\{0, 2p_{\max} - 1\}$, where p_{\max} is the maximal eigenvalue of the state.⁴¹ Given a Bell-diagonal state with correlation matrix T , the normalized volume V of the steering ellipsoid is defined as $V := |\det(T)|$.²⁰ If T is diagonal, say $T = \text{diag}(t_1, t_2, t_3)$, then $V = |t_1 t_2 t_3|$. The constraints $|t_j| \leq 1$ for $j = 1, 2, 3$ imply that $0 \leq V \leq 1$, where the upper bound is saturated only for Bell states.




Figure 3. Relations between three entanglement and steering measures for Bell-diagonal states. Here C is the concurrence, S is the steering measure, and V is the normalized volume of the steering ellipsoid. The orange region in each plot indicates the range of values. The dashed lines represent edge states and the solid lines represent Werner states. In the left and right plots, the blue regions (with some overlap with the orange regions) indicate the ranges of values of $2\|T\|_F$ versus C and $2\|T\|_F$ versus V , respectively. Here $\|T\|_F$ is the Frobenius norm of the correlation matrix T that characterizes the Bell-diagonal state and is relevant to steering under three projective measurements.

Calculation shows that C, S, V satisfy the following inequalities (see Methods section for more details):

$$\frac{2\sqrt{2}}{3}(1+2C) \leq S \leq 2\sqrt{1+C^2}, \quad (12)$$

$$C^2 \leq V \leq \left(\frac{1+2C}{3}\right)^3, \quad (13)$$

$$2\sqrt{2}\sqrt[3]{V} \leq S \leq 2\sqrt{1+V}. \quad (14)$$

Here the lower bound in (12) is applicable to entangled Bell-diagonal states, while the other five bounds in (12), (13), and (14) are applicable to all Bell-diagonal states. As an implication of the above inequalities, any Bell-diagonal state with concurrence larger than $(3 - \sqrt{2})/(2\sqrt{2})$ is steerable by two projective measurements. The normalized volume of the steering ellipsoid of any separable Bell-diagonal state is bounded from above by $1/27$, in agreement with the result in Ref.,²⁰ while that of any unsteerable Bell-diagonal state is bounded from above by $1/(2\sqrt{2})$.

Two types of Bell-diagonal states deserve special attention as they saturate certain inequalities in (12), (13), and (14). A Werner state has the form

$$W_f = f|\beta_{11}\rangle\langle\beta_{11}| + \frac{1-f}{3}(I - |\beta_{11}\rangle\langle\beta_{11}|), \quad (15)$$

where $0 \leq f \leq 1$. Note that f is equal to the singlet fraction when $f \geq 1/4$. Geometrically, the Werner state lies on a diagonal of the cube in Fig. 2; conversely, any Bell-diagonal state lying on a diagonal of the cube is equivalent to a Werner state under a local unitary transformation. The correlation matrix for the Werner state has the form $T = \text{diag}(t_1, t_2, t_3)$ with $t_1 = t_2 = t_3 = (1 - 4f)/3$. Therefore, the steering ellipsoid reduces to a sphere with radius $t_1 = t_2 = t_3 = |4f - 1|/3$; see the right plot in Fig. 1. In addition,

$$C = \max\{0, 2f - 1\}, \quad S = \frac{2\sqrt{2}}{3}|4f - 1|, \quad V = \frac{|4f - 1|^3}{27}. \quad (16)$$

The Werner state is steerable by two projective measurements iff $(3\sqrt{2} + 2)/8 < f \leq 1$. It saturates the lower bound in (14) and, when $f \geq \frac{1}{2}$, also the lower bound in (12) and the upper bound in (13).

Those states lying on an edge of the tetrahedron in Fig. 2 are called *edge states* (or rank-2 Bell-diagonal states). If an edge state has two nonzero eigenvalues p and $1 - p$ with $p \geq 1/2$, then $t_{11}^2 = 1$ and $t_{22}^2 = t_{33}^2 = (2p - 1)^2$ (assuming $t_1 \geq t_2 \geq |t_3|$). Therefore, the steering ellipsoid is rotationally symmetric with the largest semiaxis equal to 1 and the other two semiaxes equal to $2p - 1$; see the middle plot in Fig. 1. In addition,

$$C = 2p - 1, \quad S = 2\sqrt{1 + (2p - 1)^2}, \quad V = (2p - 1)^2. \quad (17)$$

The edge state is steerable by two projective measurements whenever $p \neq 1/2$, that is, when it is entangled. So entanglement is sufficient to guarantee steering and Bell nonlocality in this special case. In addition, the edge state saturates the upper bounds in (12) and (14) as well as the lower bound in (13).

Fig. 3 illustrates the relations between C, S, V . When the concurrence C is large, the three measures are closely correlated to each other, while they tend to be more independent in the opposite scenario. Quite surprisingly, the normalized volume V of the steering ellipsoid seems to have a closer relation with concurrence C rather than the steering measure S . In addition, for given concurrence $C > 0$, the volume V attains the maximum when the steering measure S attains the minimum, and vice versa.

Steering by three projective measurements

In this section we explore the steerability of Bell-diagonal states under three projective measurements by the steering party. To this end, we need a criterion for determining the compatibility of three unbiased noisy projective measurements. Fortunately, this problem has been solved in Ref.,^{42,43} according to which, three noisy binary observables $\{B_{\pm|x}\}_{x=1}^3 = \{(I \pm \mathbf{r}_x \cdot \boldsymbol{\sigma})/2\}_{x=1}^3$ are compatible iff

$$\sum_{x=0}^3 |\boldsymbol{\Lambda}_x - \boldsymbol{\Lambda}_{\text{FT}}| \leq 4. \quad (18)$$

Here $\boldsymbol{\Lambda}_0 = \mathbf{r}_1 + \mathbf{r}_2 + \mathbf{r}_3$, $\boldsymbol{\Lambda}_x = 2\mathbf{r}_x - \boldsymbol{\Lambda}_0$ for $x = 1, 2, 3$, and $\boldsymbol{\Lambda}_{\text{FT}}$ denotes the Fermat-Toricelli (FT) vector of $\{\boldsymbol{\Lambda}_x\}_{x=0}^3$, which is the vector $\boldsymbol{\Lambda}$ that minimizes the total distance $\sum_{x=0}^3 |\boldsymbol{\Lambda}_x - \boldsymbol{\Lambda}|$. In general, $\boldsymbol{\Lambda}_{\text{FT}}$ has no analytical expression.^{42,43}

Given a Bell-diagonal state with correlation matrix T , suppose Alice performs three projective measurements $\{A_{\pm|x}\}_{x=1}^3 = \{(I \pm \mathbf{e}_x \cdot \boldsymbol{\sigma})/2\}_{x=1}^3$. Then Bob's steering equivalent observables are given by $\{B_{\pm|x}\}_{x=1}^3 = \{(I \pm \mathbf{r}_x \cdot \boldsymbol{\sigma})/2\}_{x=1}^3$, where $\mathbf{r}_x = T^T \mathbf{e}_x$ for $x = 1, 2, 3$. Define

$$S_3 = \frac{1}{2} \max_{\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3 \in \mathcal{E}} \sum_{x=0}^3 |\boldsymbol{\Lambda}_x - \boldsymbol{\Lambda}_{\text{FT}}| \quad (19)$$

as a steering measure of the Bell-diagonal state under three projective measurements, where \mathcal{E} is the steering ellipsoid. Then the Bell-diagonal state is steerable by three projective measurements iff $S_3 > 2$. In general, it is not easy to compute S_3 . Here we shall derive a nontrivial lower bound, which is very useful for understanding the steerability of Bell-diagonal states by three projective measurements.

When $\mathbf{r}_3 \perp \mathbf{r}_{1,2}$, the FT vector can be determined explicitly⁴³ (note that there is a typo in Ref.⁴³ about the sign),

$$\boldsymbol{\Lambda}_{\text{FT}} = \frac{|\mathbf{r}_1 - \mathbf{r}_2| - |\mathbf{r}_1 + \mathbf{r}_2|}{|\mathbf{r}_1 - \mathbf{r}_2| + |\mathbf{r}_1 + \mathbf{r}_2|} \mathbf{r}_3, \quad (20)$$

which imply that

$$\sum_{x=0}^3 |\boldsymbol{\Lambda}_x - \boldsymbol{\Lambda}_{\text{FT}}| = 2\sqrt{(|\mathbf{r}_1 - \mathbf{r}_2| + |\mathbf{r}_1 + \mathbf{r}_2|)^2 + 4\mathbf{r}_3^2}. \quad (21)$$

Theorem 2. Any Bell-diagonal state with $\|T\|_{\text{F}} > 1$ is steerable by three projective measurements, where $\|T\|_{\text{F}} = \sqrt{\text{tr}(TT^T)} = \sqrt{\text{tr}(T^T T)}$ is the Frobenius norm of the correlation matrix T .

Proof. Let $\lambda_1, \lambda_2, \lambda_3$ be the eigenvalues of TT^T in nonincreasing order and $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ the associated orthonormal eigenvectors. Let $\mathbf{r}_x = T^T \mathbf{e}_x$ for $x = 1, 2, 3$. Then $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3$ are mutually orthogonal and

$$S_3 \geq \sqrt{(|\mathbf{r}_1 - \mathbf{r}_2| + |\mathbf{r}_1 + \mathbf{r}_2|)^2 + 4\mathbf{r}_3^2} = 2\sqrt{\lambda_1 + \lambda_2 + \lambda_3} = 2\|T\|_{\text{F}}. \quad (22)$$

If the Bell-diagonal state is not steerable by three projective measurements, then $S_3 \leq 2$, so $\|T\|_{\text{F}} \leq 1$. \square

The Frobenius norm $\|T\|_{\text{F}}$ happens to be the Euclidean norm of the vector (t_1, t_2, t_3) that represents the Bell-diagonal state in Figs. 2 and 4; its square is equal to the sum of squares of the three semiaxes of the steering ellipsoid. The set of Bell-diagonal states with the same norm $\|T\|_{\text{F}}$ lies on a sphere. It is clear from the above discussion that $S_3 \geq 2\|T\|_{\text{F}} \geq S$, so any Bell-diagonal state that is steerable by two projective measurements is also steerable by three projective measurements, as expected. The converse is not true in general, as illustrated in Fig. 4. Consider the Werner state in (15) for example, we have




Figure 4. Illustration of Bell-diagonal states steerable by three projective measurements (cf. Fig. 2). (left) The regular tetrahedron represents the set of Bell-diagonal states. The octahedron in green represents the set of separable states. The blue regions represent those states that are steerable by two projective measurements, and the red regions represent those states that are not steerable by two projective measurements but steerable by three projective measurements as specified in the proof of Theorem 2. (right) A face of the regular tetrahedron which represents the set of rank-3 Bell-diagonal states.

$\|T\|_F = |4f - 1|/\sqrt{3}$, so the Werner state is steerable by three projective measurements if $1 \geq f > (\sqrt{3} + 1)/4$. By contrast, it is steerable by two projective measurements only if $1 \geq f > (3\sqrt{2} + 2)/8$.

The relations between $\|T\|_F$ and C, V can be derived with similar methods used in deriving (12) and (14), with the results

$$\frac{1}{\sqrt{3}}(1 + 2C) \leq \|T\|_F \leq \sqrt{1 + 2C^2}, \quad (23)$$

$$\sqrt{3}V^{1/3} \leq \|T\|_F \leq \sqrt{1 + 2V}. \quad (24)$$

Here the lower bound in (23) is applicable to entangled Bell-diagonal states, while the other three bounds are applicable to all Bell-diagonal states. As in (12) and (14), the two lower bounds are saturated by Werner states, while the two upper bounds are saturated by edge states; see Fig. 3. These inequalities are quite instructive to understanding the steering of Bell-diagonal states by three projective measurements given that $S_3 \geq 2\|T\|_F$. As an implication, any unsteerable Bell-diagonal state satisfies $C \leq (\sqrt{3} - 1)/2$ and $V \leq 1/(3\sqrt{3})$.

Beyond Bell-diagonal states

Here we discuss briefly the generalization of previous results to generic two-qubit states. First consider a two-qubit state with maximally mixed marginal for Bob, that is, $\mathbf{b} = \mathbf{0}$, in which case (4) reduces to

$$\rho_{\text{cano}} = \frac{1}{4}(I \otimes I + \mathbf{a} \cdot \boldsymbol{\sigma} \otimes I + \sum_{i,j=1}^3 t_{ij} \sigma_i \otimes \sigma_j), \quad (25)$$

where $T = (t_{ij})$ is the correlation matrix. Let $\lambda_1, \lambda_2, \lambda_3$ be the eigenvalues of TT^T arranged in nonincreasing order. The following two corollaries generalize Theorems 1 and 2, respectively.

Corollary 2. Any two-qubit state in (25) with $\lambda_1 + \lambda_2 > 1$ is steerable by two projective measurements from Alice.

This corollary also holds for arbitrary two-qubit states since the condition $\lambda_1 + \lambda_2 > 1$ is both necessary and sufficient for violating the CHSH inequality.^{27,39} Here we are contented to prove this special case using the connection with the joint measurement problem, in line with our previous approach.

Proof. Suppose Alice performs two projective measurements $\{A_{\pm|x}\} = \{(I \pm \mathbf{e}_x \cdot \boldsymbol{\sigma})/2\}_{x=1}^2$. Then the steering-equivalent observables for Bob are given by $\{B_{\pm|x}\}_{x=1}^2 = \{(1 \pm \alpha_x)I \pm \mathbf{r}_x \cdot \boldsymbol{\sigma}\}/2\}_{x=1}^2$, where $\alpha_x = \mathbf{e}_x \cdot \mathbf{a}$ and $\mathbf{r}_x = T^T \mathbf{e}_x$. Unlike the proof of Theorem 1, these observables are usually biased; nevertheless, the condition in (8) is still necessary for two compatible biased observables.³⁵⁻³⁸ Now the Corollary follows from the same reasoning as in the proof of Theorem 1. \square

Corollary 3. Any two-qubit state in (25) with $\|T\|_F > 1$ is steerable by three projective measurements from Alice.

Proof. Suppose Alice performs three projective measurements $\{A_{\pm|x}\} = \{(I \pm \mathbf{e}_x \cdot \boldsymbol{\sigma})/2\}_{x=1}^3$. Then the steering-inequivalent observables for Bob are given by $\{B_{\pm|x}\}_{x=1}^3 = \{[(1 \pm \alpha_x)I \pm \mathbf{r}_x \cdot \boldsymbol{\sigma}]/2\}_{x=1}^3$, where $\alpha_x = \mathbf{e}_x \cdot \mathbf{a}$ and $\mathbf{r}_x = T^T \mathbf{e}_x$ as before. Since condition (18) is still necessary for three compatible biased observables,^{42,43} the corollary follows from the same reasoning as in the proof of Theorem 2. \square

In general, any two two-qubit state ρ with invertible ρ_B (note that ρ is necessarily separable and thus unsteerable if ρ_B is not invertible) can be turned into the form of (25) by a local filtering operation of the form^{14,25,28,44}

$$I \otimes (2\rho_B)^{-1/2}(\cdot)I \otimes (2\rho_B)^{-1/2}. \quad (26)$$

Moreover, such filtering operation does not change the steerability of the state by measurements of Alice. Therefore, Corollaries 2 and 3 also provide sufficient criteria for the steerability of general two-qubit states.

Discussion

In summary, we studied systematically the steerability of Bell-diagonal states by projective measurements on the steering party. In the simplest nontrivial scenario of two projective measurements, we solved the problem completely by deriving a necessary and sufficient criterion, which has a simple geometrical interpretation. We also introduced a steering measure and proved that it is equal to the maximal violation of the CHSH inequality. This conclusion implies that a Bell-diagonal state is steerable by two projective measurements iff it violates the CHSH inequality. In addition, we clarified the relations between entanglement and steering by deriving tight inequalities satisfied by the concurrence, our steering measure, and the volume of the steering ellipsoid. Furthermore, we explored the steerability of Bell-diagonal states under three projective measurements. A simple sufficient criterion was derived, which can detect the steerability of many states that are not steerable by two projective measurements. Finally, we derive sufficient steering criteria for arbitrary two-qubit states under two and three projective measurements by generalizing the above results.

Our study provided a number of instructive analytical results on steering, which are quite rare in the literature. These results not only furnish a simple geometric picture about steering of Bell-diagonal states, but also offer valuable insight on the relations between entanglement, steering, and Bell nonlocality. They may serve as a starting point for exploring more complicated steering scenarios. In addition, our work prompts several interesting questions, which deserve further studies. For example, is the steering criterion in Theorem 2 both necessary and sufficient? Is there any upper bound on the number of measurements that are sufficient to induce steering for all steerable Bell-diagonal states? We hope that these questions will stimulate further progress on the study of steering.

Methods

Concurrence and steering measure

Here we derive the inequalities in (12), (13), and (14) in the main text, which characterize the relations between the concurrence C , the steering measure S (under two projective measurements), and the volume V of the steering ellipsoid. We also determine those Bell-diagonal states that saturate these inequalities. Similar approach can be applied to derive (23) and (24), which are pertinent to steering of Bell-diagonal states by three projective measurements.

Without loss of generality, we may assume that ρ has the form in (6) with $|t_3| \leq t_2 \leq t_1 \leq 1$. Then the spectrum of ρ is given by

$$\frac{1}{4} \{1 - t_1 - t_2 - t_3, 1 - t_1 + t_2 + t_3, 1 + t_1 - t_2 + t_3, 1 + t_1 + t_2 - t_3\}, \quad (27)$$

where the eigenvalues are arranged in nondecreasing order. The minimal and the maximal eigenvalues are respectively given by $p_{\min} = (1 - t_1 - t_2 - t_3)/4 \geq 0$ and $p_{\max} = (1 + t_1 + t_2 - t_3)/4$.

If the Bell-diagonal state is separable, that is $C = 0$, then $0 \leq p_{\min} \leq p_{\max} \leq 1/2$,³¹ which implies that

$$t_1 + t_2 + |t_3| \leq 1, \quad t_1^2 + t_2^2 \leq 1, \quad |t_1 t_2 t_3| \leq 1/27. \quad (28)$$

So the inequalities $S \leq 2\sqrt{1+C^2}$, $V \leq (1+2C)^3/27$, and $S \leq 2\sqrt{1+V}$ in (12), (13), and (14) hold for separable Bell-diagonal states. The inequality $S \leq 2\sqrt{1+C^2}$ is saturated iff $t_1 = 1$, $t_2 = t_3 = 0$, in which case ρ is an edge state with two nonzero eigenvalues equal to $1/2$. The inequality $S \leq 2\sqrt{1+V}$ is saturated under the same condition. The inequality $V \leq (1+2C)^3/27$ is saturated iff $t_1 = t_2 = |t_3| = 1/3$, in which case ρ is a Werner state which either has singlet fraction $1/2$ or is proportional to

a projector of rank 3. Here states that are equivalent to W_f in (15) under local unitary transformations are also called Werner states. The inequality $C^2 \leq V$ in (13) is trivial for separable states; it is saturated iff $V = C = 0$, that is, $t_3 = 0$, in which case the Bell-diagonal state lies on a coordinate plane in Fig. 2. The inequality $2\sqrt{2}\sqrt[3]{V} \leq S$ follows from the definitions of S and V and is applicable to both separable and entangled states. It is saturated iff $t_1 = t_2 = |t_3|$, in which case ρ is a Werner state.

If the Bell-diagonal state is entangled, then $p_{\max} > 1/2$, $C = 2p_{\max} - 1 = (t_1 + t_2 - t_3 - 1)/2$, and $t_3 = t_1 + t_2 - 1 - 2C$. The positivity of ρ and the requirement $|t_3| \leq t_2 \leq t_1$ lead to the following set of inequalities,

$$t_2 \leq t_1 \quad t_1 + t_2 \leq 1 + C, \quad t_1 + 2t_2 \geq 1 + 2C. \quad (29)$$

These inequalities determine a triangular region in the parameter space of t_1, t_2 with the following three vertices:

$$(1, C), \quad \frac{1}{2}(1 + C, 1 + C), \quad \frac{1}{3}(1 + 2C, 1 + 2C). \quad (30)$$

The maximum $1 + C^2$ of $t_1^2 + t_2^2$ under these constraints is attained iff $t_1 = 1, t_2 = -t_3 = C$, in which case the state has two nonzero eigenvalues equal to $(1 \pm C)/2$ and is thus an edge state. The minimum $2(1 + 2C)^2/9$ is attained iff $t_1 = t_2 = -t_3 = (1 + 2C)/3$, in which case the state has one eigenvalue equal to $(1 + C)/2$ and three eigenvalues equal to $(1 - C)/6$, and is thus a Werner state. By contrast, the maximum $(1 + 2C)^3/27$ of $|t_1 t_2 t_3|$ is attained exactly when $t_1^2 + t_2^2$ attains the minimum, and the minimum C^2 of $|t_1 t_2 t_3|$ is attained when $t_1^2 + t_2^2$ attains the maximum. Therefore, (12) and (13) hold for entangled Bell-diagonal states. As an immediate corollary, (14) also holds in this case.

In summary, the lower bound in (12) is applicable to entangled Bell-diagonal states, while the other five bounds in (12), (13), (14) are applicable to all Bell-diagonal states. The two inequalities $S \leq 2\sqrt{1 + C^2}$ and $S \leq 2\sqrt{1 + V}$ are saturated only for edge states. The inequality $C^2 \leq V$ is saturated only for edge states and those states with $V = 0$. The two inequalities $2\sqrt{2}(1 + 2C)/3 \leq S$ and $V \leq (1 + 2C)^3/27$ are saturated only for Werner states that have singlet fractions at least $1/2$ or Werner states that are proportional to rank-3 projectors. The inequality $2\sqrt{2}\sqrt[3]{V} \leq S$ is saturated only for Werner states. In particular, among entangled Bell-diagonal states, only edge states and Werner states with singlet fractions larger than $1/2$ can saturate these inequalities.

References

1. Einstein, A., Podolsky, B. & Rosen, N. Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? *Phys. Rev.* **47**, 777 (1935).
2. Schrödinger, E. Discussion of Probability Relations between Separated Systems. *Math. Proc. Cambr. Philos. Soc.* **31**, 555 (1935).
3. Wiseman, H. M., Jones, S. J. & Doherty, A. C. Steering, Entanglement, Nonlocality, and the Einstein-Podolsky-Rosen Paradox. *Phys. Rev. Lett.* **98**, 140402 (2007).
4. Jones, S. J., Wiseman, H. M., & Doherty, A. C. Entanglement, Einstein-Podolsky-Rosen correlations, Bell nonlocality, and steering. *Phys. Rev. A* **76**, 052116 (2007).
5. Bowles, J., Vértesi, T., Quintino, M. T. & Brunner, N. One-way Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **112**, 200402 (2014).
6. Branciard, C., Cavalcanti, E. G., Walborn, S. P., Scarani, V. & Wiseman, H. M. One-sided device-independent quantum key distribution: Security, feasibility, and the connection with steering. *Phys. Rev. A* **85**, 010301 (2012).
7. Reid, M. D. Signifying quantum benchmarks for qubit teleportation and secure quantum communication using Einstein-Podolsky-Rosen steering inequalities. *Phys. Rev. A* **88**, 062338 (2013).
8. Piani, M. & Watrous, J. Necessary and Sufficient Quantum Information Characterization of Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **114**, 060404 (2015).
9. Reid, M. D. Demonstration of the Einstein-Podolsky-Rosen paradox using nondegenerate parametric amplification. *Phys. Rev. A* **40**, 913-923 (1989).
10. Cavalcanti, E. G., Jones, S. J., Wiseman, H. M. & Reid, M. D. Experimental criteria for steering and the Einstein-Podolsky-Rosen paradox. *Phys. Rev. A* **80**, 032112 (2009).
11. Walborn, S. P., Salles, A., Gomes, R. M., Toscano, F. & Souto Ribeiro, P. H. Revealing Hidden Einstein-Podolsky-Rosen Nonlocality. *Phys. Rev. Lett.*, **106**, 130402 (2011).
12. Pusey, M. F. Negativity and steering: A stronger Peres conjecture. *Phys. Rev. A* **88**, 032313 (2013).

13. Kogias, I., Skrzypczyk, P., Cavalcanti, D., Acín, A. & Adesso, G. Hierarchy of Steering Criteria Based on Moments for All Bipartite Quantum Systems. *Phys. Rev. Lett.*, **115**, 210401 (2015).
14. Zhu, H, Hayashi, M & Chen, L, Universal steering inequalities. arXiv:1510.04711 (2015).
15. Wittmann, B. et al. Loophole-free Einstein-Podolsky-Rosen experiment via quantum steering. *New J. Phys.* **14**, 053030 (2012).
16. Händchen, V. et al. Observation of one-way Einstein-Podolsky-Rosen steering. *Nat. Photon.* **6**, 596-599 (2012).
17. Skrzypczyk, P., Navascues, M., & Cavalcanti, D. Quantifying Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **112**, 180404 (2014).
18. Kogias, I., Lee, A. R., Ragy, S. & Adesso G., Quantification of Gaussian Quantum Steering. *Phys. Rev. Lett.* **114**, 060403 (2015).
19. Chen, J., Su, H., Ye, X., Wu, C., & Oh, C. H. Einstein-Podolsky-Rosen Steerability Criterion for Two-Qubit Density Matrices. arXiv:1112.4693 (2012).
20. Jevtic, S., Pusey, M., Jennings, D. & Rudolph, T. Quantum Steering Ellipsoids. *Phys. Rev. Lett.* **113**, 020402 (2014).
21. Jevtic, S., Hall, M. J. W., Anderson, M. R., Zwierz, M. & Wiseman, H. M. Einstein-Podolsky-Rosen steering and the steering ellipsoid. *J. Opt. Soc. Am. B* **32**, 4, A40-A49 (2015).
22. Verstraete, F., Dehaene, J. & Moor, B. D. Local filtering operations on two qubits. *Phys. Rev. A* **64**, 010101(R) (2001).
23. Quintino, M. T., Vértesi, T. & Brunner, N. Joint Measurability, Einstein-Podolsky-Rosen Steering, and Bell Nonlocality. *Phys. Rev. Lett.* **113**, 160402 (2014).
24. Uola, R., Moroder, T. & Gühne, O. Joint Measurability of Generalized Measurements Implies Classicality. *Phys. Rev. Lett.* **113**, 160403 (2014).
25. Uola, R., Budroni, C., Gühne, O. & Pellonpää, J.-P. One-to-one mapping between steering and joint measurability problems. *Phys. Rev. Lett.* **115**, 230402 (2015).
26. Clauser, J. F., Horne, M. A., Shimony, A. & Holt, R. A. Proposed Experiment to Test Local Hidden-Variable Theories. *Phys. Rev. Lett.* **23**, 880 (1969).
27. Horodecki, R., Horodecki, P. & Horodecki, M. Violating Bell inequality by mixed spin-1/2 states: necessary and sufficient condition. *Phys. Lett. A* **200**, 340-344 (1995).
28. Quintino, M. T., Vértesi, T., Cavalcanti, D., Augusiak, R., Demianowicz, M., Acín, A., & Brunner, N. Inequivalence of entanglement, steering, and Bell nonlocality for general measurements. *Phys. Rev. A* **92**, 032107 (2015).
29. Heinosaari, T. & Wolf, M. M. Nondisturbing quantum measurements. *J. Math. Phys.* **51**, 092201 (2010).
30. Zhu, H. Information complementarity: A new paradigm for decoding quantum incompatibility. *Sci. Rep.* **5**, 14317 (2015).
31. Horodecki, R. & Horodecki, M. Information-theoretic aspects of inseparability of mixed states. *Phys. Rev. A* **54**, 1838 (1996).
32. Verstraete, *A STUDY OF ENTANGLEMENT IN QUANTUM INFORMATION THEORY*. (Ph. D. thesis, Katholieke Universiteit Leuven, 2002).
33. Shi, M., Jiang, F., Sun, C. & Du, J. Geometric picture of quantum discord for two-qubit quantum states. *New J. Phys.* **13**, 073016 (2011).
34. Milne, A., Jevtic, S., Jennings, D., Wiseman, H. & Rudolph, T. Quantum steering ellipsoids, extremal physical states and monogamy. *New J. Phys.* **16**, 083017 (2014).
35. Busch, P. Unsharp reality and joint measurements for spin observables. *Phys. Rev. D* **33**, 2253 (1986).
36. Busch, P., Schmidt, H. J. Coexistence of qubit effects. *Quant. Inf. Process.* **9**, 143-169 (2010).
37. Stano, P., Reitzner, D., & Heinosaari, T. Coexistence of qubit effects. *Phys. Rev. A* **78**, 012315 (2008).
38. Yu, S., Liu, N.-L., Li, L. & Oh, C. H. Joint measurement of two unsharp observables of a qubit. *Phys. Rev. A* **81**, 062116 (2010).
39. Scarani, V. The device-independent outlook on quantum physics. *Acta Physica Slovaca* **62**, 347 (2012), arXiv:1303.3081.
40. Lang, M. D. & Caves, C. M. Quantum Discord and the Geometry of Bell-Diagonal States. *Phys. Rev. Lett.* **105**, 150501 (2010).

41. Hill, S. & Wootters, W. K. Entanglement of a Pair of Quantum Bits. *Phys. Rev. Lett.* **78**, 5022 (1997).
42. Pal, R & Ghosh, S. Approximate joint measurement of qubit observables through an Arthur-Kelly model. *J. Phys. A* **44**, 485303 (2011).
43. Yu, S. & Oh, C. H. Quantum contextuality and joint measurement of three observables of a qubit. arXiv:1312.6470 (2013).
44. Bowles, J., Hirsch, F., Quintino, M. T., & Brunner, N. Sufficient criterion for guaranteeing that a two-qubit state is unsteerable. arXiv:1510.06721 (2015).

Acknowledgements

H.Z. is grateful to Christopher Fuchs and Johan Aberg for suggestions on the title. We gratefully thank for the supports by NSFC (Grant Nos. 11375141, 11425522, 11434013, 11275131). H.Z. acknowledges financial support from the Excellence Initiative of the German Federal and State Governments (ZUK81) and the DFG as well as Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research and Innovation.

Author contributions

H.Z. and Q.Q. initiated the research project and established the main results, including Theorems 1, 2, and Corollaries 1-3. H.F., S.M.F, S.Y.L. and W.L.Y. joined some discussions and provided suggestions. Q.Q. and H.Z. wrote the manuscript with advice from H.F., S.M.F, S.Y.L. and W.L.Y.

Additional information

Competing financial interests: The authors declare no competing financial interests.