

Asymptotic behavior of the Laplacian quasi-maximum likelihood estimator of affine causal processes

Jean-Marc Bardet¹, Yakoub Boularouk² and Khedidja Djaballah²

¹ S.A.M.M., Université Panthéon-Sorbonne, 90, rue de Tolbiac, 75634, Paris, France.

² Université des Sciences et de la Technologie Houari Boumédiene, Alger, Algérie

January 5, 2016

Abstract: We prove the consistency and asymptotic normality of the Laplacian Quasi-Maximum Likelihood Estimator (QMLE) for a general class of causal time series including ARMA, AR(∞), GARCH, ARCH(∞), ARMA-GARCH, APARCH, ARMA-APARCH,..., processes. We notably exhibit the advantages (moment order and robustness) of this estimator compared to the classical Gaussian QMLE. Numerical simulations confirms the accuracy of this estimator.

2000 AMS Classification Numbers: 62M10 (primary), 62F12 (secondary)

Key words: Laplacian Quasi-Maximum Likelihood Estimator; Strong consistency; Asymptotic normality; ARMA-ARCH processes.

1 Introduction

This paper is devoted to establish the consistency and the asymptotic normality of a parametric estimator for a general class of time series. This class was already defined and studied in Doukhan and Wintenberger (2007), Bardet and Wintenberger (2009) and Bardet *et al.* (2012). Hence, we will consider an observed sample (X_1, \dots, X_n) where $(X_t)_{t \in \mathbb{Z}}$ is a solution of the following equation:

$$X_t = M_{\theta_0}(X_{t-1}, X_{t-2}, \dots) \zeta_t + f_{\theta_0}(X_{t-1}, X_{t-2}, \dots), \quad t \in \mathbb{Z}, \quad (1.1)$$

where

- $\theta_0 \in \Theta \subset \mathbb{R}^d$, $d \in \mathbb{N}^*$, is an unknown vector of parameters, also called the "true" parameters;
- $(\zeta_t)_{t \in \mathbb{Z}}$ is a sequence of centred independent identically distributed random variables (i.i.d.r.v.) with symmetric probability distribution, *i.e.* $\zeta_0 \stackrel{\mathcal{L}}{=} -\zeta_0$, satisfying $E[|\zeta_0|^r] < \infty$ with $r \geq 1$ and $E[|\zeta_0|] = 1$. If $r \geq 2$, denote $\sigma_\zeta^2 = \text{Var}(\zeta_0)$;
- $(\theta, (x_n)_{n \in \mathbb{N}}) \rightarrow M_\theta((x_n)_{n \in \mathbb{N}}) \in (0, \infty)$ and $(\theta, (x_n)_{n \in \mathbb{N}}) \rightarrow f_\theta((x_n)_{n \in \mathbb{N}}) \in \mathbb{R}$ are two known applications.

For instance, if $M_{\theta_0}(X_{t-1}, X_{t-2}, \dots) = 1$ and $f_{\theta_0}(X_{t-1}, X_{t-2}, \dots) = \alpha_0 X_{t-1}$ with $|\alpha_0| < 1$ then (X_t) is a causal AR(1) process. In Doukhan and Wintenberger (2007) and Bardet and Wintenberger (2009), it was proved that all the most famous stationary time series used in econometrics, such as ARMA, AR(∞), GARCH, ARCH(∞), TARARCH, ARMA-GARCH processes can be written as a causal stationary solution of (1.1).

In Bardet and Wintenberger (2009), it was also established that under several conditions on M_θ , f_θ and if $E[|\zeta_0|^r]$ with $r \geq 2$, the usual Gaussian Quasi-Maximum Likelihood Estimator (QMLE) of θ is strongly consistent and when $r \geq 4$ it is asymptotically normal. This estimator was first defined by Weiss (1986) for ARCH processes, and the asymptotic study of this estimator was first obtained by Lumsdaine (1996) for GARCH(1, 1) processes, Berkes *et al.* (2003) for GARCH(p, q) processes, Francq and Zakoian (2004) for ARMA-GARCH processes, Mikosch and Straumann (2006) for general heteroskedastic models, and Robinson and Zaffaroni (2006) for ARCH(∞) processes. The results of Bardet and Wintenberger (2009) devoted to processes satisfying almost everywhere (1.1) as well as its multivariate generalisation, provide a general and unified framework for studying the asymptotic properties of the Gaussian QMLE.

However, the definition of the Gaussian QMLE is explicitly obtained with the assumption that (ζ_t) is a Gaussian sequence and even if it could be applied when the probability distribution of (ζ_t) is non-Gaussian, it keeps some drawbacks of this initial assumption. Indeed, the computation of this estimators requires the minimization of a least squares contrast (typically $\sum_{t=1}^n M_\theta^{-2}(X_t - f_\theta)^2$) and this induces that $r = 2$ is required for the consistency and $r = 4$ for the asymptotic normality (and therefore confidence intervals or tests). For numerous real data such requirement is sometimes too strong (for instance the kurtosis of economic data is frequently considered as infinite). Moreover, such estimator is not robust to eventual outliers. Hence, the reference probability distribution of (ζ_t) could be a Laplace one and this allows to avoid both these drawbacks. Roughly speaking this choice implies to minimize a least absolute deviations contrast (typically $\sum_{t=1}^n M_\theta^{-1}|X_t - f_\theta|$) instead of the previous least squares contrast. And therefore, $r = 1$ will be sufficient for insuring the strong consistency of this Laplacian-QMLE, while only $r = 2$ is required for the asymptotic normality (see below).

Such probability distribution choice is not new since this leads to a Least Absolute Deviations (LAD) estimation. Hence, for ARMA processes, Davis and Dunsmuir (1997) proved the consistency and asymptotic normality of the LAD estimator. For ARCH or GARCH processes, the same results concerning the LAD estimator were already established by Peng and Yao (2003), while Berkes and Horváth (2004) proved the consistency and asymptotic normality of the Laplacian-QMLE. Newey and Steigerwald (1997) considered also the estimator for other conditional heteroskedasticity models. Recently, Francq *et al.* (2011) proposed a two-stage non-Gaussian-QML estimation for GARCH models and Francq and Zakoian (2013) proposed an alternative one-step procedure, based on an appropriate non-Gaussian-QML estimator, the asymptotic properties of both these approaches were studied.

In this paper we unified all these studies of the Laplacian-QMLE in a simple framework, *i.e.* causal stationary solutions of (1.1). This notably allows to obtain known results on ARMA or GARCH but also to establish for the first time the consistency and the asymptotic normality of the Laplacian-QMLE for APARCH, ARMA-GARCH, ARMA-ARCH(∞) and ARMA-APARCH processes.

Numerical Monte-Carlo experiments were realized to illustrate the theoretical results. And there are convincing, especially when the accuracy of Laplacian-QMLE is compared with Gaussian-QMLE: except for Gaussian distribution of (ζ_t) , the Laplacian-QMLE provides a sharper estimation than Gaussian-QMLE for all the other probability distributions we considered. This is notably the case, and this is not a surprise, for a Gaussian mixing which mimics the presence of outliers.

The following Section 2 will be devoted to provide the definitions and assumptions. In Section 3 the main results are stated with numerous examples of application, while Section 4 presents the results of Monte-Carlo experiments and Section 5 contains the proofs.

2 Definition and assumptions

2.1 Definition of the estimator

Let (X_1, \dots, X_n) be an observed trajectory of X which is an a.s. solution of (1.1) where $\theta \in \Theta \subset \mathbb{R}^d$ is unknown. For estimating θ we consider the log-likelihood of (X_1, \dots, X_n) conditionally to (X_0, X_{-1}, \dots) . If g is the probability density (with respect to Lebesgue measure) of ζ_0 , then, from the affine causal definition of X , this log-likelihood can be written:

$$\log(L_\theta(X_1, \dots, X_n)) = \sum_{t=1}^n \log\left(g\left(\frac{X_t - f_\theta^t}{M_\theta^t}\right)\right)$$

where $M_\theta^t := M_\theta(X_{t-1}, X_{t-2}, \dots)$ and $f_\theta^t := f_\theta(X_{t-1}, X_{t-2}, \dots)$, with the assumption that $M_\theta^t > 0$. However, M_θ^t and f_θ^t are generally not computable since X_0, X_{-1}, \dots are unknown. Thus, a quasi-log-likelihood is considered instead of the log-likelihood and it is defined by:

$$\log(QL_\theta(X_1, \dots, X_n)) = \sum_{t=1}^n \log\left(g\left(\frac{X_t - \widehat{f}_\theta^t}{\widehat{M}_\theta^t}\right)\right),$$

with $\widehat{f}_\theta^t := f_\theta(X_{t-1}, \dots, X_1, u)$ and $\widehat{M}_\theta^t := M_\theta(X_{t-1}, \dots, X_1, u)$, where $u = (u_n)_{n \in \mathbb{N}}$ is a finitely non-zero sequence $(u_n)_{n \in \mathbb{N}}$. The choice of $(u_n)_{n \in \mathbb{N}}$ does not have any consequences on the asymptotic behaviour of L_n , and (u_n) could typically be chosen as a sequence of zeros. Finally, a Quasi-Maximum Likelihood Estimator (QMLE) is defined by:

$$\widetilde{\theta}_n := \text{Argmax}_{\theta \in \Theta} \log(QL_\theta(X_1, \dots, X_n)).$$

Usually, the Gaussian distribution is considered, with $g(x) = (2\pi)^{-1/2} e^{-x^2/2}$ for $x \in \mathbb{R}$ and this provides the Gaussian-QMLE of θ .

Here, we chose the Laplacian distribution, *i.e.*,

$$g(x) = \frac{1}{2} e^{-|x|} \quad \text{for } x \in \mathbb{R}, \quad (2.1)$$

and this implies $E[|\zeta_0|] = 1$.

Therefore, we respectively define the Laplacian-likelihood and Laplacian-quasi-likelihood by:

$$L_n(\Theta) = - \sum_{t=1}^n q_t(\Theta) \quad \text{with} \quad q_t(\Theta) = \log |M_\theta^t| + |M_\theta^t|^{-1} |X_t - f_\theta^t| \quad (2.2)$$

$$\widehat{L}_n(\theta) = - \sum_{t=1}^n \widehat{q}_t(\theta) \quad \text{with} \quad \widehat{q}_t(\theta) := \log |\widehat{M}_\theta^t| + |\widehat{M}_\theta^t|^{-1} |X_t - \widehat{f}_\theta^t|. \quad (2.3)$$

A Laplacian-QMLE $\widehat{\theta}_n$ is a maximizer of \widehat{L}_n :

$$\widehat{\theta}_n := \arg \max_{\theta \in \Theta} \widehat{L}_n(\theta).$$

We restrict the set Θ in such a way that a stationary solution (X_t) of order 1 or 2 of (1.1) exists. Additional conditions are also required for insuring the consistency and the asymptotic normality of $\widehat{\theta}_n$. More details are given now.

2.2 Existence and stationarity

As it was already done in Doukhan and Wintenberger (2007) and Bardet and Wintenberger (2009), several Lipschitz-type inequalities on f_θ and M_θ are required for obtaining the existence and r -order stationarity of an ergodic causal solution of (1.1). First, denote $\|g_\theta\|_\Theta = \sup_{\theta \in \Theta} \|g_\theta\|$ with $m \in \mathbb{N}^*$ and $\|\cdot\|$ the usual Euclidian norm. Now, let us introduce the generic symbol K for any of the functions f or M and for $g_\theta : \mathbb{R}^{\mathbb{N}} \mapsto \mathbb{R}^m$ or $\mathcal{M}_m(\mathbb{R})$, For $k = 0, 1, 2$ and some subset Θ of \mathbb{R}^d , define a Lipschitzian assumption on function K_θ :

Assumption $(\mathbf{A}_k(K, \Theta))$ $\forall x \in \mathbb{R}^\infty, \theta \in \Theta \mapsto K_\theta(x) \in \mathcal{C}^k(\Theta)$ and $\partial_\theta^k K_\theta$ satisfies $\|\partial_\theta^k K_\theta(0)\|_\Theta < \infty$ and there exists a sequence $(\alpha_j^{(k)}(K, \Theta))_j$ of nonnegative numbers such that $\forall x, y \in \mathbb{R}^{\mathbb{N}}$

$$\|\partial_\theta^k K_\theta(x) - \partial_\theta^k K_\theta(y)\|_\Theta \leq \sum_{j=1}^{\infty} \alpha_j^{(k)}(K, \Theta) |x_j - y_j|, \text{ with } \sum_{j=1}^{\infty} \alpha_j^{(k)}(K, \Theta) < \infty$$

For ensuring a stationary r -order solution of (1.1), for $r \geq 1$, define the set

$$\Theta(r) := \left\{ \theta \in \mathbb{R}^d, (\mathbf{A}_0(f, \{\theta\})) \text{ and } (\mathbf{A}_0(M, \{\theta\})) \text{ hold,} \right. \\ \left. \sum_{j=1}^{\infty} \alpha_j^{(0)}(f, \{\theta\}) + (\mathbf{E}[|\zeta_0|^r])^{1/r} \sum_{j=1}^{\infty} \alpha_j^{(0)}(M, \{\theta\}) < 1 \right\}.$$

Then, from Doukhan and Wintenberger (2007), we obtain:

Proposition 2.1. *If $\theta_0 \in \Theta(r)$ for some $r \geq 1$, then there exists a unique causal (X_t) is independent of $(\zeta_i)_{i>t}$ for $t \in \mathbb{Z}$) solution X of (1.1), which is stationary, ergodic and satisfies $\mathbf{E}[|X_0|^r] < \infty$.*

Finally, the following lemma insures that if a process X satisfies Proposition 2.1, a causal predictable ARMA process with X as innovation also satisfies Proposition 2.1. We first provide the classical following notion for a sequence $(u_n)_{n \in \mathbb{N}}$ of real numbers:

$(u_n)_{n \in \mathbb{N}}$ is an exponentially decreasing sequence (EDS)

\iff

there exists $\rho \in [0, 1[$ such as $u_n = \mathcal{O}(\rho^n)$ when $n \rightarrow \infty$.

Lemma 2.1. *Let X be a.s. a causal stationary solution of (1.1) for $\theta_0 \in \mathbb{R}^d$. Let \widetilde{X} be such as $\widetilde{X}_t = \Lambda_\beta(L) X_t$ for $t \in \mathbb{Z}$ with $\Lambda_{\beta_0}(L) = P_{\beta_0}^{-1}(L) Q_{\beta_0}(L)$ where $(P_{\beta_0}, Q_{\beta_0})$ are the coprime polynomials of a causal invertible ARMA(p, q) processes with a vector of parameters $\beta_0 \in \mathbb{R}^{p+q}$.*

Denote $\Lambda_{\beta_0}^{-1}(x) = Q_{\beta_0}^{-1}(x) P_{\beta_0}(x) = 1 + \sum_{j=1}^{\infty} \psi_j(\beta_0)x^j$. Then \tilde{X} is a.s. a causal stationary solution of the equation

$$\tilde{X}_t = \tilde{M}_{\tilde{\theta}_0}((\tilde{X}_{t-i})_{i \geq 1}) \zeta_t + \tilde{f}_{\tilde{\theta}_0}((\tilde{X}_{t-i})_{i \geq 1}) \quad \text{for } t \in \mathbb{Z},$$

where $\tilde{f}_{\tilde{\theta}_0}$ and $\tilde{M}_{\tilde{\theta}_0}$ are given in (5.1) and $\tilde{\theta}_0 = (\tilde{\theta}_0, \beta_0)$. Moreover, for $i = 0, 1, 2$ and with $K = f$ or M and $\tilde{K} = \tilde{f}$ or \tilde{M} ,

- if $\alpha_j^{(i)}(K, \{\theta_0\}) = O(j^{-\beta})$ and $\beta > 1$, then $\alpha_j^{(i)}(\tilde{K}, \{\tilde{\theta}_0\}) = O(j^{-\beta})$;
- if $\alpha_j^{(i)}(K, \{\theta_0\})$ is EDS, then $\alpha_j^{(i)}(\tilde{K}, \{\tilde{\theta}_0\})$ is EDS.

2.3 Assumptions required for the convergence of the Laplacian-QMLE

The Laplacian-QMLE could converge and be asymptotically Gaussian but this requires some additional assumptions on Θ and functions f_θ and M_θ :

- **Condition C1** (Compactness) Θ is a compact set.
- **Condition C2** (Lower bound of the conditional variance) There exists a deterministic constant $\underline{M} > 0$ such that for all $\theta \in \Theta$, then $M_\theta(x) > \underline{M}$ for all $x \in \mathbb{R}^{\mathbb{N}}$.
- **Condition C3** (Identifiability) The functions M_θ and f_θ are such that: for all $\theta_1, \theta_2 \in \Theta$, then $M_{\theta_1} = M_{\theta_2}$ and $f_{\theta_1} = f_{\theta_2}$ implies that $\theta_1 = \theta_2$.

3 Consistency and asymptotic normality of the estimator

3.1 Consistency and asymptotic normality

First we prove the strong consistency of a sequence of Laplacian-QMLE for a solution of (1.1). The proof of this theorem, is postponed in Section 5, as well as the other proofs.

Theorem 3.1. *Assume Conditions C1, C2 and C3 hold and $\theta_0 \in \Theta(r) \cap \Theta$ with $r \geq 1$. Let X be the stationary solution of (1.1). If $(A_0(f, \Theta))$ and $(A_0(M, \Theta))$ hold with*

$$\alpha_j^{(0)}(f, \Theta) + \alpha_j^{(0)}(M, \Theta) = \mathcal{O}(j^{-\ell}) \quad \text{for some } \ell > \frac{2}{\min(r, 2)} \quad (3.1)$$

then a sequence of Laplacian-QMLE $(\hat{\theta}_n)_n$ strongly converges, that is $\hat{\theta}_n \xrightarrow[n \rightarrow \infty]{a.s.} \theta_0$.

Of course, the conditions required for this strong consistency of a sequence of Laplacian-QMLE are almost the same than the ones required for the strong consistency of a sequence of Gaussian-QMLE except that $r \in [1, 2)$ is proved to be possible in Theorem 3.1 and not in case of Gaussian-QMLE (see Bardet and Wintenberger, 2009). Moreover, if $r = 2$, the condition (3.1) on Lipschitzian coefficients is weaker for Laplacian-QMLE than for Gaussian-QMLE. As we will see below, many usual time series can satisfy the assumptions of Theorem 3.1; for example, an AR(∞) process can be defined for satisfying the strong consistency of Laplacian-QMLE while the conditions given in Bardet and Wintenberger (2009) do not ensure the strong consistency of Gaussian-QMLE.

Now we state an extension of Theorem 1 proved in Davis and Dunsmuir (1997) which will be an essential step of the proof of the asymptotic normality of the estimator.

Theorem 3.2. Let $(Z_t)_{t \in \mathbb{Z}}$ be a sequence of i.i.d.r.v such as $\text{Var}(Z_0) = \sigma^2 < \infty$, with common distribution function which is symmetric ($F(-x) = 1 - F(x)$ for $x \in \mathbb{R}$) and is continuously differentiable in a neighborhood of 0 with derivative $f(0)$ in 0. Denote $\mathcal{F}_t = \sigma(Z_t, Z_{t-1}, \dots)$ for $t \in \mathbb{Z}$ and let $(Y_t)_{t \in \mathbb{Z}}$ and $(V_t)_{t \in \mathbb{Z}}$ two stationary processes adapted to $(\mathcal{F}_t)_t$ and such as $\mathbb{E}[Y_0^2 V_0^2] < \infty$.

Then

$$\sum_{t=1}^n V_{t-1} (|Z_t - n^{-1/2} Y_{t-1}| - |Z_t|) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \mathcal{N}\left(f(0) \mathbb{E}[V_0 Y_0^2], \mathbb{E}[V_0^2 Y_0^2]\right) \quad (3.2)$$

Then, the asymptotic normality of the Laplacian-QMLE can be established using additional assumptions:

Theorem 3.3. Assume that $\theta_0 \in \overset{\circ}{\Theta} \cap \Theta(r)$ where $r \geq 2$ and $\overset{\circ}{\Theta}$ denotes the interior of Θ . Let X be the stationary solution of the equation (1.1). Assume that the conditions of Theorem 3.1 hold and for $i = 1, 2$, assume $(A_i(f, \Theta))$ and $(A_i(M, \Theta))$ hold. Then, if the cumulative probability function of ζ_0 is continuously differentiable in a neighborhood of 0 with derivative $g(0)$ in 0 and if matrix Γ_F or Γ_M , defined in (5.21), are definite positive symmetric matrix, then

$$\sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \mathcal{N}_d\left(0, (\Gamma_M + 2g(0) \Gamma_F)^{-1} ((\sigma_\zeta^2 - 1) \Gamma_M + \Gamma_F) (\Gamma_M + 2g(0) \Gamma_F)^{-1}\right). \quad (3.3)$$

As it was already proved for the median estimator (see van der Vaart, 1996) or for least absolute deviations estimator of ARMA process (see Davis and Dunsmuir, 1997), it is not surprising that the probability density function g of the white noise $(\zeta_i)_i$ impacts the asymptotic covariance of (3.3). However, when $f_\theta = 0$, this is not such the case and this is what happens for GARCH processes (see Francq *et al.*, 2011, or Francq and Zakoian, 2015) where g does not appear in the asymptotic covariance.

3.2 Examples

In this section, several examples of time series satisfying the conditions of previous results are considered. Like it could be boring to state the results for all sufficiently famous processes, we refer, *mutatis mutandis*, to Bardet and Wintenberger (2009) and Bardet *et al.* (2012) for ARCH(∞) and TARARCH(∞).

1/ APARCH processes. APARCH(δ, p, q) model has been introduced (see Ding *et al.*, 1993) as the solution of equations

$$\begin{cases} X_t = \sigma_t \zeta_t, \\ \sigma_t^\delta = \omega + \sum_{i=1}^p \alpha_i (|X_{t-i}| - \gamma_i X_{t-i})^\delta + \sum_{j=1}^q \beta_j \sigma_{t-j}^\delta, \end{cases} \quad (3.4)$$

where $\delta \geq 1$, $\omega > 0$, $-1 < \gamma_i < 1$ and $\alpha_i \geq 0$ for $i = 1, \dots, p$, $\beta_j \geq 0$ for $j = 1, \dots, q$ with α_p, β_q strictly positive and $\sum_{j=1}^q \beta_j < 1$. Hence, we denote here $\theta = (\delta, \omega, \alpha_1, \dots, \alpha_p, \gamma_1, \dots, \gamma_p, \beta_1, \dots, \beta_q)$. Using L the usual backward operator such as $LX_t = X_{t-1}$, $(1 - \sum_{j=1}^q \beta_j L^j)^{-1}$ exists and simple computations imply for $t \in \mathbb{Z}$:

$$\begin{aligned} \sigma_t^\delta &= \left(1 - \sum_{j=1}^q \beta_j L^j\right)^{-1} \left[\omega + \sum_{i=1}^p \alpha_i (1 - \gamma_i)^\delta (\max(X_{t-i}, 0))^\delta + \alpha_i (1 + \gamma_i)^\delta (-\min(X_{t-i}, 0))^\delta\right] \\ &= b_0 + \sum_{i \geq 1} b_i^+ (\max(X_{t-i}, 0))^\delta + \sum_{i \geq 1} b_i^- (\max(-X_{t-i}, 0))^\delta. \end{aligned}$$

where $b_0 = w(1 - \sum_{j=1}^q \beta_j)^{-1}$ and the coefficients $(b_i^+, b_i^-)_{i \geq 1}$ are defined by the recursion relations

$$\begin{cases} b_i^+ = \sum_{k=1}^q \beta_k b_{i-k}^+ + \alpha_i(1 - \gamma_i)^\delta & \text{with } \alpha_i(1 - \gamma_i) = 0 \text{ for } i > p \\ b_i^- = \sum_{k=1}^q \beta_k b_{i-k}^- + \alpha_i(1 + \gamma_i)^\delta & \text{with } \alpha_i(1 + \gamma_i) = 0 \text{ for } i > p \end{cases} \quad (3.5)$$

with $b_i^+ = b_i^- = 0$ for $i \leq 0$. As a consequence, for APARCH model, $f_\theta^t \equiv 0$ and $M_\theta^t = \sigma_t$. It is clear that $\alpha_j^{(0)}(f, \Theta) = 0$ and simple computations imply $\alpha_j^{(0)}(M, \Theta) = \sup_{\theta \in \Theta} \max(|b_j^+(\theta)|^{1/\delta}, |b_j^-(\theta)|^{1/\delta})$. Therefore $A_0(f, \Theta)$ holds and $\sum_{j=1}^q \beta_j < 1$ implies that a sequence defined by $u_n = \sum_{k=1}^q \beta_k u_{n-k}$ for n large enough is such as $(u_n)_{n \in \mathbb{N}}$ is an exponentially decreasing sequence and therefore $A_0(M, \Theta)$ holds. Thus for $r \geq 1$, the stationarity set $\Theta(r)$ is defined by

$$\Theta(r) = \left\{ \theta \in \mathbb{R}^{2p+q+2} \mid (\mathbb{E}[|\zeta_0|^r])^{1/r} \sum_{j=1}^{\infty} \max(|b_j^+|^{1/\delta}, |b_j^-|^{1/\delta}) < 1 \right\}. \quad (3.6)$$

Now the strong consistency and asymptotic normality of the Laplacian-QMLE for APARCH models can be established (see the proof in Section 5):

Proposition 3.1. *Assume that X is a stationary solution of (3.4) with $\theta_0 \in \Theta$ where Θ is a compact subset of $\Theta(r)$ defined in (3.6). Then,*

1. *If $r = 1$, then $\hat{\theta}_n \xrightarrow[n \rightarrow \infty]{a.s.} \theta_0$.*

2. *If $r = 2$, and if Γ_M defined in (5.21) is a definite positive symmetric matrix, then*

$$\sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \mathcal{N}_{2p+q+2}(0, (\sigma_\zeta^2 - 1) \Gamma_M^{-1}).$$

To our knowledge, this is the first statement the asymptotic properties of Laplacian-QMLE for APARCH processes.

2/ ARMA-GARCH processes. ARMA(p, q)-GARCH(p', q') processes have been introduced by Ding *et al.* (1993) and Ling and McAleer (2003) as the solution of the system of equations

$$\begin{cases} P_\theta(L) X_t = Q_\theta(L) \varepsilon_t, \\ \varepsilon_t = \sigma_t \zeta_t, \quad \text{with } \sigma_t^2 = c_0 + \sum_{i=1}^{p'} c_i \varepsilon_{t-i}^2 + \sum_{i=1}^{q'} d_i \sigma_{t-i}^2 \end{cases} \quad (3.7)$$

where

- $c_0 > 0$, $c_i \geq 0$ for $i = 1, \dots, p'$, $d_i \geq 0$ for $i = 1, \dots, q'$, $\sum_{i=1}^{q'} d_i < 1$ and $c_{p'}, d_{q'}$ positive;
- $P_\theta(x) = 1 - a_1 x - \dots - a_p x^p$ and $Q_\theta(x) = 1 - b_1 x - \dots - b_q x^q$ are coprime polynomials with $\sum_{i=1}^p |a_i| < 1$ and $\sum_{i=1}^q |b_i| < 1$.

Let $\theta = (c_0, c_1, \dots, c_{p'}, d_1, \dots, d_{q'}, a_1, \dots, a_p, b_1, \dots, b_q)$. We are going to use Lemma 2.1. Since (ε_t) is supposed to be a GARCH(p', q'), then $f_\theta^\varepsilon = 0$ and $M_\theta^\varepsilon = ((1 - \sum_{j=1}^{q'} d_j L^j)^{-1} (c_0 + c_1 \varepsilon_{t-1}^2 + \dots + c_{p'} \varepsilon_{t-p'}^2))^{1/2}$ and direct computations imply that the Lipschitz coefficients of (ε_t) are such as $\alpha_j^{(0)}(f^\varepsilon, \{\theta_0\}) = 0$ and $\alpha_j^{(0)}(M^\varepsilon, \{\theta_0\}) = |\beta_j|$ with $(1 + \sum_{j=1}^{\infty} \beta_j x^j)(1 - \sum_{j=1}^{q'} d_j x^j) = \sum_{j=0}^{p'} c_j x^j$. Therefore $(\alpha_j^{(0)}(f^\varepsilon, \{\theta_0\}))_j$ and $(\alpha_j^{(0)}(M^\varepsilon, \{\theta_0\}))_j$ are EDS (see for instance Berkes and Horvath,

2004). Thus $(A_0(f^\varepsilon, \{\theta_0\}))$ and $(A_0(M^\varepsilon, \{\theta_0\}))$ hold.

Considering the ARMA part and denoting (ψ_j) such as $(1 + \sum_{j=1}^{\infty} \psi_j x^j)(1 - \sum_{j=1}^{\infty} a_j x^j) = (1 - \sum_{j=1}^{\infty} b_j x^j)$, then from Lemma 2.1 we deduce that:

$$\begin{cases} \alpha_j^{(0)}(f, \{\theta_0\}) &= |\psi_j| \\ \alpha_j^{(0)}(M, \{\theta_0\}) &\leq \sum_{k=1}^j |\psi_k| \times |\beta_{j-k}| \end{cases}.$$

Then we deduce that $(\alpha_j^{(0)}(f, \{\theta_0\}))_j$ and $(\alpha_j^{(0)}(M, \{\theta_0\}))_j$ are EDS, $(A_0(f, \{\theta_0\}))$ and $(A_0(M, \{\theta_0\}))$ hold, and X is a.s. a solution of (1.1) for θ included in the r -order stationarity set $\Theta(r)$ defined by

$$\Theta(r) = \left\{ \theta \in \mathbb{R}^{p+q+p'+q'+1} \mid \sum_{i=1}^{\infty} |\psi_i(\theta)| + (\mathbb{E}[|\zeta_0|^r])^{1/r} \sum_{j=1}^{\infty} \sum_{k=1}^j |\psi_k| \times |\beta_{j-k}| < 1 \right\}. \quad (3.8)$$

Now the strong consistency and asymptotic normality of the Laplacian-QMLE for ARMA-GARCH processes can be established:

Proposition 3.2. *Assume that X is a stationary solution of (3.7) with $\theta_0 \in \Theta$ where Θ is a compact subset of $\Theta(r)$ defined in (3.8). Then,*

1. *If $r = 1$, then $\hat{\theta}_n \xrightarrow[n \rightarrow \infty]{a.s.} \theta_0$.*
2. *If $r = 2$, and if Γ_f and Γ_M defined in (5.21) are definite positive symmetric matrix, then the asymptotic normality (3.3) of $\hat{\theta}_n$ holds.*

This result is a new one and extends the previous results already obtained with Gaussian-QMLE for such processes (see for instance, Ling and McAleer, 2003, and Bardet and Wintenberger, 2009).

3/ ARMA-ARCH(∞) processes. ARMA(p, q)-ARCH(∞) processes are a natural extension of ARMA-GARCH processes. They are the solution of the system of equations

$$\begin{cases} P_\theta(L) X_t = Q_\theta(L) \varepsilon_t, \\ \varepsilon_t = \sigma_t \zeta_t, \quad \text{with} \quad \sigma_t^2 = c_0 + \sum_{i=1}^{\infty} c_i \varepsilon_{t-i}^2 \end{cases} \quad (3.9)$$

where

- $c_0 > 0$, $c_i \geq 0$ for $i \geq 1$;
- $P_\theta(x) = 1 - a_1 x - \dots - a_p x^p$ and $Q_\theta(x) = 1 - b_1 x - \dots - b_q x^q$ are coprime polynomials with $\sum_{i=1}^p |a_i| < 1$ and $\sum_{i=1}^q |b_i| < 1$.

ARCH(∞) processes were introduced by Robinson (1991) and the asymptotic properties of Gaussian-QMLE were studied in Robinson and Zaffaroni (2006), Straumann and Mikosch (2006) or Bardet and Wintenberger (2009). Hence, we assume that there exists $\beta = (\beta_1, \dots, \beta_m)$ such as for all $i \in \mathbb{N}$, $c_i = c(i, \beta)$, with $c(\cdot)$ a known function. Let $\theta = (\beta, a_1, \dots, a_p, b_1, \dots, b_q)$. We are going to use Lemma 2.1. Since (ε_t) is supposed to be an ARCH(∞), then $f_\theta^\varepsilon = 0$ and $M_\theta^\varepsilon = (c(0, \beta) + \sum_{i=1}^{\infty} c(i, \beta) \varepsilon_{t-i}^2)^{1/2}$ and direct computations imply that the Lipschitz coefficients

of (ε_t) are such as $\alpha_j^{(0)}(f^\varepsilon, \{\theta_0\}) = 0$ and $\alpha_j^{(0)}(M^\varepsilon, \{\theta_0\}) = c(j, \beta_0)$. Therefore we assume that there exists $\ell > 1$ such as

$$c(j, \beta_0) = \mathcal{O}(j^{-\ell}) \text{ when } j \rightarrow \infty. \quad (3.10)$$

Thus $(A_0(f^\varepsilon, \{\theta_0\}))$ and $(A_0(M^\varepsilon, \{\theta_0\}))$ hold.

Considering the ARMA part and denoting (ψ_j) such as $(1 + \sum_{j=1}^{\infty} \psi_j x^j)(1 - \sum_{j=1}^{\infty} a_j x^j) = (1 - \sum_{j=1}^{\infty} b_j x^j)$, then from Lemma 2.1 we deduce that:

$$\begin{cases} \alpha_j^{(0)}(f, \{\theta_0\}) &= |\psi_j| \\ \alpha_j^{(0)}(M, \{\theta_0\}) &\leq \sum_{k=1}^j |\psi_k| \times c(j, \beta_0) \end{cases}.$$

Then we deduce that $(\alpha_j^{(0)}(f, \{\theta_0\}))_j$ is EDS and $(\alpha_j^{(0)}(M, \{\theta_0\}))_j = \mathcal{O}(j^{-\ell})$. Then $(A_0(f, \{\theta_0\}))$ and $(A_0(M, \{\theta_0\}))$ hold, and X is a.s. a solution of (1.1) for θ included in the r -order stationarity set $\Theta(r)$ defined by

$$\Theta(r) = \left\{ \theta \in \mathbb{R}^{p+q+m} \mid \sum_{i=1}^{\infty} |\psi_i(\theta)| + (\mathbb{E}[|\zeta_0|^r])^{1/r} \sum_{j=1}^{\infty} \sum_{k=1}^j |\psi_k| \times c(j, \beta_0) < 1 \right\}. \quad (3.11)$$

Now the strong consistency and asymptotic normality of the Laplacian-QMLE for ARMA-ARCH(∞) processes can be established:

Proposition 3.3. *Assume that X is a stationary solution of (3.9) where (3.10) holds and with $\theta_0 \in \Theta$ where Θ is a compact subset of $\Theta(r)$ defined in (3.11). Then,*

1. *If $r \geq 1$ and $\ell \geq 2/\min(r, 2)$, then $\widehat{\theta}_n \xrightarrow[n \rightarrow \infty]{a.s.} \theta_0$.*
2. *If $r = 2$, $\ell > 1$ and if $\partial_\beta^i c(j, \beta) = \mathcal{O}(j^{-\ell})$ for $i = 1, 2$, and if Γ_f and Γ_M defined in (5.21) are definite positive symmetric matrix, then the asymptotic normality (3.3) of $\widehat{\theta}_n$ holds.*

This result is a new one. Note that $\ell > 1$ and $r = 2$ is required for the asymptotic normality of Laplacian-QMLE while $r = 4$ and $\ell > 2$ is required for Gaussian-QMLE for such processes (see for instance Bardet and Wintenberger, 2009). This confers a clear advantage to Laplacian-QMLE.

4/ ARMA-APARCH processes. The ARMA(p, q)-APARCH(p', q') processes have been also introduced by Ding *et al.* (1993) as the solutions of the equations

$$\begin{cases} P_\theta(L) X_t = Q_\theta(L) \varepsilon_t, \\ \varepsilon_t = \sigma_t \zeta_t, \text{ with } \sigma_t^\delta = \omega + \sum_{i=1}^{p'} \alpha_i (|\varepsilon_{t-i}| - \gamma_i \varepsilon_{t-i})^\delta + \sum_{j=1}^{q'} \beta_j \sigma_{t-j}^\delta \end{cases} \quad (3.12)$$

where:

- $\delta \geq 1$, $\omega > 0$, $-1 < \gamma_i < 1$ and $\alpha_i \geq 0$ for $i = 1, \dots, p'-1$, $\beta_j \geq 0$ for $j = 1, \dots, q'-1$, $\alpha_{p'}$, $\beta_{q'}$ positive real numbers and $\sum_{j=1}^{p'} \alpha_j < 1$;
- $P_\theta(x) = 1 - a_1 x - \dots - a_p x^p$ and $\Psi_\theta(x) = 1 - b_1 x - \dots - b_q x^q$ are coprime polynomials with $\sum_{i=1}^p |a_i| < 1$ and $\sum_{i=1}^q |b_i| < 1$.

Let $\theta = (\delta, \omega, \alpha_1, \dots, \alpha_{p'}, \gamma_1, \dots, \gamma_{p'}, \beta_1, \dots, \beta_{q'}, a_1, \dots, a_p, b_1, \dots, b_q)$. Then, as for ARMA-GARCH processes, we are going to use Lemma 2.1. Thanks to the computations realized for APARCH processes, we obtain $\alpha_j^{(0)}(f^\varepsilon, \{\theta_0\}) = 0$ and $\alpha_j^{(0)}(M^\varepsilon, \{\theta_0\}) = \max(|b_j^+|^{1/\delta}, |b_j^-|^{1/\delta})$ with $(b_i^+, b_i^-)_{i \geq 1}$ defined in (3.5).

Then, we have

$$\begin{cases} \alpha_j^{(0)}(f, \{\theta_0\}) & \leq |\psi_j| \\ \alpha_j^{(0)}(M, \{\theta_0\}) & \leq \sum_{k=1}^j |\psi_k| \times \max(|b_{j-k}^+|^{1/\delta}, |b_{j-k}^-|^{1/\delta}) \end{cases} .$$

(ψ_j) such as $(1 + \sum_{j=1}^{\infty} \psi_j x^j)(1 - \sum_{j=1}^{\infty} a_j x^j) = (1 - \sum_{j=1}^{\infty} b_j x^j)$. From Lemma 2.1, $(A_0(f, \Theta))$ and $(A_0(M, \Theta))$ hold since $(\alpha_j^{(0)}(f^\varepsilon, \{\theta_0\}))_j = 0$ and $(\alpha_j^{(0)}(M^\varepsilon, \{\theta_0\}))_j$ are EDS. As a consequence, for $r \geq 1$, the stationarity set $\Theta(r)$ is defined by

$$\Theta(r) = \left\{ \theta \in \mathbb{R}^{p+q+p'+q'+2} \mid \sum_{j=1}^{\infty} |\psi_j| + (\mathbb{E}|\zeta_0|^r)^{1/r} \sum_{j=1}^{\infty} \sum_{k=1}^j |\psi_k| \times \max(|b_{j-k}^+|^{1/\delta}, |b_{j-k}^-|^{1/\delta}) < 1 \right\}.$$

Now, we are able to provide the asymptotic properties of QMLE for ARMA-APARCH models.

Proposition 3.4. *Assume that X is a stationary solution of (3.12) with $\theta_0 \in \Theta$ where Θ is a compact subset of $\Theta(r)$ defined in (3.8). Then,*

1. *If $r = 1$, then $\hat{\theta}_n \xrightarrow[n \rightarrow \infty]{a.s.} \theta_0$.*
2. *If $r = 2$, and if Γ_f and Γ_M defined in (5.21) are definite positive symmetric matrix, then the asymptotic normality (3.3) of $\hat{\theta}_n$ holds.*

This result is stated for the first time for Laplacian-QMLE. The case of Gaussian-QMLE for ARMA-APARCH could be also obtained following the previous decomposition and the paper Bardet and Wintenberger (2009). Once again, the asymptotic normality of Laplacian-QMLE only requires $r = 2$ while this requires $r = 4$ for Gaussian-QMLE.

4 Numerical Results

To illustrate the asymptotic results stated previously, we realized Monte-Carlo experiments on the behavior of Laplacian-QMLE (denoted $\hat{\theta}_n^{LQL}$) for several time series models, sample sizes and probability distributions. A comparison with the results obtained by Gaussian QMLE (denoted $\hat{\theta}_n^{GQL}$) is also proposed.

More precisely, the considered probability distributions of (ζ_t) are:

- Centred Gaussian distribution denoted \mathcal{N} ;
- Centred Laplacian distribution denoted \mathcal{L} ;
- Centred Uniform distribution denoted \mathcal{U} ;
- Centred Student distribution with 3 freedom degrees, denoted t_3 ;
- Normalized centred Gaussian mixture with probability distribution $0.05 * \mathcal{N}(-2, 0.16) + \mathcal{N}(0, 1) + 0.05 * \mathcal{N}(2, 0.16)$ and denoted \mathcal{M} .

All these probability distributions are normalized such as $E[|\zeta_0|] = 1$, required for Laplacian-QMLE. For using Gaussian-QMLE requiring $\sigma_\zeta^2 = 1$, it is necessary to consider the model with $M'_\theta = \frac{E[|\zeta_0|]}{\sigma_\zeta} M_\theta$ instead of M_θ .

Several models of time series satisfying (1.1) and the assumptions of Theorem 3.1 and 3.3 are considered:

- a ARMA(1, 1) process defined by $X_t = \phi X_{t-1} + \zeta_t + \theta\zeta_{t-1}$ with $\phi = 0.4$ and $\theta = 0.6$;
- a ARCH(1) process defined by $X_t = \zeta_t \sqrt{\omega + \alpha X_{t-1}^2}$ with $\omega = 0.4$ and $\alpha = 0.2$;
- a GARCH(1, 1) process defined by $X_t = \zeta_t \sigma_t$ where $\sigma_t^2 = \alpha_0 + \alpha_1 X_{t-1}^2 + \beta \sigma_{t-1}^2$ with $\alpha_0 = 0.2$, $\alpha_1 = 0.4$ and $\beta = 0.2$;
- a ARMA(1, 1)-GARCH(1, 1) process defined by $X_t = \phi X_{t-1} + \varepsilon_t + \theta \varepsilon_{t-1}$ where $\varepsilon_t = \zeta_t \sigma_t$ and $\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \beta \sigma_{t-1}^2$ with $\phi = 0.4$, $\theta = 0.6$, $\alpha_0 = 0.2$, $\alpha_1 = 0.4$ and $\beta = 0.1$;
- a ARMA(1, 1)-APARCH(1, 1) process defined by $X_t = \phi X_{t-1} + \varepsilon_t + \theta \varepsilon_{t-1}$ where $\varepsilon_t = \zeta_t \sigma_t$ and $\sigma_t^\delta = \alpha_0 + \alpha_1 (|\varepsilon_{t-1}| - \gamma \varepsilon_{t-1})^\delta + \beta \sigma_{t-1}^\delta$ and $\phi = 0.4$, $\theta = 0.6$, $\alpha_0 = 0.2$, $\alpha_1 = 0.4$, $\gamma = 0.5$, $\beta = 0.1$ and $\delta = 1.2$.

Hence we computed the root-mean-square error (RMSE) from 1000 independent replications of $\widehat{\theta}_n^{LQL}$ and $\widehat{\theta}_n^{LQL}$ for those processes and the results are presented in Table 1 and 2.

Conclusion of the numerical results: On the one hand, it is clear that the RMSE decreases as the sample size increases, which validates the theoretical results (consistency of the estimators). On the other hand, Table 1 and 2 show that the Laplacian-QMLE provides more accurate estimation than the Gaussian-QMLE for several types of noise, except of course in the case of a Gaussian distribution (even in this case the RSME of both the estimators are almost the same).

5 Proofs

Proof of Lemma 2.1. First, as X is a stationary process and the ARMA(p, q) process is causal invertible then \widetilde{X} is also a stationary process (the coefficients of Λ_{β_0} are EDS). Moreover, it is well known that $(\psi_j(\beta_0))_{j \in \mathbb{N}}$ is EDS. Then we have:

$$\begin{aligned}
\widetilde{X}_t &= \Lambda_{\beta_0}(L) \left(M_{\theta_0}((X_{t-i})_{i \geq 1}) \zeta_t + f_{\theta_0}((X_{t-i})_{i \geq 1}) \right) \\
\widetilde{X}_t + \sum_{j=1}^{\infty} \psi_j(\beta_0) \widetilde{X}_{t-j} &= M_{\theta_0}((\Lambda_{\beta_0}^{-1}(L) \widetilde{X}_{t-i})_{i \geq 1}) \zeta_t + f_{\theta_0}((\Lambda_{\beta_0}^{-1}(L) \widetilde{X}_{t-i})_{i \geq 1}) \\
\widetilde{X}_t &= \widetilde{M}_{\widetilde{\theta}_0}((\widetilde{X}_{t-i})_{i \geq 1}) \zeta_t + \widetilde{f}_{\widetilde{\theta}_0}((\widetilde{X}_{t-i})_{i \geq 1}) \\
\text{with } \begin{cases} \widetilde{M}_{\widetilde{\theta}_0}((x_{t-i})_{i \geq 1}) &= M_{\widetilde{\theta}_0}((\Lambda_{\beta_0}^{-1}(L) x_{t-i})_{i \geq 1}) \\ \widetilde{f}_{\widetilde{\theta}_0}((x_{t-i})_{i \geq 1}) &= f_{\theta_0}((\Lambda_{\beta_0}^{-1}(L) x_{t-i})_{i \geq 1}) - \sum_{j=1}^{\infty} \psi_j(\beta_0) x_{t-j} \end{cases} \quad (5.1)
\end{aligned}$$

		\mathcal{L}		\mathcal{N}		t_3		\mathcal{U}		\mathcal{M}		
		$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	
ARMA(1,1)	θ	100	0.106	0.091	0.114	0.117	0.113	0.090	0.112	0.059	0.110	0.078
		1000	0.031	0.024	0.032	0.032	0.036	0.027	0.031	0.014	0.031	0.023
		5000	0.014	0.010	0.014	0.015	0.016	0.011	0.016	0.012	0.013	0.010
	ϕ	100	0.119	0.102	0.121	0.128	0.123	0.102	0.120	0.067	0.121	0.090
		1000	0.037	0.028	0.036	0.036	0.040	0.030	0.036	0.017	0.036	0.027
		5000	0.016	0.012	0.014	0.016	0.017	0.013	0.016	0.007	0.014	0.006
ARCH(1)	ω	100	0.068	0.061	0.048	0.049	0.254	0.085	0.035	0.025	0.062	0.052
		1000	0.020	0.018	0.015	0.015	0.134	0.049	0.011	0.016	0.036	0.018
		5000	0.010	0.009	0.006	0.006	0.115	0.044	0.005	0.015	0.031	0.008
	α	100	0.161	0.155	0.141	0.142	0.979	0.418	0.102	0.064	0.484	0.423
		1000	0.063	0.058	0.043	0.043	0.852	0.169	0.029	0.033	0.157	0.133
		5000	0.016	0.014	0.012	0.012	0.378	0.109	0.013	0.031	0.087	0.062
GARCH(1,1)	α_0	100	0.112	0.105	0.095	0.100	0.211	0.126	0.081	0.047	0.134	0.114
		1000	0.036	0.032	0.028	0.028	0.098	0.058	0.023	0.018	0.066	0.051
		5000	0.016	0.014	0.012	0.012	0.055	0.043	0.010	0.015	0.040	0.023
	α_1	100	0.162	0.157	0.149	0.150	0.453	0.364	0.115	0.070	0.507	0.429
		1000	0.061	0.056	0.449	0.449	0.333	0.150	0.030	0.033	0.160	0.136
		5000	0.029	0.026	0.020	0.020	0.193	0.095	0.013	0.030	0.086	0.058
	β	100	0.225	0.209	0.188	0.190	0.499	0.429	0.163	0.105	0.483	0.390
		1000	0.060	0.055	0.051	0.051	0.285	0.174	0.044	0.022	0.170	0.169
		5000	0.027	0.024	0.022	0.022	0.180	0.075	0.019	0.009	0.072	0.075

Table 1: Root Mean Square Error of the components of $\hat{\theta}_n^{LQL}$ and $\hat{\theta}_n^{GQL}$ for ARMA(1, 1), ARCH(1) and GARCH(1, 1) processes.

		\mathcal{L}		\mathcal{N}		t_3		\mathcal{U}		\mathcal{M}		
		$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	$\hat{\theta}_n^{GQL}$	$\hat{\theta}_n^{LQL}$	
ARMA(1,1) -GARCH(1,1)	θ	100	0.120	0.097	0.107	0.107	0.121	0.098	0.097	0.067	0.123	0.087
		1000	0.035	0.024	0.028	0.028	0.048	0.029	0.024	0.015	0.035	0.026
		5000	0.016	0.010	0.012	0.012	0.023	0.012	0.015	0.011	0.011	0.007
	ϕ	100	0.135	0.109	0.117	0.119	0.141	0.116	0.110	0.077	0.132	0.102
		1000	0.044	0.030	0.033	0.033	0.063	0.035	0.029	0.023	0.053	0.046
		5000	0.020	0.014	0.015	0.015	0.029	0.015	0.013	0.012	0.019	0.014
	α_0	100	0.104	0.096	0.085	0.084	0.158	0.129	0.073	0.055	0.131	0.116
		1000	0.031	0.028	0.025	0.025	0.241	0.060	0.021	0.019	0.053	0.046
		5000	0.014	0.012	0.010	0.010	0.052	0.042	0.009	0.016	0.036	0.019
	α_1	100	0.179	0.177	0.166	0.167	0.469	0.385	0.134	0.107	0.494	0.405
		1000	0.064	0.060	0.045	0.045	0.328	0.161	0.031	0.046	0.160	0.137
		5000	0.031	0.027	0.020	0.020	0.182	0.096	0.013	0.038	0.090	0.062
	β	100	0.302	0.269	0.252	0.233	0.604	0.497	0.217	0.164	0.553	0.472
		1000	0.057	0.051	0.051	0.051	0.312	0.187	0.045	0.049	0.165	0.170
		5000	0.025	0.022	0.020	0.020	0.199	0.073	0.062	0.066	0.019	0.025
ARMA(1,1) -APARCH(1,1)	θ	100	0.110	0.086	0.096	0.101	0.112	0.090	0.097	0.068	0.125	0.091
		1000	0.029	0.021	0.023	0.024	0.031	0.021	0.022	0.014	0.033	0.024
		5000	0.013	0.008	0.010	0.010	0.014	0.009	0.010	0.006	0.015	0.011
	ϕ	100	0.138	0.114	0.121	0.126	0.128	0.107	0.111	0.086	0.146	0.107
		1000	0.040	0.027	0.032	0.032	0.041	0.028	0.029	0.026	0.043	0.030
		5000	0.018	0.012	0.012	0.012	0.020	0.012	0.013	0.014	0.019	0.013
	ω	100	0.198	0.192	0.199	0.210	0.254	0.262	0.221	0.170	0.290	0.272
		1000	0.079	0.067	0.056	0.056	0.226	0.218	0.044	0.045	0.142	0.129
		5000	0.033	0.028	0.025	0.025	0.209	0.207	0.017	0.029	0.061	0.056
	α	100	0.206	0.201	0.183	0.184	0.464	0.449	0.167	0.131	0.352	0.327
		1000	0.060	0.053	0.041	0.041	0.447	0.432	0.029	0.043	0.143	0.134
		5000	0.025	0.023	0.018	0.018	0.421	0.414	0.012	0.027	0.071	0.059
	γ	100	0.413	0.386	0.346	0.356	0.439	0.426	0.310	0.233	0.613	0.601
		1000	0.105	0.094	0.071	0.070	0.101	0.092	0.057	0.041	0.217	0.220
		5000	0.042	0.039	0.029	0.029	0.045	0.038	0.024	0.018	0.086	0.089
β	100	0.297	0.282	0.255	0.238	0.312	0.288	0.186	0.145	0.476	0.468	
	1000	0.074	0.067	0.058	0.058	0.074	0.066	0.043	0.033	0.151	0.150	
	5000	0.033	0.031	0.024	0.024	0.034	0.029	0.018	0.012	0.061	0.063	
δ	100	0.732	0.732	0.712	0.717	0.740	0.739	0.703	0.613	0.830	0.815	
	1000	0.402	0.352	0.296	0.296	0.394	0.338	0.235	0.290	0.542	0.534	
	5000	0.170	0.147	0.132	0.131	0.169	0.145	0.092	0.168	0.251	0.262	

Table 2: Root Mean Square Error of the components of $\hat{\theta}_n^{LQL}$ and $\hat{\theta}_n^{GQL}$ for ARMA(1,1)-GARCH(1,1) and ARMA(1,1)-APARCH(1,1) processes.

Finally, for $i = 0$,

$$\begin{aligned}
|\tilde{f}_{\tilde{\theta}_0}((x_{t-i})_{i \geq 1}) - \tilde{f}_{\tilde{\theta}_0}((y_{t-i})_{i \geq 1})| &\leq \sum_{j=1}^{\infty} \alpha_j^{(0)}(f, \{\theta_0\}) |(\Lambda_{\beta_0}^{-1}(L)x_{t-j-i})_{i \geq 1} - (\Lambda_{\beta_0}^{-1}(L)y_{t-j-i})_{i \geq 1}| \\
&\quad + |\psi_j(\beta_0)| |x_{t-j} - y_{t-j}| \\
&\leq \sum_{j=1}^{\infty} \alpha_j^{(0)}(f, \{\theta_0\}) \left| \sum_{k=0}^{\infty} |\psi_k(\beta_0)| |x_{t-k-j} - y_{t-k-j}| + |\psi_j(\beta_0)| |x_{t-j} - y_{t-j}| \right| \\
&\leq \sum_{j=1}^{\infty} \left(|\psi_j(\beta_0)| + \sum_{k=1}^j \alpha_k^{(0)}(f, \{\theta_0\}) |\psi_{j-k}(\beta_0)| \right) |x_{t-j} - y_{t-j}| \\
\implies \alpha_j^{(0)}(\tilde{f}, \{\tilde{\theta}_0\}) &\leq |\psi_j(\beta_0)| + \sum_{k=1}^j \alpha_k^{(0)}(f, \{\theta_0\}) |\psi_{j-k}(\beta_0)|. \tag{5.2}
\end{aligned}$$

Moreover, we also have:

$$\begin{aligned}
|\tilde{M}_{\tilde{\theta}_0}((x_{t-i})_{i \geq 1}) - \tilde{M}_{\tilde{\theta}_0}((y_{t-i})_{i \geq 1})| &\leq \sum_{j=1}^{\infty} \alpha_j^{(0)}(M, \{\theta_0\}) |(\Lambda_{\beta_0}^{-1}(L)x_{t-j-i})_{i \geq 1} - (\Lambda_{\beta_0}^{-1}(L)y_{t-j-i})_{i \geq 1}| \\
\implies \alpha_j^{(0)}(\tilde{M}, \{\tilde{\theta}_0\}) &\leq \sum_{k=1}^j \alpha_k^{(0)}(M, \{\theta_0\}) |\psi_{j-k}|. \tag{5.3}
\end{aligned}$$

The same kinds of computations could also be done by considering the first and second derivatives of \tilde{f} and \tilde{M} with respect to $\tilde{\theta}$. Note, and this is important, that the first and second derivatives of Λ_{β}^{-1} with respect to $\tilde{\theta}$ are also EDS. Finally,

- if when $j \rightarrow \infty$, $\alpha_j^{(0)}(K, \{\theta_0\}) = O(j^{-\beta})$ with $\beta > 1$ and $\psi_j = O(\rho^j)$ with $0 \leq \rho < 1$, then there exists $C > 0$ such as $\sum_{k=1}^j \alpha_k^{(0)}(K, \{\theta_0\}) |\psi_{j-k}| \leq C \sum_{k=1}^j k^{-\beta} \rho^{j-k} \sim -C(\log \rho)^{-1} j^{-\beta}$ and therefore $\alpha_j^{(0)}(\tilde{K}, \{\tilde{\theta}_0\}) = O(j^{-\beta})$.
- if when $j \rightarrow \infty$, $\alpha_j^{(0)}(K, \{\theta_0\}) = O(r^j)$ with $0 \leq r < 1$ and $\psi_j = O(\rho^j)$ with $0 \leq \rho < 1$, then there exists $C > 0$ such as $\sum_{k=1}^j \alpha_k^{(0)}(K, \{\theta_0\}) |\psi_{j-k}| \leq C \sum_{k=1}^j r^{-k} \rho^{j-k} = O(j \max(r, \rho)^j)$ and therefore $\alpha_j^{(0)}(\tilde{K}, \{\tilde{\theta}_0\})$ is EDS.

The same kind of computation can be also done for $(\alpha_j^{(i)}(\tilde{K}, \{\tilde{\theta}_0\}))_j$ since the derivatives and second-derivatives of $\Lambda_{\beta_0}^{-1}$ with respect to β and therefore to $\tilde{\theta}$ are also EDS. \square

Now we remind two lemmas already proved in Bardet and Wintenberger (2009):

Lemma 5.1. *Assume that $\theta_0 \in \Theta(r)$ for $r \geq 1$ and X is the causal stationary solution of the equation (1.1). If $(A_0(K, \Theta))$ holds (with $K = f$ or $K = M$) then $K_{\theta}^t \in L^r(\mathcal{C}(\Theta, R^m))$ and there exists $C > 0$ not depending on t such that*

$$\mathbb{E}[\|\hat{K}_{\theta}^t - K_{\theta}^t\|_{\Theta}^r] \leq C \mathbb{E}[|X_0|^r] \left(\sum_{j \geq t} \alpha_j(K, \Theta) \right)^r \quad \text{for all } t \in N^*. \tag{5.4}$$

Lemma 5.2. *Let $\mathcal{D}^{(2)}(\Theta)$ denote the Banach space of 2 times continuously differentiable functions on Θ equipped with the uniform norm*

$$\|h\|_{2, \Theta} = \|h\|_{\Theta} + \left\| \frac{\partial h}{\partial \theta} \right\|_{\Theta} + \left\| \frac{\partial^2 h}{\partial \theta \partial \theta'} \right\|_{\Theta}.$$

Let $\theta_0 \in \Theta(r)$ ($r \geq 1$) and assume that for $i = 0, 1, 2$, $(A_i(f, \Theta))$ and $(A_i(M, \Theta))$ hold. Then $f_\theta^t \in \mathbb{L}^r(\mathcal{D}^{(2)}(\Theta))$ and $M_\theta^t \in \mathbb{L}^r(\mathcal{D}^{(2)}(\Theta))$.

Now, we begin with the proofs of Theorem 3.1, 3.2 and 3.3.

Proof of Theorem 3.1. The proof of the theorem is divided into two parts and follows the same procedure than in Bardet and Wintenberger (2009). In (i), a uniform (on Θ) strong law of large numbers satisfied by $\frac{1}{n} \widehat{L}_n(\theta)$ converging to $L(\theta) := -\mathbb{E}[q_0(\theta)]$ is established. In (ii), it is proved that $L(\theta)$ admits a unique maximum in θ_0 . Those two conditions lead to the strong consistency of $\widehat{\theta}_n$ (from Jeantheau, 1998).

(i) In the same way and for the same reason in the proof of Theorem 1 of Bardet and Wintenberger (2009), the uniform strong law of large numbers satisfied by the sample mean of $(\widehat{q}_t)_{t \in N^*}$ (defined in (2.3)) is implied by establishing $\mathbb{E}[\|q_t(\theta)\|_\Theta] < \infty$. But from Lemma 5.1, for all $t \in \mathbb{Z}$,

$$\begin{aligned} |q_t(\theta)| &= |(M_\theta^t)^{-1}|X_t - f_\theta^t| + \log(M_\theta^t)| \\ &\leq \frac{|X_t - f_\theta^t(\theta)|}{\underline{M}} + |\log(\underline{M})| + M_\theta^t \\ \implies \sup_{\theta \in \Theta} |q_t(\theta)| &\leq \frac{1}{\underline{M}}(|X_t| + \|f_\theta^t(\theta)\|_\Theta) + |\log(\underline{M})| + \|M_\theta^t\|_\Theta. \end{aligned}$$

With $r \geq 1$, we have $\forall t \in \mathbb{Z}$, $\mathbb{E}[|X_t|] < \infty$ from Proposition 2.1 and $\mathbb{E}[\|f_\theta^t\|_\Theta^r + \|M_\theta^t\|_\Theta^r] < \infty$ from Lemma 5.1, implying $\mathbb{E}[\|f_\theta^t\|_\Theta + \|M_\theta^t\|_\Theta] < \infty$. As a consequence, for all $t \in \mathbb{Z}$,

$$\mathbb{E}[\|q_t(\theta)\|_\Theta] < \infty.$$

Hence, the uniform strong law of large numbers for $(q_t(\theta))$ follows:

$$\left\| \frac{L_n(\theta)}{n} - L(\theta) \right\|_\Theta \xrightarrow[n \rightarrow \infty]{a.s.} 0. \quad (5.5)$$

Now, we are going to establish $\frac{1}{n} \|\widehat{L}_n(\theta) - L_n(\theta)\|_\Theta \xrightarrow[n \rightarrow \infty]{a.s.} 0$. Indeed, for all $\theta \in \Theta$ and $t \in N^*$,

$$\begin{aligned} |\widehat{q}_t(\theta) - q_t(\theta)| &\leq |\log(\widehat{M}_\theta^t) - \log(M_\theta^t) + (\widehat{M}_\theta^t)^{-1}|X_t - \widehat{f}_\theta^t| - (M_\theta^t)^{-1}|X_t - f_\theta^t|| \\ &\leq |\log(\widehat{M}_\theta^t) - \log(M_\theta^t)| + |((\widehat{M}_\theta^t)^{-1} - (M_\theta^t)^{-1})|X_t - f_\theta^t| + (\widehat{M}_\theta^t)^{-1}|\widehat{f}_\theta^t - f_\theta^t| \\ &\leq |\widehat{M}_\theta^t - M_\theta^t| \underline{M}^{-1} + |\widehat{M}_\theta^t - M_\theta^t| \underline{M}^{-2} |X_t - f_\theta^t| + \underline{M}^{-1} |\widehat{f}_\theta^t - f_\theta^t| \\ &\leq C(1 + |X_t| + |f_\theta^t|)(|\widehat{M}_\theta^t - M_\theta^t| + |\widehat{f}_\theta^t - f_\theta^t|), \end{aligned}$$

with $C > 0$. Hence, we have:

$$\|\widehat{q}_t(\theta) - q_t(\theta)\|_\Theta \leq C(1 + |X_t| + \|f_\theta^t\|_\Theta)(\|\widehat{M}_\theta^t - M_\theta^t\|_\Theta + \|\widehat{f}_\theta^t - f_\theta^t\|_\Theta).$$

By Corollary 1 of Kounias and Weng (1969), the proof is achieved if there exists $s \in (0, 1]$ such as

$$\sum_{t \geq 1} \frac{1}{t^s} \mathbb{E}[\|q_t(\theta) - \widehat{q}_t(\theta)\|_\Theta^s] < \infty. \quad (5.6)$$

Let us prove (5.6) with $s = r/2$ when $r \in [1, 2]$.

From Cauchy-Schwarz Inequality and assumptions $A_0(f, \Theta)$ and $A_0(M, \Theta)$,

$$\mathbb{E}[\|\widehat{q}_t(\theta) - q_t(\theta)\|_{\Theta}^{r/2}] \leq C (\mathbb{E}[(1 + |X_t| + \|f_{\theta}^t\|_{\Theta})^r])^{\frac{1}{2}} (\mathbb{E}[(\|\widehat{M}_{\theta}^t - M_{\theta}^t\|_{\Theta} + \|f_{\theta}^t - f_{\theta}^t\|_{\Theta})^r])^{\frac{1}{2}}.$$

Using Lemma 5.1 and previous proved results implying $\mathbb{E}[|X_t|^r] < \infty$, $\mathbb{E}[\|f_{\theta}^t\|_{\Theta}^r + \|M_{\theta}^t\|_{\Theta}^r] < \infty$, we obtain

$$\begin{aligned} \mathbb{E}[\|\widehat{q}_t(\theta) - q_t(\theta)\|_{\Theta}^{r/2}] &\leq C \left(\sum_{j>t} \alpha_j^{(0)}(f, \Theta) + \alpha_j^{(0)}(M, \Theta) \right)^{\frac{r}{2}} \\ &\leq C t^{-\frac{(\ell-1)r}{2}}, \end{aligned}$$

where the last inequality is obtained from the condition (3.1) of Theorem 3.1.

Hence, we have

$$\sum_{t \geq 1} \frac{1}{t^{r/2}} \mathbb{E}[\|\widehat{q}_t(\theta) - q_t(\theta)\|_{\Theta}^{r/2}] \leq A \sum_{t \geq 1} t^{-r \ell/2},$$

which is finite when $r \ell > 2$. When $r \geq 2$, it is sufficient to consider the case $r = 2$. As a consequence, we obtain

$$\frac{1}{n} \sum_{t=1}^n \|\widehat{q}_t(\theta) - q_t(\theta)\|_{\Theta} \xrightarrow[n \rightarrow \infty]{a.s.} 0 \quad \frac{1}{n} \|\widehat{L}_n(\theta) - L_n(\theta)\|_{\Theta} \xrightarrow[n \rightarrow \infty]{a.s.} 0, \quad (5.7)$$

and therefore, using (5.5),

$$\frac{1}{n} \|\widehat{L}_n(\theta) - L(\theta)\|_{\Theta} \xrightarrow[n \rightarrow \infty]{a.s.} 0. \quad (5.8)$$

(ii) Now for $\theta \in \Theta$, we study

$$L(\theta) = -\mathbb{E}[q_0(\theta)].$$

which can also be consider as a Kullback-Lieber discrepancy. We have

$$\begin{aligned} L(\theta) &= -\mathbb{E}[\log(M_{\theta}^t) + (M_{\theta}^t)^{-1} |X_t - f_{\theta}^t|] \\ &= -\mathbb{E}\left[\log(M_{\theta}^t) + \frac{M_{\theta_0}^t}{M_{\theta}^t} \left| \zeta_t + \frac{f_{\theta_0}^t - f_{\theta}^t}{M_{\theta_0}^t} \right|\right]. \end{aligned}$$

Hence, using $\mathbb{E}[|\zeta_t|] = 1$, we obtain:

$$\begin{aligned} L(\theta_0) - L(\theta) &= \mathbb{E}\left[\log\left(\frac{M_{\theta}^t}{M_{\theta_0}^t}\right) + \frac{M_{\theta_0}^t}{M_{\theta}^t} \left| \zeta_t + \frac{f_{\theta_0}^t - f_{\theta}^t}{M_{\theta_0}^t} \right| - 1\right] \\ &= \mathbb{E}\left[\log\left(\frac{M_{\theta}^t}{M_{\theta_0}^t}\right) - 1 + \frac{M_{\theta_0}^t}{M_{\theta}^t} \mathbb{E}\left[\left| \zeta_t + \frac{f_{\theta_0}^t - f_{\theta}^t}{M_{\theta_0}^t} \right| \mid (X_{t-k})_{k \geq 1}\right]\right]. \end{aligned}$$

But for ζ_t following a symmetric probability distribution, for any $m \in \mathbb{R}^*$, $\mathbb{E}[|\zeta_t + m|] > \mathbb{E}[|\zeta_t|] = 1$.

Therefore, for $\theta \neq \theta_0$, if $f_{\theta} \neq f_{\theta_0}$ (else $>$ is replaced by \geq),

$$\begin{aligned} L(\theta_0) - L(\theta) &> \mathbb{E}\left[\log\left(\frac{M_{\theta}^t}{M_{\theta_0}^t}\right) - 1 + \frac{M_{\theta_0}^t}{M_{\theta}^t}\right] \\ &> h\left(\frac{M_{\theta}^t}{M_{\theta_0}^t}\right), \end{aligned}$$

with $h(x) = \log(x) - 1 + x$. But for any $x \in (0, 1) \cup (1, \infty)$, $h(x) > 0$ and $h(1) = 0$. Therefore if $M_\theta \neq M_{\theta_0}$, $h\left(\frac{M_\theta^t}{M_{\theta_0}^t}\right) > 0$ (> 0 is replaced by $= 0$ if $M_\theta = M_{\theta_0}$). This implies from Condition C3 (Identifiability) that $L(\theta_0) - L(\theta) > 0$ almost surely for all $\theta \in \Theta$, $\theta \neq \theta_0$. Hence a supremum of $L(\theta)$ is only reached for $\theta = \theta_0$ which is the unique maximum. \square

Proof of Theorem 3.2. We follow the same scheme of proof than in Davis and Dunsmuir (1997). Hence, denote

$$\begin{aligned} S_n &= \sum_{t=1}^n V_{t-1} (|Z_t - n^{-1/2} Y_{t-1}| - |Z_t|) \\ &= -n^{-1/2} \sum_{t=1}^n V_{t-1} Y_{t-1} \operatorname{sgn}(Z_t) \\ &\quad + 2 \sum_{t=1}^n V_{t-1} (n^{-1/2} Y_{t-1} - Z_t) (\mathbf{1}_{0 < Z_t < n^{-1/2} Y_{t-1}} - \mathbf{1}_{n^{-1/2} Y_{t-1} < Z_t < 0}) \\ &= A_n + B_n. \end{aligned}$$

Since $\mathbb{E}[V_{t-1} Y_{t-1} \operatorname{sgn}(Z_t) / \mathcal{F}_{t-1}] = \mathbb{E}[\operatorname{sgn}(Z_t)] \mathbb{E}[V_{t-1} Y_{t-1}] = 0$ and $\mathbb{E}[V_0^2 Y_0^2] < \infty$, we can apply a central limit theorem for stationary martingale difference sequence (see Billingsley, 1968) and

$$A_n \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \mathcal{N}(0, \mathbb{E}[V_0^2 Y_0^2]). \quad (5.9)$$

Now, considering B_n , define also $W_{nt} = V_{t-1} (n^{-1/2} Y_{t-1} - Z_t) \mathbf{1}_{0 < Z_t < n^{-1/2} Y_{t-1}}$. Using the same arguments as in Davis and Dunsmuir (1997), we also obtain

- $\limsup_{n \rightarrow \infty} n \mathbb{E}[W_{nt}^2] = 0$;
- $\mathbb{E}[W_{nt} / \mathcal{F}_{t-1}] \simeq \frac{1}{2n} f(0) V_{t-1} Y_{t-1}^2$ for $|n^{-1/2} Y_{t-1}| < \varepsilon$;
- $\sum_{t=1}^n W_{nt} \xrightarrow[n \rightarrow \infty]{\mathcal{P}} \frac{1}{2} f(0) \mathbb{E}[V_0 Y_0^2 \mathbf{1}_{Y_0 > 0}]$.

Then we deduce

$$B_n \xrightarrow[n \rightarrow \infty]{\mathcal{P}} f(0) \mathbb{E}[V_0 Y_0^2]. \quad (5.10)$$

The proof is achieved from (5.9) and (5.10). \square

Proof of Theorem 3.3. We follow a proof which is similar to the one of Theorem 2 in Davis and Dunsmuir (1997) or Li and Li (2008).

Let $v = \sqrt{n}(\theta - \theta_0) \in \mathbb{R}^d$. Then we are going to prove in 2/ that maximizing $\widehat{L}_n(\theta)$ is equivalent to maximizing $L_n(\theta)$ which is equivalent to maximizing

$$\begin{aligned} W_n(v) &= - \sum_{t=1}^n (q_t(\theta_0 + n^{-1/2} v) - q_t(\theta_0)) \\ &= \sum_{t=1}^n \log \left(\frac{(M_{\theta_0 + n^{-1/2} v}^t)^{-1}}{(M_{\theta_0}^t)^{-1}} \right) + (M_{\theta_0}^t)^{-1} |X_t - f_{\theta_0}^t| - (M_{\theta_0 + n^{-1/2} v}^t)^{-1} |X_t - f_{\theta_0 + n^{-1/2} v}^t| \end{aligned} \quad (5.11)$$

with respect to v . As a consequence, there exists a sequence $(\widehat{v}_n)_n$ where \widehat{v}_n is a maximizer of $W_n(v)$ such as $\widehat{v}_n = \sqrt{n}(\widehat{\theta}_n - \theta_0)$. In 1/ we will provide a limit theorem satisfied by $W_n(v)$. Then

we are going to prove in 3/ that $(W_n(\cdot))_n$ converges as a process of $\mathcal{C}(\mathbb{R}^d)$ (space of continuous functions on \mathbb{R}^d) to a limit process W . Hence $(\widehat{v}_n)_n$ converges to the maximizer of W .

1/ First, we are going to study the asymptotic behavior of $W_n(v)$. We have

$$\begin{aligned} W_n(v) &= \sum_{t=1}^n \log \left(\frac{(M_{\theta_0+n^{-1/2}v}^t)^{-1}}{(M_{\theta_0}^t)^{-1}} \right) + |X_t - f_{\theta_0}^t| \left((M_{\theta_0}^t)^{-1} - (M_{\theta_0+n^{-1/2}v}^t)^{-1} \right) \\ &\quad + \sum_{t=1}^n (M_{\theta_0+n^{-1/2}v}^t)^{-1} (|X_t - f_{\theta_0}^t| - |X_t - f_{\theta_0+n^{-1/2}v}^t|) \\ &= I_1(v) + I_2(v). \end{aligned}$$

We have:

$$I_1(v) = - \sum_{t=1}^n \log \left(\frac{M_{\theta_0+n^{-1/2}v}^t}{M_{\theta_0}^t} \right) + |\zeta_t| \left(1 - \frac{M_{\theta_0}^t}{M_{\theta_0+n^{-1/2}v}^t} \right)$$

Using Taylor expansions, we deduce that for each $t \in \{1, \dots, n\}$, there exists $\bar{\theta}_1^t$ and $\bar{\theta}_2^t$ in the segment $[\theta_0, \theta_0 + n^{-1/2}v]$ such as:

$$\begin{aligned} \log \left(\frac{M_{\theta_0+n^{-1/2}v}^t}{M_{\theta_0}^t} \right) &= n^{-1/2} (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial M_{\theta_0}^t}{\partial \theta} \right)_{\theta_0} + \frac{1}{2} n^{-1} \left\{ v' \left(\frac{\partial^2 M_{\theta_0}^t}{\partial \theta^2} \right)_{\bar{\theta}_1^t} - (M_{\theta_0}^t)^{-2} \left(v' \left(\frac{\partial M_{\theta_0}^t}{\partial \theta} \right)_{\bar{\theta}_1^t} \right)^2 \right\} \\ \frac{M_{\theta_0}^t}{M_{\theta_0+n^{-1/2}v}^t} &= 1 - n^{-1/2} (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial M_{\theta_0}^t}{\partial \theta} \right)_{\theta_0} + \frac{1}{2} n^{-1} \left\{ 2 (M_{\theta_0}^t)^{-2} \left(v' \left(\frac{\partial M_{\theta_0}^t}{\partial \theta} \right)_{\bar{\theta}_2^t} \right)^2 - v' \left(\frac{\partial^2 M_{\theta_0}^t}{\partial \theta^2} \right)_{\bar{\theta}_2^t} \right\} \end{aligned}$$

Then,

$$\begin{aligned} I_1(v) &= n^{-1/2} \sum_{t=1}^n (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial M_{\theta_0}^t}{\partial \theta} \right)_{\theta_0} (|\zeta_t| - 1) + \frac{1}{2n} \sum_{t=1}^n (M_{\theta_0}^t)^{-1} \left\{ v' \left(\frac{\partial^2 M_{\theta_0}^t}{\partial \theta^2} \right)_{\bar{\theta}_2^t} v |\zeta_t| - v' \left(\frac{\partial^2 M_{\theta_0}^t}{\partial \theta^2} \right)_{\bar{\theta}_1^t} v \right\} \\ &\quad + \frac{1}{2n} \sum_{t=1}^n (M_{\theta_0}^t)^{-2} \left\{ \left(v' \left(\frac{\partial M_{\theta_0}^t}{\partial \theta} \right)_{\bar{\theta}_1^t} \right)^2 - 2 \left(v' \left(\frac{\partial M_{\theta_0}^t}{\partial \theta} \right)_{\bar{\theta}_2^t} \right)^2 |\zeta_t| \right\} \\ &= I_1^{(1)}(v) + I_1^{(2)}(v) + I_1^{(3)}(v). \end{aligned}$$

Using a Central Limit Theorem for martingale-differences (see [4]), since $\mathbb{E} \left[(M_{\theta_0}^t)^{-1} v' \left(\frac{\partial M_{\theta_0}^t}{\partial \theta} \right)_{\theta_0} (|\zeta_t| - 1) / \mathcal{F}_{t-1} \right] = 0$ and $\mathbb{E} \left[\left\| (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial M_{\theta_0}^t}{\partial \theta} \right)_{\theta_0} \right\|_{\Theta}^2 \right] < \infty$ (from Lemma 5.2), we have:

$$I_1^{(1)}(v) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \mathcal{N} \left(0, \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial M_{\theta_0}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right] (\sigma_{\zeta}^2 - 1) \right). \quad (5.12)$$

Now, using that $\theta \in \Theta \mapsto \frac{\partial M_{\theta}^t}{\partial \theta}$ and $\theta \mapsto \frac{\partial^2 M_{\theta}^t}{\partial \theta^2}$ are continuous functions, $\bar{\theta}_1^t \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \theta_0$ and $\bar{\theta}_2^t \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \theta_0$, we claim that $I_1^{(2)}(v)$ have the same limit distribution that $\frac{1}{2n} \sum_{t=1}^n (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial^2 M_{\theta_0}^t}{\partial \theta^2} \right)_{\theta_0} v (|\zeta_t| - 1)$. Note that $\mathbb{E} \left[(M_{\theta_0}^t)^{-1} v' \left(\frac{\partial^2 M_{\theta_0}^t}{\partial \theta^2} \right)_{\theta_0} v (|\zeta_t| - 1) / \mathcal{F}_{t-1} \right] = 0$ and $\mathbb{E} \left[\left\| (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial^2 M_{\theta_0}^t}{\partial \theta^2} \right)_{\theta_0} v \right\| \right] < \infty$ (from Lemma 5.2). Thus, from the strong large number law for martingale-differences (see [4]), we obtain:

$$\frac{1}{2n} \sum_{t=1}^n (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial^2 M_{\theta_0}^t}{\partial \theta^2} \right)_{\theta_0} v (|\zeta_t| - 1) \xrightarrow[n \rightarrow \infty]{a.s.} 0,$$

and this implies:

$$I_1^{(2)}(v) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} 0. \quad (5.13)$$

Finally similar arguments as previously induce that $I_1^{(3)}(v)$ have the same limit distribution that $\frac{1}{2n} \sum_{t=1}^n (M_{\theta_0}^t)^{-2} \left(v' \left(\frac{\partial M_{\theta}^t}{\partial \theta} \right)_{\theta_0} \right)^2 (1 - 2|\zeta_t|)$. From the strong large number law for martingale-differences (see [4]), we obtain:

$$\begin{aligned} \frac{1}{2n} \sum_{t=1}^n (M_{\theta_0}^t)^{-2} \left(v' \left(\frac{\partial M_{\theta}^t}{\partial \theta} \right)_{\theta_0} \right)^2 (1 - 2|\zeta_t|) &\xrightarrow[n \rightarrow \infty]{a.s.} \frac{1}{2} \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial M_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 (1 - 2|\zeta_0|) \right] \\ &\xrightarrow[n \rightarrow \infty]{a.s.} -\frac{1}{2} \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial M_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right], \end{aligned}$$

and this implies:

$$I_1^{(3)}(v) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} -\frac{1}{2} \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial M_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right]. \quad (5.14)$$

Finally, from (5.12), (5.13) and (5.14), we obtain:

$$I_1(v) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \mathcal{N} \left(-\frac{1}{2} \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial M_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right], \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial M_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right] (\sigma_{\zeta}^2 - 1) \right). \quad (5.15)$$

Now we consider $I_2(v) = \sum_{t=1}^n (M_{\theta_0+n^{-1/2}v}^t)^{-1} (|X_t - f_{\theta_0}^t| - |X_t - f_{\theta_0+n^{-1/2}v}^t|)$. Using again Taylor expansion, we can write:

$$\begin{aligned} I_2(v) &= \sum_{t=1}^n (M_{\theta_0}^t)^{-1} (|X_t - f_{\theta_0}^t| - |X_t - f_{\theta_0+n^{-1/2}v}^t|) \\ &\quad - n^{-1/2} \sum_{t=1}^n (M_{\bar{\theta}_M^t}^t)^{-2} v' \left(\frac{\partial M_{\theta}^t}{\partial \theta} \right)_{\bar{\theta}_M^t} (|X_t - f_{\theta_0}^t| - |X_t - f_{\theta_0+n^{-1/2}v}^t|) \\ &= I_2^{(1)}(v) + I_2^{(2)}(v), \end{aligned}$$

with $\bar{\theta}_M^t$ in the segment $[\theta_0, \theta_0 + n^{-1/2}v]$.

First we have:

$$I_2^{(1)}(v) = \sum_{t=1}^n (|\zeta_t| - |\zeta_t - n^{-1/2}(M_{\theta_0}^t)^{-1} v' \left(\frac{\partial f_{\theta}^t}{\partial \theta} \right)_{\bar{\theta}_f^t}|)$$

with $\bar{\theta}_f^t$ in the segment $[\theta_0, \theta_0 + n^{-1/2}v]$. Using Theorem 3.2, which an extension of Theorem 1 established in Davis and Dunsmuir (1997), denoting $Z_t = \zeta_t$, $Y_t = (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial f_{\theta}^t}{\partial \theta} \right)_{\bar{\theta}_f^t}$ and $V_t = 1$ for $t \in \mathbb{Z}$,

$$I_2^{(1)} \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \mathcal{N} \left(-g(0) \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial f_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right], \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial f_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right] \right). \quad (5.16)$$

since $\mathbb{E}[Y_t^2 V_t^2] \leq \underline{M}^{-2} \mathbb{E}[\|v' \frac{\partial f_{\theta}^t}{\partial \theta}\|_{\Theta}^2] < \infty$ from Lemma 5.2.

Then, we have:

$$\begin{aligned}
I_2^{(2)}(v) &\sim n^{-1/2} \sum_{t=1}^n (M_{\theta_0}^t)^{-2} v' \left(\frac{\partial M_{\theta}^t}{\partial \theta} \right)_{\theta_0} (|X_t - f_{\theta_0}^t| - |X_t - f_{\theta_0+n^{-1/2}v}^t|) \\
&\sim n^{-1/2} \sum_{t=1}^n (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial M_{\theta}^t}{\partial \theta} \right)_{\theta_0} (|\zeta_t| - |\zeta_t - n^{-1/2} (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial f_{\theta}^t}{\partial \theta} \right)_{\theta_f^t}|) \\
&\xrightarrow[n \rightarrow \infty]{\mathcal{P}} 0,
\end{aligned} \tag{5.17}$$

using the proof of Theorem 3.2 and denoting $Z_t = \zeta_t$, $Y_t = (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial f_{\theta}^t}{\partial \theta} \right)_{\theta_f^t}$ and $V_t = (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial M_{\theta}^t}{\partial \theta} \right)_{\theta_0}$ for $t \in \mathbb{Z}$ and condition $\mathbb{E}[|V_t Y_t|] < \infty$ insuring a strong law of large number instead of central limit theorem for a martingale difference process. Therefore, from (5.16) and (5.17), we deduce

$$I_2(v) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \mathcal{N} \left(-g(0) \mathbb{E}[(M_{\theta_0}^0)^{-2} (v' \left(\frac{\partial f_{\theta}^0}{\partial \theta} \right)_{\theta_0})^2], \mathbb{E}[(M_{\theta_0}^0)^{-2} (v' \left(\frac{\partial f_{\theta}^0}{\partial \theta} \right)_{\theta_0})^2] \right). \tag{5.18}$$

Finally, we obtain the behavior of $W_n(v)$ defined in (5.11) from (5.15) and (5.18). However, we have to specify the asymptotic dependency relation between $I_1^{(1)}$ and $I_2^{(1)}$. Indeed these two terms converge to a Gaussian law. This implies to consider the asymptotic behavior of the sum of these two terms which could be reduced to the asymptotic behavior of:

$$n^{-1/2} \sum_{t=1}^n (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial M_{\theta}^t}{\partial \theta} \right)_{\theta_0} (|\zeta_t| - 1) + n^{-1/2} \sum_{t=1}^n (M_{\theta_0}^t)^{-1} v' \left(\frac{\partial f_{\theta}^t}{\partial \theta} \right)_{\theta_0} \text{sgn}(\zeta_t),$$

from the proof of Theorem 3.2. Using again a central limit theorem for martingale differences, we obtain as asymptotic variance:

$$\begin{aligned}
\mathbb{E} \left[(M_{\theta_0}^t)^{-2} \left(v' \left(\frac{\partial M_{\theta}^t}{\partial \theta} \right)_{\theta_0} (|\zeta_t| - 1) + v' \left(\frac{\partial f_{\theta}^t}{\partial \theta} \right)_{\theta_0} \text{sgn}(\zeta_t) \right)^2 \right] &= \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial M_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right] (\sigma_{\zeta}^2 - 1) \\
&+ 2 \mathbb{E} \left[(M_{\theta_0}^0)^{-2} v' \left(\frac{\partial M_{\theta}^0}{\partial \theta} \right)_{\theta_0} v' \left(\frac{\partial f_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right] \mathbb{E} [(|\zeta_t| - 1) \text{sgn}(\zeta_t)] + \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(v' \left(\frac{\partial f_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right] \\
&= \mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left\{ (\sigma_{\zeta}^2 - 1) \left(v' \left(\frac{\partial M_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 + \left(v' \left(\frac{\partial f_{\theta}^0}{\partial \theta} \right)_{\theta_0} \right)^2 \right\} \right]
\end{aligned} \tag{5.19}$$

since $(\zeta_t)_t$ admits a symmetric probability distribution with a null median and expectation. Therefore, there is no covariance term and finally we obtain:

$$W_n(v) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} W(v) = v' \left(-\frac{1}{2} \Gamma_M - g(0) \Gamma_F \right) v + v' N \tag{5.20}$$

$$\text{with } \begin{cases} N \stackrel{\mathcal{L}}{=} \mathcal{N}(0, ((\sigma_{\zeta}^2 - 1) \Gamma_M + \Gamma_F)) \\ \Gamma_F = \left(\mathbb{E} \left[(M_{\theta_0}^0)^{-2} \left(\frac{\partial f_{\theta}^0}{\partial \theta_i} \right)_{\theta_0} \left(\frac{\partial f_{\theta}^0}{\partial \theta_j} \right)_{\theta_0} \right] \right)_{1 \leq i, j \leq d} \\ \Gamma_M = \left(\mathbb{E} \left[\left(\frac{\partial \log(M_{\theta}^0)}{\partial \theta_i} \right)_{\theta_0} \left(\frac{\partial \log(M_{\theta}^0)}{\partial \theta_j} \right)_{\theta_0} \right] \right)_{1 \leq i, j \leq d} \end{cases} \tag{5.21}$$

2/ Now, we consider the approximation $\widehat{W}_n(v)$ of $W_n(v)$ defined by:

$$\widehat{W}_n(v) = - \sum_{t=1}^n (\widehat{q}_t(\theta_0 + n^{-1/2}v) - \widehat{q}_t(\theta_0)) \quad \text{for any } v \in \mathbb{R}^d.$$

From the assumptions of Theorem 3.1 and (5.7) we have $\frac{1}{n} \sum_{t=1}^n \|\widehat{q}_t(\theta) - q_t(\theta)\|_{\Theta} \xrightarrow[n \rightarrow \infty]{a.s.} 0$. Then we have $\widehat{W}_n(v) = W_n(v) + R_n(v)$ with $[\sup_{v \in \mathbb{R}^d} |R_n(v)|] \leq 2 \sum_{t=1}^n [\|\widehat{q}_t(\theta) - q_t(\theta)\|_{\Theta}] \xrightarrow[n \rightarrow \infty]{a.s.} 0$ and then:

$$\widehat{W}_n(v) \xrightarrow[n \rightarrow \infty]{\mathcal{D}} W(v) \quad (5.22)$$

with W defined in (5.20).

3/ Now, from (5.22), the proof of Theorem 3.2 and the same arguments than in the proof of Theorem 2 of Davis and Dunsmuir (1997), we deduce that finite distributions $(\widehat{W}_n(v_1), \dots, \widehat{W}_n(v_k))$ converge to $(W(v_1), \dots, W(v_k))$ for any $(v_1, \dots, v_k) \in (\mathbb{R}^d)^k$. Moreover, always following the proof of Theorem 2, $(W_n(v))_v$ converges to $(W(v))_v$ as a process on the continuous function space \mathcal{C}^0 . As a consequence, a maximum \widehat{v} of $\widehat{W}_n(v)$ satisfies:

$$\widehat{v} = (\Gamma_M + 2g(0)\Gamma_F)^{-1}N,$$

and this implies (3.3). \square

Proof of Proposition 3.1. First, Condition **C2** is satisfied since $b_0 > 0$. Other conditions on Lipschitzian coefficients are also satisfied from Lemma 2.1 (see the arguments above). The identifiability condition **C3** is also satisfied from the following which are divided into two parts. In (i) we proof that $(\delta, b_0, (b_i^+(\theta), b_i^-(\theta))_{i \geq 1})$ (defined in (3.5)) are unique, thereafter in (ii) we proof that $\theta = (\omega, (\alpha_i)_{1 \leq i \leq p}, (\gamma_i)_{1 \leq i \leq p}, (\beta_i)_{1 \leq i \leq q})$ is also unique.

(i) The proof of this result follow the same reasoning in Berkes *et al.* (2003). First we have

$$\sigma_t^\delta = b_0(\theta) + \sum_{i \geq 1} b_i^+(\theta)(\max(X_{t-i}, 0))^\delta + \sum_{i \geq 1} b_i^-(\theta)(\max(-X_{t-i}, 0))^\delta. \quad (5.23)$$

We prove the result by contradiction. Suppose that there exist two vectors $\beta = (\delta, b_0, (b_i^+)_{i \geq 1}, (b_i^-)_{i \geq 1})$ and $\beta' = (\delta', b'_0, (b_i^{+'})_{i \geq 1}, (b_i^{-'})_{i \geq 1})$ verifying (5.23). Let $m > 0$ be the smallest integer satisfying $b_m^+ \neq b_m^{+'}$ or $b_m^- \neq b_m^{-'}$ (if $b_i^+ = b_i^{+'}$ and $b_i^- = b_i^{-'}$ $\forall i \geq 1$ then $b_0 = b'_0$). In one hand, since $x \in (0, \infty) \mapsto x^\delta$ is a one-to-one map and since $P(X_t = \pm 1, \forall t \in \mathbb{Z}) = 0$, we have $\delta = \delta'$. In the other hand, by definition of m , we have

$$\begin{aligned} (b_m^{+'} - b_m^+)(\max(X_{t-i}, 0))^\delta + (b_m^{-'} - b_m^-)(\max(-X_{t-m}, 0))^\delta \\ = b_0 - b'_0 + \sum_{i \geq m+1} (b_i^+ - b_i^{+'})(\max(X_{t-i}, 0))^\delta + \sum_{i \geq m+1} (b_i^- - b_i^{-'})(\max(-X_{t-i}, 0))^\delta. \end{aligned} \quad (5.24)$$

From (3.4), we have $X_{t-m} = \sigma_{t-m}\zeta_{t-m}$, therefore

$$(b_m^{+'} - b_m^+)(\max(X_{t-m}, 0))^\delta + (b_m^{-'} - b_m^-)(\max(-X_{t-m}, 0))^\delta = \begin{cases} (b_m^{+'} - b_m^+)\sigma_{t-m}^\delta \zeta_{t-m}^\delta & \text{when } \zeta_{t-m} \geq 0 \\ (b_m^{-'} - b_m^-)\sigma_{t-m}^\delta (-\zeta_{t-m})^\delta & \text{when } \zeta_{t-m} < 0 \end{cases}$$

Moreover (5.24) and the fact that $b_m^+ \neq b_m^{+'}$ or $b_m^- \neq b_m^{-'}$ implies that at least one of the following equalities hold

$$\begin{cases} \zeta_{t-m}^\delta = ((b_m^{+'} - b_m^+)\sigma_{t-m}^\delta)^{-1} \left(\sum_{i \geq m+1} (b_i^+ - b_i^{+'})(\max(X_{t-i}, 0))^\delta \right) & \text{when } \zeta_{t-m} \geq 0 \\ \text{or} \\ (-\zeta_{t-m})^\delta = ((b_m^{-'} - b_m^-)\sigma_{t-m}^\delta)^{-1} \left(\sum_{i \geq m+1} (b_i^- - b_i^{-'})(\max(X_{t-i}, 0))^\delta \right) & \text{when } \zeta_{t-m} < 0 \end{cases}$$

Since $\sigma_{t-m}^\delta > b_0 > 0$, ζ_{t-m}^δ is well defined. Let F_k be the F -algebra generated by $(\zeta_i, i < k)$. The causal representation of the APARCH(δ, p, q) shows that X_j is F_j -measurable and thus the right-hand side of the above equations (and consequently also ζ_{t-m}^δ in the case $\zeta_{t-m} \geq 0$ or the case $\zeta_{t-m} < 0$) is a real-valued random variable, measurable with respect to F_{t-m-1} . Since (ζ_j) is a sequence of independent random variables, this implies that ζ_{t-m} is a.s. constant when $\zeta_{t-m} \geq 0$ or when $\zeta_{t-m} < 0$, contradicting the hypothesis saying ζ_0^δ has a non-degenerate distribution. This achieves (i).

(ii) The representation (5.23) is the same as

$$\sigma_t^\delta = b_0 + \Psi^+(L)(\max(X_t, 0))^\delta + \Psi^-(L)(\max(-X_t, 0))^\delta.$$

with $\Psi^+ = \Upsilon_{\theta_1}^{-1} \Delta_{\theta_2}^+$, $\Psi^- = \Upsilon_{\theta_1}^{-1} \Delta_{\theta_2}^-$ and $\Delta_{\theta_2}^+(L) = \sum_{i=1}^p \alpha_i (1 - \gamma_i) L^i$, $\Delta_{\theta_2}^-(L) = \sum_{i=1}^p \alpha_i (1 + \gamma_i) L^i$ and $\Upsilon_{\theta_1}(L) = \sum_{i=1}^q \beta_i L^i$, where $(\Delta_{\theta_2}^+, \Upsilon_{\theta_1})$ and $(\Delta_{\theta_2}^-, \Upsilon_{\theta_1})$ respectively coprime and $\theta_1 = (\beta_i)_{1 \leq i \leq q}$, $\theta_2 = ((\alpha_i)_{1 \leq i \leq p}, (\gamma_i)_{1 \leq i \leq p})$, then $\theta = (\omega, \theta_1, \theta_2)$.

Suppose that there exist others polynomials $\Delta_{\theta_2}^+ = \sum_{i=1}^p \alpha'_i (1 - \gamma'_i) L^i$, $\Delta_{\theta_2}^- = \sum_{i=1}^p \alpha'_i (1 + \gamma'_i) L^i$, $\Upsilon_{\theta_2} = \sum_{i=1}^q \beta'_i L^i$ satisfying $\Psi^+ = \Upsilon_{\theta_1}^{-1} \Delta_{\theta_2}^+$, $\Psi^- = \Upsilon_{\theta_1}^{-1} \Delta_{\theta_2}^-$ with $(\Delta_{\theta_2}^+, \Upsilon_{\theta_1})$, $(\Delta_{\theta_2}^-, \Upsilon_{\theta_1})$ respectively coprime. Then

$$\begin{cases} \Upsilon_{\theta_1}^{-1} \Delta_{\theta_2}^+ = \Upsilon_{\theta_1}^{-1} \Delta_{\theta_2}^+ \\ \Upsilon_{\theta_1}^{-1} \Delta_{\theta_2}^- = \Upsilon_{\theta_1}^{-1} \Delta_{\theta_2}^- \end{cases} \Rightarrow \begin{cases} \Delta_{\theta_2}^+ = (\Upsilon_{\theta_1} \Upsilon_{\theta_1}^{-1}) \Delta_{\theta_2}^+ \\ \Delta_{\theta_2}^- = (\Upsilon_{\theta_1} \Upsilon_{\theta_1}^{-1}) \Delta_{\theta_2}^- \end{cases}$$

from the first equality, since $\deg(\Delta_{\theta_2}^+) = \deg(\Delta_{\theta_2}^+) = q$, we conclude that $\Upsilon_{\theta_1} \Upsilon_{\theta_1}^{-1} = 1$, therefore $\Upsilon_{\theta_1} = \Upsilon_{\theta_1}$ and so $\Delta_{\theta_2}^+ = \Delta_{\theta_2}^+$, likewise from the second equality we conclude that $\Delta_{\theta_2}^- = \Delta_{\theta_2}^-$.

- The equalities $\Delta_{\theta_2}^+ = \Delta_{\theta_2}^+$, $\Delta_{\theta_2}^- = \Delta_{\theta_2}^-$ implies that $\alpha_i (1 - \gamma_i) = \alpha'_i (1 - \gamma'_i)$ and $\alpha_i (1 + \gamma_i) = \alpha'_i (1 + \gamma'_i)$ which give $\alpha_i = \alpha'_i$ and $\gamma_i = \gamma'_i$.
- The equality $\Upsilon_{\theta_1} = \Upsilon_{\theta_1}$ implies that $\beta_i = \beta'_i$.
- Since $(\beta_i)_{i=1,p}, b_0 = w(1 - \sum_{j=1}^p \beta_j)^{-1}$ are unique then ω is unique.

Thus, Condition **C3** is established and the proof of proposition is achieved. \square

Proof of Proposition 3.2. Since we prove that Lemma 2.1 implies that conditions on Liptshitzian coefficients $(\alpha_j^{(i)}(f, \Theta))_j$ and $(\alpha_j^{(i)}(M, \Theta))_j$, it remains to prove conditions **C2** and **C3**. Condition **C2** holds since c_0 is supposed to be a positive number. Finally, condition **C3** also holds since $f_\theta = f_{\theta'}$ implies $\psi_j(\theta) = \psi_j(\theta')$ for all $j \in \mathbb{Z}$. Therefore the parameters of the ARMA part of the process are identified and then the identification of the parameters GARCH can be deduced from the proof of Proposition 3.1. \square

Proof of Proposition 3.4. This proofs mimics exactly the proof of Proposition 3.2. \square

References

- [1] Bardet, J.-M. and Wintenberger, O. (2009) Asymptotic normality of the Quasi-Maximum likelihood estimator for multidimensional causal process, *The Annals of Statistics*, 37, 2730-2759.

- [2] Berkes, I., Horváth, L. and Kokoszka, P. (2003), GARCH processes: structure and estimation, *Bernoulli*, 9, 201-227.
- [3] Berkes, I. and Horváth, L. (2004) The efficiency of the estimators of the parameters in GARCH processes, *The Annals of Statistics*, 32, 633-655.
- [4] Billingsley, P. (1968) *Convergence of Probability Measures*. Wiley, New York.
- [5] Bollerslev, T. (1986) Generalized autoregressive conditional heteroskedasticity. *J. Econometrics*, 31,307-327.
- [6] Ding, Z., Granger C.W.J. and Engle R.F. (1993); A Long Memory Property of Stock Market Returns and a New Model, *Journal of Empirical Finance* 1, 83-106.
- [7] Engle, R.F. (1982) Autoregressive conditional heteroskedasticity with estimates of the variance of U.K. inflation, *Econometrica* 50, 987-1008.
- [8] Francq, C. and Zakoian, J.-M. (2004) Maximum likelihood estimation of pure GARCH and ARMA-GARCH processes, *Bernoulli*, 10, 605-637.
- [9] Francq, C., Lepage, G. and Zakoian, J-M. (2011) Two-stage non Gaussian QML estimation of GARCH models and testing the efficiency of the Gaussian QMLE. *Journal of Econometrics*, 165, 246-257.
- [10] Francq, C. and Zakoian, J-M. (2013) Optimal predictions of powers of conditionally heteroskedastic processes. *Journal of the Royal Statistical Society B*, 75, 345-367.
- [11] Francq, C. and Zakoian, J-M. (2015) Risk-parameter estimation in volatility models. *Journal of Econometrics*, 184, 158-173.
- [12] Jeantheau, T. (1998). Strong consistency of estimators for multivariate arch models. *Econometric Theory* ,14, 70-86.
- [13] Kounias, E.G. and Weng, T.-S. (1969) An inequality and almost sure convergence. *Annals of Mathematical Statistics*, 40, 1091-1093.
- [14] Li, G. and Li, W.K. (2008) Least Absolute Deviation Estimation for Fractionally Integrated Autoregressive Moving Average Time Series Models with Conditional Heteroscedasticity. *Biometrika*, 95, 399-414.
- [15] Ling, S. and McAleer, M. (2003). Asymptotic theory for a vector ARMA-GARCH model. *Econometric Theory*, 19, 280-310.
- [16] Lumsdaine, R.L. (1996) Consistency and asymptotic normality of the quasi-maximum likelihood in IGARCH(1,1) and covariance stationary GARCH(1,1) models. *Econometrica*, 64, 575-596.
- [17] Newey, W.K. and Steigerwald, D.G. (1997). Asymptotic bias for quasi-maximum-likelihood estimators in conditional heteroskedasticity models. *Econometrica* 65, 587-599.

- [18] Peng, L. and Yao, Q. (2003). Least absolute deviations estimation for ARCH and GARCH models. *Biometrika* 90, 967-975.
- [19] Robinson, P.M. (1991). Testing for strong serial correlation and dynamic conditional heteroskedasticity in multiple regression, *Journal of Econometrics*, 47, 67-84.
- [20] Robinson, P.M. and Zaffaroni, P. (2006) Pseudo-maximum likelihood estimation of ARCH(∞) models, *Ann. Statist.* 34, 1049-1074.
- [21] Straumann, D. and Mikosch, T. (2006). Quasi-maximum-likelihood estimation in conditionally heteroscedastic time series: A stochastic recurrence equations approach. *Ann. Statist.* 34, 2449-2495.
- [22] van der Vaart, A.W. (2000) *Asymptotic Statistics*, Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press.
- [23] Weiss, A.A. (1986) Asymptotic theory for ARCH models estimation and testing. *Econometric Theory*, 2, 107-131.