

# HOPF AUTOMORPHISMS AND TWISTED EXTENSIONS

SUSAN MONTGOMERY, MARIA D. VEGA, AND SARAH WITHERSPOON

ABSTRACT. We give some applications of a Hopf algebra constructed from a group acting on another Hopf algebra  $A$  as Hopf automorphisms, namely Molnar’s smash coproduct Hopf algebra. We find connections between the exponent and Frobenius-Schur indicators of a smash coproduct and the twisted exponents and twisted Frobenius-Schur indicators of the original Hopf algebra  $A$ . We study the category of modules of the smash coproduct.

## 1. INTRODUCTION

Molnar [Ml] defined smash coproducts of Hopf algebras, putting them on equal footing with the better-known smash products by viewing both as generalizations of semidirect products of groups. Recently smash coproducts have made an appearance as examples of new phenomena in representation theory [BW, DE]. In this paper we propose several applications of smash coproducts. In particular, the smash coproduct construction will allow us to “untwist” some invariants defined via the action of a Hopf algebra automorphism, such as the twisted exponents and the twisted Frobenius-Schur indicators.

We note that considering Hopf automorphisms is a timely topic, since there has been recent progress in determining the automorphism groups of some Hopf algebras [AD, Ke, R3, SV, Y]. There has also been much recent work on indicators; their importance lies in the fact that they are invariants of the category of representations of the Hopf algebra, and may be defined for more abstract categories [NSc]. Moreover the notion of twisted indicators can be extended to pivotal categories [SV3].

We start by defining the smash coproduct  $A \bowtie k^G$ , for any Hopf algebra  $A$  with an action of a finite group  $G$  by Hopf automorphisms, in the next section. In Section 3 we recall the notions of exponent and twisted exponent [SV2] of a Hopf algebra, and find connections between the exponent of  $A \bowtie k^G$  and twisted exponents of  $A$  itself. In Section 4 we assume the Hopf algebra  $A$  is semisimple. We recall definitions of Frobenius-Schur indicators [KSZ] and twisted Frobenius-Schur indicators [SV] for simple modules over the Hopf algebra, and give relationships between the indicators of the smash coproduct  $A \bowtie k^G$  and twisted indicators of  $A$  itself.

In Section 5 we do not assume the Hopf algebra is semisimple. We introduce the twisted Frobenius-Schur indicators of the regular representation of such a Hopf algebra, simultaneously generalizing indicators for not necessarily semisimple Hopf algebras [KMN] and twisted indicators for semisimple Hopf algebras [SV]. Again we find a connection with the Frobenius-Schur indicator of a smash coproduct. We compute an example for which the Hopf algebra

---

*Date:* June 5, 2015.

The first author was supported by NSF grant DMS-1301860, the second author by an Alliance Post-Doctoral Fellowship supported by NSF grant DMS-0946431, and the third author by NSF grants DMS-1101399 and DMS-1401016.

$A$  is of dimension 8 in Section 6. Finally in Section 7 we study the structure of categories of modules of  $A \bowtie k^G$ , showing that they are equivalent to semidirect product tensor categories  $\mathcal{C} \rtimes G$ , where  $\mathcal{C}$  is a category of  $A$ -modules.

Throughout,  $k$  will be an algebraically closed field of characteristic 0.

## 2. THE SMASH COPRODUCT

Our Hopf algebra was defined by Molnar [Ml, Theorem 2.14], who called it the smash coproduct, although our definition seems different at first glance. See also [R2, p. 357].

Let  $A$  be a Hopf algebra over a field  $k$  and let a finite group  $G$  act as Hopf algebra automorphisms of  $A$ . Let  $k^G$  be the algebra of set functions from  $G$  to  $k$  under pointwise multiplication; that is, if  $\{p_x \mid x \in G\}$  denotes the basis of  $k^G$  dual to  $G$ , then  $p_x p_y = \delta_{x,y} p_x$  for all  $x, y \in G$ . Recall that  $k^G$  is a Hopf algebra with comultiplication given by  $\Delta(p_x) = \sum_{y \in G} p_y \otimes p_{y^{-1}x}$ , counit  $\varepsilon(p_x) = \delta_{1,x}$  and antipode  $S(p_x) = p_{x^{-1}}$  for all  $x \in G$ .

Then we may form the *smash coproduct Hopf algebra*

$$K = A \bowtie k^G$$

with algebra structure the usual tensor product of algebras. Denote by  $a \bowtie p_x$  the element  $a \otimes p_x$  in  $K$ , for each  $a \in A$  and  $x \in G$ . Comultiplication is given by

$$\Delta(a \bowtie p_x) = \sum_{y \in G} (a_1 \bowtie p_y) \otimes ((y^{-1} \cdot a_2) \bowtie p_{y^{-1}x})$$

for all  $x \in G$ ,  $a \in A$ . The counit and antipode are determined by

$$\varepsilon(a \bowtie p_x) = \delta_{1,x} \varepsilon(a) 1 \quad \text{and} \quad S(a \bowtie p_x) = (x^{-1} \cdot S(a)) \bowtie p_{x^{-1}}.$$

If  $\Lambda_A$  is an integral for  $A$ , then  $\Lambda_K = \Lambda_A \bowtie p_1$  is an integral for  $K$ .

Note that Molnar defines the smash coproduct for the right coaction of any commutative Hopf algebra  $H$ . We show that our construction is actually his smash coproduct with  $H = k^G$ , by dualizing our  $G$ -action to a  $k^G$ -coaction.

**Lemma 2.1.** (1)  $K$  as above is isomorphic to the smash coproduct as in [Ml, Theorem 2.14], and thus is a Hopf algebra.

(2) If  $A$  is finite-dimensional, then  $K^* \cong A^* \# kG$ , the smash product Hopf algebra as in [Ml, Theorem 2.13].

*Proof.* (1) Given the left action of  $G$  on  $A$ , we define  $\rho : A \rightarrow A \otimes k^G$  by  $a \mapsto \sum_{x \in G} (x \cdot a) \otimes p_x$ . Then  $\rho$  is a right comodule map, using the fact that the  $G$ -action on  $A$  satisfies  $x \cdot (y \cdot a) = (xy \cdot a)$  and  $1 \cdot a = a$  for all  $x, y \in G$  and  $a \in A$ .

Next we note that  $A$  is a right comodule algebra under  $\rho$  since the  $G$ -action is multiplicative, that is  $(x \cdot a)(x \cdot b) = x \cdot (ab)$ . Also  $A$  is a right comodule coalgebra, as the  $G$ -action preserves the coalgebra structure of  $A$ , that is,  $x \cdot (\sum_a a_1 \otimes a_2) = \sum_a (x \cdot a)_1 \otimes (x \cdot a)_2$ . Thus  $A$  is a right  $k^G$ -comodule bialgebra.

Finally the antipode also dualizes to the antipode given by Molnar, and thus Molnar's theorem [Ml, Theorem 2.14] applies.

(2) This is a special case of Molnar's result [Ml, Theorem 5.4]. □

### 3. HOPF POWERS AND EXPONENTS

In any Hopf algebra  $H$ , we denote the  $n$ th Hopf power of an element  $x \in H$  by  $x^{[n]} = \sum_x x_1 x_2 x_3 \dots x_n$ ; that is, first apply  $\Delta_H$   $n - 1$  times to  $x$  and then multiply. Note that  $x \mapsto x^{[n]}$  is a linear map.

For  $H$  semisimple, recall that the exponent of  $H$ ,  $\exp(H)$ , is the smallest positive integer  $n$ , if it exists, such that  $x^{[n]} = \varepsilon(x)1$  for all  $x \in H$ . More generally, this definition makes sense whenever  $S^2 = id$ . We assume this property of  $S$  unless stated otherwise.

Recently [SV2] introduced the *twisted exponent*, where  $\exp$  is twisted by an automorphism of  $H$  of finite order. Assume that  $\tau \in \text{Aut}(H)$  and that  $n$  is a multiple of the order of  $\tau$ . Define the  $n$ th  $\tau$ -twisted Hopf power of  $x$  to be

$$x^{[n,\tau]} := \sum_x x_1(\tau \cdot x_2)(\tau^2 \cdot x_3) \dots (\tau^{n-1} \cdot x_n).$$

**Definition 3.1.**  $\exp_\tau(H)$  is the smallest positive integer  $n$ , if it exists, such that  $n$  is a multiple of the order of  $\tau$  and  $x^{[n,\tau]} = \varepsilon(x)1$  for all  $x \in H$ .

Since  $\tau$  is a Hopf automorphism,  $\varepsilon(\tau \cdot x) = \varepsilon(x)$  for any  $x \in H$ , and thus  $\varepsilon(x^{[n,\tau]}) = \varepsilon(x^{[n]}) = \varepsilon(x)$ . If  $H$  is not semisimple and  $S^2 \neq id$  yet  $S$  is still bijective, there is a more general definition of the twisted exponent in [SV2].

We will need the following proposition which is a special case of [SV2, Proposition 3.4].

**Proposition 3.2.** *Suppose that the Hopf automorphism  $\tau$  of the semisimple Hopf algebra  $H$  has order  $r$ ,  $\exp_\tau(H)$  is finite, and  $m$  is a positive integer. Then  $x^{[mr,\tau]} = \varepsilon(x)1$  for all  $x \in H$  if and only if  $\exp_\tau(H)$  divides  $m$ .*

Next we give some formulas for our Hopf algebras  $K = A \bowtie k^G$ .

**Lemma 3.3.** *Let  $w = a \bowtie p_x \in A \bowtie k^G$ , the smash coproduct as above. Then*

$$(a \bowtie p_x)^{[n]} = \sum_{z \in G, z^n = x} a^{[n, z^{-1}]} \bowtie p_z.$$

*In particular for  $w = \Lambda_K = \Lambda_A \bowtie p_1$ , replace  $z$  by  $z^{-1}$ . Then*

$$\Lambda_K^{[n]} = \sum_{z \in G, z^n = 1} \Lambda_A^{[n, z]} \bowtie p_{z^{-1}}.$$

*Proof.* A calculation shows that

$$(a \bowtie p_x)^{[n]} = \sum_{z \in G, z^n = x} a_1(z^{-1} \cdot a_2)(z^{-2} \cdot a_3) \dots (z^{-(n-1)} \cdot a_n) \bowtie p_z,$$

which gives the first equation in the lemma. The second follows from the first. □

We now find a relation among the (twisted) exponents of  $A$ ,  $G$ , and  $K = A \bowtie k^G$ .

**Theorem 3.4.** *Assume that  $S^2 = id$  in  $A$ . Then the exponent of  $K$  is the least common multiple of  $\exp(G)$  and  $\exp_z(A)$  for all  $z \in G$ .*

*Proof.* Let  $n = \exp(K)$ , so that

$$(a \bowtie p_x)^{[n]} = \varepsilon(a \bowtie p_x)1 = \varepsilon(a)\delta_{x,1}1 = \varepsilon(a)\delta_{x,1} \sum_z p_z$$

for all  $a \in A$  and  $x \in G$ . When  $a = 1$ , then  $(p_x)^{[n]} = \delta_{x,1}1$  implies that  $\exp(G) = \exp(k^G)$  divides  $n$ . Thus  $z^n = 1$  for all  $z \in G$ . By the above calculation,  $(a \natural p_1)^{[n]} = \varepsilon(a)1$ , and so by Lemma 3.3,  $a^{[n, z^{-1}]} = \varepsilon(a)$  for all  $z \in G$  and  $a \in A$ . Therefore by Proposition 3.2,  $\exp(K)$  is a common multiple of  $\exp(G)$  and  $\exp_z(A)$  for all  $z \in G$ .

Now let  $m$  be any common multiple of  $\exp(G)$  and  $\exp_z(A)$  for all  $z \in G$ . By Lemma 3.3 and Proposition 3.2,

$$\begin{aligned} (a \natural p_x)^{[m]} &= \sum_{z \in G, z^m = x} a^{[m, z^{-1}]} \natural p_z \\ &= \delta_{1,x} \sum_{z \in G} a^{[m, z^{-1}]} \natural p_z \\ &= \delta_{1,x} \varepsilon(a) \sum_{z \in G} p_z = \varepsilon(a \natural p_x) 1_K. \end{aligned}$$

Again by Proposition 3.2,  $\exp(K)$  divides  $m$ . □

We will use the following lemma in calculations.

**Lemma 3.5.** *Let  $H$  be a Hopf algebra and let  $\tau$  be a Hopf automorphism of  $H$  whose order divides  $n$ . Then  $S(x^{[n, \tau]}) = \tau^{-1} \cdot (S(x)^{[n, \tau^{-1}]})$  for all  $x \in H$ .*

*Proof.* Since  $S$  is an anti-algebra and anti-coalgebra map and  $\tau^n = 1$  by hypothesis,

$$\begin{aligned} S(x^{[n, \tau]}) &= S \left( \sum_x x_1(\tau \cdot x_2)(\tau^2 \cdot x_3) \cdots (\tau^{n-1} \cdot x_n) \right) \\ &= \sum_x (\tau^{n-1} \cdot S(x_n))(\tau^{n-2} \cdot S(x_{n-1})) \cdots (\tau^2 \cdot S(x_3))(\tau \cdot S(x_2))S(x_1) \\ &= \sum_x (\tau^{-1} \cdot S(x_n))(\tau^{-2} \cdot S(x_{n-1})) \cdots (\tau^{-1(n-2)} \cdot S(x_3))(\tau^{-1(n-1)} \cdot S(x_2))S(x_1) \\ &= \tau^{-1} \cdot \left( \sum_x S(x_n)(\tau^{-1} \cdot S(x_{n-1})) \cdots (\tau^{-1(n-3)} \cdot S(x_3))(\tau^{-1(n-2)} \cdot S(x_2))(\tau^{-1(n-1)} \cdot S(x_1)) \right) \\ &= \tau^{-1} \cdot (S(x)^{[n, \tau^{-1}]}). \end{aligned}$$

□

**Corollary 3.6.** *Let  $H$  be a Hopf algebra for which  $S^2 = id$  and let  $\tau$  be a Hopf automorphism of  $H$ . Then  $\exp_{\tau^{-1}}(H) = \exp_{\tau}(H)$ .*

*Proof.* It is clear from Lemma 3.5 that  $x^{[n, \tau]} = \varepsilon(x)1 \iff S(x)^{[n, \tau^{-1}]} = \varepsilon(x)1$  since  $\tau$  and  $S$  are bijective. Thus the two twisted exponents are the same. □

**Question 3.7.** We ask if Corollary 3.6 is true more generally. That is, if the order of  $\tau$  is  $n$  and  $m$  is relatively prime to  $n$ , then is  $\exp_{\tau^m}(H) = \exp_{\tau}(H)$ ?

#### 4. MODULES AND FROBENIUS-SCHUR INDICATORS

In this section, we assume  $A$  is a semisimple Hopf algebra, and thus we may assume that  $\Lambda_A$  is a normalized integral, that is,  $\varepsilon(\Lambda_A) = 1$ . Then the integral  $\Lambda_K = \Lambda_A \natural p_1$  of  $K = A \natural k^G$  is a normalized integral of  $K$ .

For any (left)  $K$ -module  $M$ , we may write

$$M = \bigoplus_{x \in G} M_x$$

where  $M_x = p_x \cdot M$  is a  $K$ -submodule of  $M$  for each  $x \in G$ . Note that each  $M_x$  is also an  $A$ -module, by restricting the action to  $A$ .

Let  $\nu_m^K$  denote the  $m$ th *Frobenius-Schur indicator* for  $K$ -modules as in [KSZ], and let  $\nu_{m,x}^A$  denote the  $m$ th *twisted Frobenius-Schur indicator* for  $A$ -modules, twisted by  $x$ , as in [SV]. That is, if  $V$  is a  $K$ -module with character (or trace function)  $\chi_V$ , then

$$\nu_m^K(V) = \chi_V(\Lambda_K^{[m]}).$$

If  $W$  is an  $A$ -module with character  $\chi_W$  and  $x$  is an automorphism of  $A$  whose order divides  $m$ , then

$$\nu_{m,x}^A(W) = \chi_W(\Lambda_A^{[m,x]}).$$

See [SV] for general results on twisted indicators and for computations of  $\nu_{m,x}^A$  when  $A = H_8$ , the smallest semisimple noncommutative, noncocommutative Hopf algebra.

Our next theorem gives a relationship between the Frobenius-Schur indicators of  $K$  and the twisted Frobenius-Schur indicators of  $A$ .

**Theorem 4.1.** *For every  $K$ -module  $M$ ,*

$$\nu_m^K(M) = \sum_{x \in G, x^m=1} \nu_{m,x^{-1}}^A(M_x).$$

*Proof.* Write  $M = \bigoplus_{x \in G} M_x$  as before. Then  $\nu_m^K(M) = \sum_{x \in G} \nu_m^K(M_x)$ , and we will now compute  $\nu_m^K(M_x)$  for an element  $x$  of  $G$ , writing  $\Lambda = \Lambda_A$  for ease of notation: By Lemma 3.3,

$$\begin{aligned} \nu_m^K(M_x) &= \chi_{M_x}(\Lambda_K^{[m]}) \\ &= \chi_{M_x} \left( \sum_{z \in G, z^m=1} \Lambda^{[m,z]} \natural p_{z^{-1}} \right) \\ &= \delta_{x^m,1} \chi_{M_x}(\Lambda^{[m,x^{-1}]}) = \delta_{x^m,1} \nu_{m,x^{-1}}^A(M_x). \end{aligned}$$

Summing over all elements of  $G$ , we obtain the stated formula. □

As a consequence, for example, if  $x$  is an element of  $G$  of order  $n$  and  $M$  is a  $K$ -module for which  $M = M_x$  (i.e.  $M_y = 0$  for all  $y \neq x$ ), then  $\nu_m^K(M) = 0$  for all  $m < n$ .

In our next result, we show that a twisted Frobenius-Schur indicator may always be realized as a Frobenius-Schur indicator for a smash coproduct. Let  $\tau$  be any Hopf automorphism of  $A$  of finite order  $n$ , and let  $G = \langle \tau \rangle$  be the cyclic subgroup of the automorphism group generated by  $\tau$ . Set  $K = A \natural k^G$ .

**Theorem 4.2.** *For any  $A$ -module  $N$ , extend  $N$  to be a  $K$ -module  $M$  by letting  $M_{\tau^{-1}} = N$  and  $M_x = 0$  for all  $x \in G, x \neq \tau^{-1}$ . Then for every positive integer multiple  $m$  of  $n$ ,*

$$\nu_{m,\tau}^A(N) = \nu_m^K(M).$$

Thus every value of a twisted indicator for  $A$  is the value of an ordinary indicator for a smash coproduct over  $A$ .

*Proof.* By Theorem 4.1,

$$\nu_m^K(M) = \sum_{x \in G, x^m=1} \nu_{m,x^{-1}}^A(M_x) = \nu_{m,\tau}^A(M_{\tau^{-1}}) = \nu_{m,\tau}^A(N).$$

□

**Example 4.3.** We illustrate the theorem using a non-trivial automorphism of  $A = H_8$ , the Kac-Palyutkin algebra of dimension 8 which is neither commutative nor cocommutative. The Hopf automorphism group was found in [SV], Section 4.2. Let  $A$  be generated by  $x, y, z$  with the usual relations  $x^2 = y^2 = 1$ ,  $z^2 = \frac{1}{2}(1 + x + y - xy)$ ,  $xy = yx$ ,  $xz = zy$  and  $yz = zx$ , where  $x, y$  are group-like and  $\Delta(z) = \frac{1}{2}(1 \otimes 1 + 1 \otimes x + y \otimes 1 - y \otimes x)(z \otimes z)$ .

Let  $\tau = \tau_4$  be the automorphism of  $A$  of order 2 that interchanges  $x$  and  $y$  and sends  $z$  to  $\frac{1}{2}(-z + xz + yz + xyz)$ , and let  $\chi$  be the character of the unique two-dimensional simple module  $N$  of  $A$ . Then from [SV],  $\nu_{2,\tau}^A(N) = -1$ .

Letting  $G = \langle \tau \rangle$  and  $K = A \bowtie k^G$ ,  $N$  becomes a  $K$ -module  $M$  by setting  $M_\tau = N$  and  $M_1 = 0$ . Then  $\nu_2^K(M) = -1$ .

## 5. FROBENIUS-SCHUR INDICATORS FOR NON-SEMISIMPLE HOPF ALGEBRAS

Let  $A$  be a finite-dimensional Hopf algebra that is not necessarily semisimple and for which  $S^2$  is not necessarily the identity map. When  $A$  is not semisimple, there does not exist a normalized integral, and so we cannot use the definition of indicator from the previous section. Instead we extend the work in [KMN] and define twisted Frobenius-Schur indicators for  $A$  itself and obtain connections to Frobenius-Schur indicators of smash coproducts. Fix  $\tau$ , a Hopf automorphism of  $A$  whose order divides the positive integer  $m$ . We define a variant of the  $m$ th twisted Hopf power map of  $A$  to be  $P_{m-1,\tau} : A \rightarrow A$ , given by

$$P_{m-1,\tau}(a) = \sum_a (\tau^{m-1} \cdot a_1)(\tau^{m-2} \cdot a_2) \cdots (\tau^2 \cdot a_{m-2})(\tau \cdot a_{m-1})$$

for all  $a \in A$ . We will use this map to define twisted Frobenius-Schur indicators, and then we will show how it relates to the twisted Hopf power maps defined in Section 3, by giving equivalent definitions of twisted Frobenius-Schur indicators in Theorem 5.1 and Corollary 5.2.

The  $m$ th twisted Frobenius-Schur indicator of  $A$  is

$$\nu_{m,\tau}(A) := \text{Tr}(S \circ P_{m-1,\tau}),$$

the trace of the map  $S \circ P_{m-1,\tau}$  from  $A$  to  $A$ , where  $S$  is the antipode of  $A$ .

We choose this definition as it specializes to the definition of the Frobenius-Schur indicator of the regular representation  $A$  for an arbitrary finite-dimensional Hopf algebra in [KMN] when  $\tau$  is the identity, and also to the definition of twisted Frobenius-Schur indicators in the semisimple case given in [SV, Theorem 5.1]. The indicator of the regular representation has also been considered in [Sh].

The following theorem generalizes part of [KMN, Theorem 2.2].

**Theorem 5.1.** *Let  $\Lambda$  be a left integral of  $A$  and  $\lambda$  a right integral of  $A^*$  for which  $\lambda(\Lambda) = 1$ . Then*

$$\nu_{m,\tau}(A) = \lambda(S(\Lambda)^{[m,\tau]}).$$

*Proof.* By [R, Theorem 1],

$$\begin{aligned}
\text{Tr}(S \circ P_{m-1, \tau}) &= \sum \lambda(S(\Lambda_2) S \circ P_{m-1, \tau}(\Lambda_1)) \\
&= \sum \lambda(S(\Lambda_m) S((\tau^{m-1} \cdot \Lambda_1)(\tau^{m-2} \cdot \Lambda_2) \cdots (\tau \cdot \Lambda_{m-1}))) \\
&= \sum \lambda(S(\Lambda_m)(\tau \cdot S(\Lambda_{m-1})) \cdots (\tau^{m-1} \cdot S(\Lambda_1))) \\
&= \sum \lambda(S(\Lambda)_1(\tau \cdot S(\Lambda)_2) \cdots (\tau^{m-1} \cdot S(\Lambda)_m)) = \lambda(S(\Lambda)^{[m, \tau]}).
\end{aligned}$$

□

A similar proof to that of [KMN, Corollary 2.6] yields the following result that will be useful for computations.

**Corollary 5.2.** *Let  $\Lambda_r$  be a right integral of  $A$  and  $\lambda_r$  be a right integral of  $A^*$  for which  $\lambda_r(\Lambda_r) = 1$ . Then*

$$\nu_{m, \tau}(A) = \lambda_r(\Lambda_r^{[m, \tau]}).$$

*Similarly let  $\Lambda_l$  be a left integral of  $A$  and  $\lambda_l$  be a left integral of  $A^*$  for which  $\lambda_l(\Lambda_l) = 1$ . Then*

$$\nu_{m, \tau}(A) = \lambda_l(\tau^{-1} \cdot \Lambda_l^{[m, \tau^{-1}]}).$$

*Proof.* The first statement follows immediately from Theorem 5.1 and the fact that if  $\Lambda_l$  is a left integral, then  $\Lambda_r := S(\Lambda_l)$  is a right integral, and the value of  $\lambda_r$  on each is the same.

For the second statement, if  $\lambda_r$  is a right integral, let  $\lambda_l := \lambda_r \circ S$ , a left integral of  $A^*$ . Then again by Theorem 5.1 and also Lemma 3.5,

$$\begin{aligned}
\lambda_l(\tau^{-1} \cdot \Lambda_l^{[m, \tau^{-1}]}) &= \lambda_r(S(\tau^{-1} \cdot \Lambda_l^{[m, \tau^{-1}]})) \\
&= \lambda_r(\tau^{-1} \cdot (S(\Lambda_l^{[m, \tau^{-1}]})) \\
&= \lambda_r(S(\Lambda_l)^{[m, \tau]}) = \lambda_r(\Lambda_r^{[m, \tau]}).
\end{aligned}$$

□

Now let  $G$  be a group of Hopf algebra automorphisms of  $A$ , as in Section 2. The next result is a connection between twisted indicators of  $A$  and indicators of the smash coproduct  $K = A \bowtie k^G$ .

**Theorem 5.3.**  $\nu_m(K) = \sum_{g \in G, g^m=1} \nu_{m, g}(A).$

*Proof.* Note that  $\Lambda_K = \Lambda \natural p_1$  and  $\lambda_{K^*} = \lambda \otimes (\sum_{z \in G} z)$  (since e.g.  $\varepsilon(z \cdot a) = \varepsilon(a)$ ). By [KMN, Theorem 2.2] and our Lemmas 3.3 and 3.5,

$$\begin{aligned}
\nu_m(K) &= \lambda_{K^*}(S_K(\Lambda_K^{[m]})) \\
&= \left( \lambda \otimes \left( \sum_{z \in G} z \right) \right) \left( S_K \left( \sum_{g \in G, g^m=1} \Lambda^{[m,g]} \otimes p_{g^{-1}} \right) \right) \\
&= \left( \lambda \otimes \left( \sum_{z \in G} z \right) \right) \left( S_K \left( \sum_{g \in G, g^m=1} \Lambda_1(g \cdot \Lambda_2) \cdots (g^{m-1} \cdot \Lambda_m) \right) \otimes p_{g^{-1}} \right) \\
&= \sum_{g \in G, g^m=1} \lambda(g \cdot S(\Lambda_1(g \cdot \Lambda_2) \cdots (g^{m-1} \cdot \Lambda_m))) \\
&= \sum_{g \in G, g^m=1} \lambda(S(\Lambda)^{[m, g^{-1}]}) \\
&= \sum_{g \in G, g^m=1} \nu_{m, g^{-1}}(A) = \sum_{g \in G, g^m=1} \nu_{m, g}(A).
\end{aligned}$$

□

In the next section, we compute an example, a non-semisimple Hopf algebra of dimension 8 and its Hopf automorphism group.

## 6. A NON-SEMISIMPLE EXAMPLE

Let  $A$  be the Hopf algebra defined as

$$A = k\langle g, x, y \mid gx = -xg, gy = -yg, xy = -yx, g^2 = 1, x^2 = y^2 = 0 \rangle$$

with coalgebra structure given by:

$$\begin{aligned}
\Delta(g) &= g \otimes g, \quad \varepsilon(g) = 1, \quad S(g) = g, \\
\Delta(x) &= x \otimes g + 1 \otimes x, \quad \varepsilon(x) = 0, \quad S(x) = gx, \\
\Delta(y) &= y \otimes g + 1 \otimes y, \quad \varepsilon(y) = 0, \quad S(y) = gy.
\end{aligned}$$

The element  $\Lambda = xy + yx$  is both a right and left integral for  $A$ , and  $\lambda = (xy)^*$  is both a right and left integral for  $A^*$  such that  $\lambda(\Lambda) = 1$ .

**Lemma 6.1.** *Let  $V$  be the  $k$ -span of  $x$  and  $y$ . Then  $\text{Aut}(A) \cong \text{Gl}_2(V)$ .*

*Proof.* This is close to the examples considered in [AD], as  $A$  is pointed and generated by its group-like and skew-primitive elements. However we provide an elementary proof for completeness.

The coradical of  $A$  is given by  $A_0 = k\langle g \rangle$ . Any automorphism  $\tau$  of  $A$  stabilizes  $A_0$  and so fixes  $g$ . The next term of the coradical filtration is

$$A_1 = A_0 \oplus V \oplus gV,$$

since  $V$  is the set of  $(g, 1)$ -primitives and  $gV$  is the set of  $(1, g)$ -primitives. Consequently  $V$  and  $gV$  are each stable under the action of  $\tau$ . But the  $\tau$ -action on  $V$  determines the  $\tau$ -action on  $gV$ , and also on  $A = A_1 \oplus W$ , where  $W$  is the span of  $xy$  and  $gxy$ .

Conversely it is easy to check that any invertible linear action on  $V$  preserves all of the relations of  $A$ , and thus gives an automorphism. □



For an automorphism  $\tau$  of order 2 or 3, we are able to compute some values of the indicators, using Corollary 5.2. We identify  $\tau$  with a matrix

$$\begin{pmatrix} x \\ y \end{pmatrix} \xrightarrow{\tau} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix},$$

where  $a, b, c, d \in k$ , such that  $\text{Det}(\tau) = ad - bc \neq 0$ .

**Proposition 6.2.** *Case (1). If  $\tau^2 = 1$  and  $m$  is even, then  $\nu_{m,\tau}(A) = \frac{m^2}{2}(1 + \text{Det}(\tau))$ . Consequently,*

$$\nu_{m,\tau}(A) = \begin{cases} m^2, & \text{if } \text{Det}(\tau) = 1 \\ 0, & \text{if } \text{Det}(\tau) = -1. \end{cases}$$

*Case (2). If  $\tau^3 = 1$ , then  $\nu_{3,\tau}(A) = (\text{Tr}(\tau) + \text{Det}(\tau))^2 + (\text{Tr}(\tau) + 1)(1 - \text{Det}(\tau))$ . Consequently,*

$$\nu_{3,\tau}(A) = \begin{cases} 9, & \text{if } \tau = \text{id} \\ 0, & \text{if } \tau \neq \text{id}. \end{cases}$$

*Proof.* We verify the formulas by using the first part of Corollary 5.2.

Case (1): Recall that  $\Lambda = xy + xyg$  and  $\lambda = (xy)^*$  are right integrals. We must find  $\lambda(\Lambda^{[m,\tau]})$ . First we will show that  $\lambda((xy)^{[m,\tau]}) = \frac{m^2}{4}(1 + \text{Det}(\tau))$ , and then we will argue that  $\lambda((xyg)^{[m,\tau]}) = \lambda((xy)^{[m,\tau]})$ . In order to find  $(xy)^{[m,\tau]}$ , first note that

$$\begin{aligned} \Delta^{m-1}(x) &= x \otimes g^{\otimes m-1} + 1 \otimes x \otimes g^{\otimes m-2} + \cdots + 1^{\otimes i-1} \otimes x \otimes g^{\otimes m-i} + \cdots + 1^{\otimes m-1} \otimes x, \\ \Delta^{m-1}(y) &= y \otimes g^{\otimes m-1} + 1 \otimes y \otimes g^{\otimes m-2} + \cdots + 1^{\otimes i-1} \otimes y \otimes g^{\otimes m-i} + \cdots + 1^{\otimes m-1} \otimes y, \end{aligned}$$

each sum consisting of  $m$  terms. Set

$$x_1 = x \otimes g^{\otimes m-1}, \dots, x_i = 1^{\otimes i-1} \otimes x \otimes g^{\otimes m-i}, \dots, x_m = 1^{\otimes m-1} \otimes x,$$

the index indicating the position of  $x$  in the tensor product, and similarly define  $y_1, y_2, \dots, y_m$ . Letting  $\mu$  denote the multiplication map, by definition we have

$$(xy)^{[m,\tau]} = \mu((1 \otimes \tau)^{\otimes \frac{m}{2}}) \left( \sum_{i,j=1}^m x_i y_j \right).$$

Since  $\tau \cdot g = g$  and  $\lambda = (xy)^*$ , in computing  $\lambda((xy)^{[m,\tau]})$ , the only terms in the above expansion of  $(xy)^{[m,\tau]}$  yielding a nonzero value of  $\lambda$  are those with an even number of factors of  $g$ . These are precisely the terms  $x_i y_j$  for which  $i, j$  have the same parity, of which there are  $\frac{m^2}{2}$  terms. If  $i, j$  are both odd (of which there are  $\frac{m^2}{4}$  pairs), then in  $(xy)^{[m,\tau]}$ , the  $(i, j)$  term is simply  $xy$  by the following observations: (1)  $\tau$  is applied only to factors of  $g$  or 1, which are fixed by  $\tau$ , (2) if  $i \leq j$ , there are an even number of factors of  $g$  between  $x$  and  $y$  after applying  $\mu$ , and (3) if  $i > j$ , there are an odd number of factors of  $g$  between  $x$  and  $y$  after applying  $\mu$  (since  $x_i$  is to the left of  $y_j$ ), so moving factors of  $g$  to the right, past  $x$ , results in a factor of  $(-1)$ , and then applying the relation  $yx = -xy$  results in another factor of  $(-1)$ , so that the end result is a term  $xy$ . If  $i, j$  are both even (of which there are  $\frac{m^2}{4}$  pairs), then in  $(xy)^{[m,\tau]}$ , the  $(i, j)$  term is  $\tau \cdot xy = \text{Det}(\tau)xy$ , by similar reasoning. Therefore

$$\lambda((xy)^{[m,\tau]}) = \lambda \left( \frac{m^2}{4} xy + \frac{m^2}{4} \text{Det}(\tau) xy \right) = \frac{m^2}{4} (1 + \text{Det}(\tau)).$$

Finally, in order to compute  $\lambda((xyg)^{[m,\tau]})$ , note that we need only include an extra factor of  $g^{\otimes m}$  on the right:

$$(xyg)^{[m,\tau]} = \mu((1 \otimes \tau)^{\otimes \frac{m}{2}}) \left( \sum_{i,j=1}^m x_i y_j \right) (g^{\otimes m}).$$

Since  $m$  is even, the number of new factors of  $g$  to be included, in comparison to our previous calculation, is even, and so a similar analysis applies. One checks that the extra factors of  $g$  do not affect the result, and so

$$\lambda((xyg)^{[m,\tau]}) = \lambda((xy)^{[m,\tau]}) = \frac{m^2}{4} (1 + \text{Det}(\tau)).$$

Consequently,  $\nu_{m,\tau}(A) = \lambda(\Lambda^{[m,\tau]}) = \frac{m^2}{2} (1 + \text{Det}(\tau))$ .

To see the conclusion of Case (1), note that since  $\tau^2 = 1$ , the determinant of  $\tau$  is either 1 or  $-1$ .

Case (2): A similar analysis applies. Note that  $\lambda((xy)^{[3,\tau]}) = \mu(1 \otimes \tau \otimes \tau^2) (\sum_{i,j=1}^3 x_i y_j)$  and that  $\tau^2(x) = (a^2 + bc)x + b(a + d)y$ ,  $\tau^2(y) = c(a + d)x + (d^2 + bc)y$ . In evaluating  $\lambda((xy)^{[3,\tau]})$ , we again need only consider  $(i, j)$  terms for which  $i, j$  have the same parity. By contrast, in evaluating  $\lambda((xyg)^{[3,\tau]})$ , we need only consider  $(i, j)$  terms for which  $i, j$  have different parity. Thus we find

$$\begin{aligned} \lambda((xy)^{[3,\tau]}) &= \lambda(xy + x(\tau^2 \cdot y) + yg(\tau^2 \cdot x)g + (\tau^2 \cdot xy) + (\tau \cdot xy)) \\ &= 1 + (d^2 + bc) + (a^2 + bc) + (a^2 + bc)(d^2 + bc) - bc(a + d)^2 + (ad - bc), \\ \lambda((xyg)^{[3,\tau]}) &= \lambda(xg^2(\tau \cdot y)g^4 + yg(\tau \cdot x)g^5 + g(\tau \cdot x)g^2(\tau^2 \cdot y)g + g(\tau \cdot y)g(\tau^2 \cdot x)g^2) \\ &= d + a + a(d^2 + bc) - bc(a + d) - bc(a + d) + d(a^2 + bc). \end{aligned}$$

Adding these together, we have

$$\begin{aligned} \lambda(\Lambda^{[3,\tau]}) &= 1 + a + d + a^2 + ad + d^2 + a^2d + ad^2 + a^2d^2 - abc - 2abcd - bcd + bc + b^2c^2 \\ &= (\text{Tr}(\tau) + \text{Det}(\tau))^2 + (1 + \text{Tr}(\tau))(1 - \text{Det}(\tau)). \end{aligned}$$

To see the conclusion in Case (2), one can check the possible Jordan forms of the matrix for  $\tau$ .  $\square$

## 7. TENSOR PRODUCTS AND CATEGORY OF MODULES

The following theorem generalizes [BW, Theorem 2.1] from the case that  $A$  is a group algebra, to the case that  $A$  is a Hopf algebra. Let  $K = A \natural k^G$  as before, and recall that for  $M$  a  $K$ -module and  $x \in G$ ,  $M_x$  denotes  $p_x \cdot M$ , a  $K$ -submodule of  $M$ , and  $M = \bigoplus_{x \in G} M_x$ . If  $y \in G$ , define  ${}^y M_x$  to be  $M_x$  as a vector space, with  $A$ -module structure given by  $a \cdot_y m = (y^{-1} \cdot a) \cdot m$  for all  $a \in A$ ,  $m \in M$ .

**Theorem 7.1.** *Let  $M, N$  be  $K$ -modules. Then*

- (i)  $(M \otimes N)_x \cong \bigoplus_{\substack{y,z \in G \\ yz=x}} M_y \otimes {}^y N_z$ , and
- (ii)  $(M^*)_x = {}^x(M_{x^{-1}})^*$ .

*Proof.* The proof is a straightforward generalization of that of [BW, Theorem 2.1]. We include details for completeness. We will prove the statement for modules of the form  $M = M_y$ ,  $N = N_z$ . Let  $\phi : M_y \otimes N_z \rightarrow M_y \otimes^y N_z$ , where the target module is a  $K$ -module on which  $p_{yz}$  acts as the identity and  $p_w$  acts as 0 for  $w \neq yz$ , be defined by  $\phi(m \otimes n) = m \otimes n$  for all  $m \in M_y$ ,  $n \in N_z$ . We check that  $\phi$  is a  $K$ -module homomorphism: Let  $x \in G$ ,  $a \in A$ . Apply  $\Delta$  to  $a \natural p_x$  to obtain

$$\phi((a \natural p_x)(m \otimes n)) = \sum \delta_{x,yz} \phi(a_1 m \otimes (y^{-1} \cdot a_2) n).$$

On the other hand,

$$(a \natural p_x)\phi(m \otimes n) = \sum \delta_{x,yz} a_1 m \otimes (y^{-1} \cdot a_2) n.$$

As  $\phi$  is a bijection by its definition, it is an isomorphism of  $K$ -modules.

We will prove that since  $M = M_y$ , its dual satisfies  $M^* = (M^*)_{y^{-1}}$ , and that the corresponding underlying  $A$ -module structure on the vector space  $(M^*)_{y^{-1}}$  is isomorphic to  $y^{-1}(M_y)^*$ . To see this, first let  $x \in G$ ,  $f \in M^*$ , and  $m \in M$ . Then

$$((1 \natural p_x)(f))(m) = f((1 \natural p_{x^{-1}})m) = \delta_{x^{-1},y} f(m).$$

It follows that  $(M_y)^* = M^* = (M^*)_{y^{-1}}$ , as claimed. The  $A$ -module structure on  $(M^*)_{y^{-1}}$  may be determined by considering the action on  $M^*$  of all elements of  $K$  of the form  $a \natural p_{y^{-1}}$  where  $a \in A$ . Let  $f \in M^*$  and  $m \in M$ . Then

$$((a \natural p_{y^{-1}})(f))(m) = f(S(a \natural p_{y^{-1}})m) = f((y \cdot S(a))m).$$

Considering the restriction of  $M^* = (M^*)_{y^{-1}}$  to an  $A$ -module in this way, we see that the action of  $a$  on the vector space  $(M_y)^*$  is that of  $a$  on the  $A$ -module  $y^{-1}(M_y)^*$ :

$$(a \cdot_{y^{-1}} f)(m) = ((y \cdot a)f)(m) = f(S(y \cdot a)m) = f((y \cdot S(a))m).$$

Therefore the  $A$ -module structure on the vector space  $(M^*)_{y^{-1}}$  is that of the  $A$ -module  $y^{-1}(M_y)^*$ .  $\square$

**Remark 7.2.** As a consequence of the theorem, the category of  $K$ -modules is equivalent to the *semidirect product tensor category*  $\mathcal{C} \rtimes G$  where  $\mathcal{C}$  is the category of  $A$ -modules. By definition,  $\mathcal{C} \rtimes G$  is the category  $\bigoplus_{g \in G} \mathcal{C}$ , with objects  $\bigoplus_{g \in G} (M_g, g)$  where each  $M_g$  is an object of  $\mathcal{C}$ , and tensor product  $(M, g) \otimes (N, h) = (M \otimes^g N, gh)$ . See [T], where the notation  $\mathcal{C}[G]$  is used instead for this semidirect product category. For other occurrences of  $\mathcal{C} \rtimes G$  in the literature, see, for example, [GNaNi, Ni].

## REFERENCES

- [AD] N. Andruskiewitsch and F. Dumas, On the automorphisms of  $U_q^+(\mathfrak{g})$ , In: Quantum groups, IRMA Lectures on Mathematical and Theoretical Physics, vol. **12**, pp. 107–133. European Mathematical Society, Zurich (2008).
- [BW] D. Benson and S. Witherspoon, Examples of support varieties for Hopf algebras with non-commutative tensor products, *Archiv der Mathematik* **102** (2014), no. 6, 513–520.
- [DE] S. Danz and K. Erdmann, Crossed products as twisted category algebras, *Algebr. Represent. Theor.* DOI 10.1007/S10468-014-9493-8.
- [EG] P. Etingof and S. Gelaki, On the exponent of finite-dimensional Hopf algebras. *Math. Res. Lett.*, **6**(2):131–140, 1999.
- [GNaNi] S. Gelaki, D. Naidu, and D. Nikshych, Centers of graded fusion categories, *Algebra Number Theory* **3** (2009), no. 8, 959–990.

- [KMN] Y. Kashina, S. Montgomery, and S. Ng, On the trace of the antipode and higher indicators, *Israel J. Math.* **188** (2012), 57–89.
- [KSZ] Y. Kashina, Y. Sommerhuser, and Y. Zhu, *On Higher Frobenius-Schur Indicators*, *Mem. Amer. Math. Soc.* **181** (2006), no. 855, viii+65 pp.
- [Ke] M. Keilberg, Automorphisms of the doubles of purely non-abelian finite groups, *Algebras and Representation Theory*, to appear; arXiv:1311.0575.
- [LM] V. Linchenko and S. Montgomery, A Frobenius-Schur theorem for Hopf algebras, *Algebr. Represent. Theory* **3** (2000), 347–355.
- [Ml] R. Molnar, Semi-direct products of Hopf algebras, *J. Algebra* **47** (1977), 29–51.
- [M] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, CBMS Lectures Vol. **82**, Amer. Math. Soc., Providence, 1997.
- [NSc] S.-H. Ng and P. Schauenburg, Higher Frobenius-Schur indicators for pivotal categories, *Hopf Algebras and Generalizations*, AMS Contemp. Math. 441, AMS, Providence, RI, 2007, 63–90.
- [Ni] D. Nikshych, Non-group-theoretical semisimple Hopf algebras from group actions on fusion categories, *Selecta Math.* **14** (2008), no. 1, 145–161.
- [R] D. E. Radford, The group of automorphisms of a semisimple Hopf algebra over a field of characteristic 0 is finite, *Amer. J. Math.* **112** (1990), 331–357.
- [R2] D. E. Radford, *Hopf Algebras*, World Scientific Publishing, 2012.
- [R3] D. E. Radford, On automorphisms of biproducts, arXiv:1503.00381.
- [SV] D. Sage and M. Vega, Twisted Frobenius-Schur indicators for Hopf algebras, *J. Algebra* **354** (2012), 136–147.
- [SV2] D. Sage and M. Vega, Twisted exponents and twisted Frobenius-Schur indicators for Hopf algebras, *Communications in Algebra*, to appear; arXiv:1402.5201.
- [SV3] D. Sage and M. Vega, Twisted Frobenius-Schur indicators for pivotal categories, in preparation.
- [Sh] K. Shimizu, Some computations of Frobenius-Schur indicators of the regular representations of Hopf algebras, *Algebr. Represent. Theory* **15** (2012), 325–357.
- [T] D. Tambara, Invariants and semi-direct products for finite group actions on tensor categories, *J. Math. Soc. Japan* **53** (2001), no. 2, 429–456.
- [Y] M. Yakimov, Rigidity of quantum tori and the Andruskiewitsch-Dumas conjecture, *Selecta Math (N.S.)* **20** (2014), no. 2, 421–464.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA  
*E-mail address:* smontgom@usc.edu

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC  
*E-mail address:* mdvega@ncsu.edu

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX  
*E-mail address:* sjw@math.tamu.edu