

Atom interferometry in the presence of an external test mass

B. Dubetsky,¹ M. Shverdin,¹ S. B. Libby,² P. R. Berman³

¹*AOSense Inc. 929 E Arques Ave, Sunnyvale, CA 94085*

²*Physics Division, Physical and Life Sciences, Lawrence Livermore National Laboratory, Livermore, California 94550, USA*

³*Physics Department, University of Michigan, Ann Arbor, Michigan 48109-1040*

(Dated: January 8, 2016)

The influence of an external test mass on the phase of the signal of an atom interferometer is studied theoretically. Using traditional techniques in atom optics based on the density matrix equations in the Wigner representation, we are able to extract the various contributions to the phase of the signal associated with the classical motion of the atoms, the quantum correction to this motion resulting from atomic recoil that is produced when the atoms interact with Raman field pulses, and quantum corrections to the atomic motion that occur in the time between the Raman field pulses. By increasing the effective wave vector associated with the Raman field pulses using modified field parameters, we can increase the sensitivity of the signal to the point where such quantum corrections can be measured. The expressions that are derived can be evaluated numerically to isolate the contribution to the signal from an external test mass. The regions of validity of the exact and approximate expressions are determined.

PACS numbers: 03.75.Dg, 37.25.+k, 04.80.-y

I. INTRODUCTION

Since its birth about 30 years ago [1], the field of atom interferometry (AI) has matured significantly. Experiments based on AI have been used to measure fundamental constants [2–5], the acceleration of gravity near the Earth’s surface [6–9], the gradient of the Earth’s gravitational field [4, 10, 11], and the curvature of the gravitational field produced by source masses [12]. Atom interferometer gyroscopes allow one to measure rotation rates; experiments have utilized optical fields [13], nanofabricated structures [14], and three or four spatially or temporally separated sets of fields that drive Raman transitions to split and recombine the matter waves [15–18]. There have been limits set on a non-Newtonian Yukawa-type fifth force [19] and on dark energy [20] using AI, as well as theoretical proposals for using AI to measure general relativity effects [21, 22], including gravitational waves [23]. A detailed theoretical analysis of the combined effect of rotation and gravity on the AI signal has been given [17], based on three- and four-pulse Raman schemes.

Atom interferometry has also been used to probe the gravitational field produced by a heavy test mass [4, 5, 12, 19, 20]. Using a double-difference technique [4] one can extract that part of the phase of the AI signal caused by the gravitational field of the test mass. This article provides a theoretical calculation of this contribution to the phase, based on an atom interferometer using three Raman field pulses. The results can be used to optimize measurements of the Newtonian gravitational constant G and to provide a complete derivation of results outlined in a previous paper [24]. Additionally, recently, an analytic, semi-classical expression for the phase response of an atom interferometer to an arbitrarily placed, stationary point mass has been derived in [25].

A. Estimated Phase Corrections resulting from the Test Mass

The phase in an atom interferometer depends on the interactions of the atoms with the applied Raman fields as well as the motion of the atoms between and following the applied Raman pulses. The presence of a gravitational potential modifies the atomic trajectories, leading to a modification of the AI phase. It is this modification of the phase that serves as a measure of the sensitivity of AI to gravitational effects. Since the Earth’s gravitational potential is only slightly inhomogeneous over the physical extent of the atom interferometer, it can be approximated by a Taylor series in which only the lead and gradient terms are retained [26]. Approximate solutions for the atomic trajectories were obtained in Refs. [17, 27], where effects related to the Earth’s rotation (centripetal and Coriolis forces) were also included. An exact expression for the atom trajectories with these combination of forces has also been derived when the Earth is replaced by a non-spherical mass, rotating with constant angular velocity [28].

The situation can change dramatically if a massive test object is brought close to the interferometer. The accumulated phase produced by the test mass’ gravitational field, $\delta\mathbf{g}(\mathbf{x},t)$, increases with decreasing distance y_{\min} between the test mass and the trajectories of the atoms in the interferometer and also increases with increasing delay times T between the Raman pulses. For sufficiently long T and small y_{\min} , it is no longer a good approximation to retain only the lead and gradient terms when considering the gravitational potential associated with the test mass.

The maximum value of T is limited by experimental considerations; the largest delay time that has been achieved is $T = 1.15$ s [29]. Even for smaller delay times, the inhomogeneity of the field can be significant. For example, with $T = 200$ ms, in a symmetric fountain geometry [30], the length of the atomic trajectory is longer than

$$L = \frac{1}{2}gT^2 = 0.196 \text{ m}, \quad (1)$$

where \mathbf{g} is Earth's gravitational field. With $y_{\min} \lesssim 0.1$ m [see Eqs. (103) and (104) in Sec. III], the usually accepted assumption that the gravitational potential is constant or slightly inhomogeneous along the atom trajectory becomes invalid.

In calculating the atomic trajectories, we can assume [31] that the magnitude δg of the gravitational field of the test mass at the position of the atoms in the interferometer is much less than that of the Earth's field [32],

$$\delta g \ll g. \quad (2)$$

Nevertheless, both the average field and field gradient associated with the test mass can modify the phase of the AI signal. Let us denote the average field of the test mass over the interferometric path as $\overline{\delta g}$. The interferometric phase $\delta\phi$ associated with this average field strength is of order [6–9]

$$\delta\phi \sim k\overline{\delta g}T^2, \quad (3)$$

where k is an effective wave vector of the Raman field and T is the time delay between Raman pulses. This phase change arises owing to the acceleration of the atoms produced by the field of the test mass.

In addition to this "classical" contribution to the phase, there are quantum corrections whose effect we would now like to estimate. Atom interferometers that make use of copropagating optical fields or copropagating Raman pulses as their beam splitters and combiners have a signal phase that is insensitive to quantum corrections if the gravity field is homogeneous. Quantum corrections arise as a result of rotation [33] or inhomogeneous field terms [27]. Quantum corrections ϕ_q to the phase from an inhomogeneous gravitational field are of order

$$\phi_q \sim \frac{\hbar k^2}{M_a} \gamma T^3, \quad (4)$$

where M_a is an atomic mass and γ is the magnitude of the relevant terms in the gravity-gradient tensor. One can understand the estimate (4) as the quantum part of the phase addition $k\gamma v T^3$ associated with the change of atomic velocity $v = \hbar k/M_a$ owing to recoil [34] after interaction with a Raman pulse. A reasonable estimate for γ is $\gamma \sim \overline{\delta g}/L$, where L is the length of the atomic trajectory in the interferometer. As a consequence, we find

$$\frac{\phi_q}{\delta\phi} \sim \frac{\hbar k T}{M_a L}. \quad (5)$$

For Rb⁸⁷ and $k \approx 1.61 \times 10^7 \text{ m}^{-1}$

$$\frac{\phi_q}{\delta\phi} \sim 1.2 \times 10^{-2}. \quad (6)$$

Calculations [24] indicate that $\delta\phi$ can be as large as 1 rad, implying that ϕ_q can be as large as 10^{-2} rad. Since a lower limit for the phase noise in the interferometer is of order [29]

$$\phi_{err} = 10^{-3} \text{ rad}, \quad (7)$$

one sees that quantum corrections ϕ_q are small but measurable; we will include them in our considerations.

Another type of quantum correction is produced during the free evolution of the atomic coherence between the Raman pulses. We formulate the problem in terms of the Wigner representation for the atomic density matrix, $\rho(\mathbf{x}, \mathbf{p}, t)$. This is a standard approach for studying phenomena related to the quantization of the atomic center of mass motion [34] and has also been used in the context of AI [17, 35]. The convenience of this approach is that, for the time between Raman pulses, $\rho(\mathbf{x}, \mathbf{p}, t)$ obeys an equation that is similar to the classical Liouville equation for the distribution function [17].

The Wigner distribution function can be written as,

$$\rho(\mathbf{x}, \mathbf{p}, t) = \frac{1}{(2\pi\hbar)^3} \int d\mathbf{s} \rho_x \left(\mathbf{x} + \frac{1}{2}\mathbf{s}, \mathbf{x} - \frac{1}{2}\mathbf{s}, t \right) \exp(-i\mathbf{p} \cdot \mathbf{s}/\hbar) \quad (8a)$$

$$= \frac{1}{(2\pi\hbar)^3} \int d\mathbf{u} \rho_p \left(\mathbf{p} + \frac{1}{2}\mathbf{u}, \mathbf{p} - \frac{1}{2}\mathbf{u}, t \right) \exp(i\mathbf{x} \cdot \mathbf{u}/\hbar) \quad (8b)$$

where $\rho_x(\mathbf{x}, \mathbf{x}', t)$ is the density matrix in the coordinate representation and $\rho_p(\mathbf{p}, \mathbf{p}', t)$ is the density matrix in the momentum representation. To estimate the quantum corrections we start from the time evolution equation for the Wigner function for times between the application of the Raman pulses. In the absence of the Earth's rotation, this equation can be written as [17]

$$\left\{ \partial_t + \frac{\mathbf{p}}{M_a} \partial_{\mathbf{x}} - \partial_{\mathbf{x}} U \partial_{\mathbf{p}} + Q \right\} \rho(\mathbf{x}, \mathbf{p}, t) = 0, \quad (9a)$$

$$Q = -(i\hbar)^{-1} \left[U \left(\mathbf{x} + \frac{1}{2} i\hbar \partial_{\mathbf{p}} \right) - U \left(\mathbf{x} - \frac{1}{2} i\hbar \partial_{\mathbf{p}} \right) \right] + \partial_{\mathbf{x}} U \partial_{\mathbf{p}} \quad (9b)$$

where $U(\mathbf{x})$ is the gravitational potential. For nearly homogeneous fields, such as the Earth's field, the potential functions in Eq. (9b) can be expanded to first order in \hbar . In that limit, one finds that $Q \sim 0$ and that the Wigner function obeys the same Liouville equation as the classical density matrix in the time between pulses. In the presence of a test mass, however, the gravitational potential is strongly inhomogeneous and higher order terms in the expansion are needed.

Let us estimate the correction from these higher order terms. If the term $\partial_{\mathbf{x}} U \partial_{\mathbf{p}}$ in the Liouville equation (9a) is responsible for the phase $\delta\phi$ in Eq. (3), then the Q -term results in a quantum correction

$$\phi_Q \sim \frac{Q}{\partial_{\mathbf{x}} U \partial_{\mathbf{p}}} \delta\phi. \quad (10)$$

The density matrix depends on atomic momentum p in two characteristic ways. There is both a thermal momentum

$$p_0 = \sqrt{2M_a k_B T_C} \quad (11)$$

(k_B is Boltzmann constant, T_C is atom cloud temperature) and a momentum associated with the Doppler phase,

$$p_D \sim \frac{M_a}{kT}. \quad (12)$$

For Rb at temperature $T_C \approx 1\mu\text{K}$, $k = 1.61 \times 10^7 \text{ m}^{-1}$, and $T = 200 \text{ ms}$,

$$\frac{p_D}{p_0} = \frac{1}{k v_0 T} \sim 2.2 \times 10^{-5}, \quad (13)$$

where $v_0 = p_0/M_a = 0.014 \text{ m/s}$ is the thermal velocity. In qualitative terms, we can think of the dependence of $\rho(\mathbf{x}, \mathbf{p}, t)$ on momentum to vary as

$$e^{-p^2/p_0^2} e^{ip/p_D},$$

where the first factor gives the thermal distribution and the second a phase factor resulting from the accumulated Doppler phase between Raman pulses.

If $\rho(\mathbf{x}, \mathbf{p}, t) \sim e^{-p^2/p_0^2} e^{ip/p_D}$, it follows that the Doppler phase factor makes the dominant contribution to the momentum gradient since

$$\partial_{\mathbf{p}} \sim p_D^{-1} \gg p_0^{-1}. \quad (14)$$

To estimate the quantum corrections, we expand, Eq. (9b) to second order in \hbar to obtain

$$Q \sim \frac{2\hbar^2}{3!} \partial_{\mathbf{x}}^3 U \left(\frac{\partial_{\mathbf{p}}}{2} \right)^3. \quad (15)$$

Replacing $\partial_{\mathbf{x}}^n U$ by U/L^n and $\partial_{\mathbf{p}}$ by p_D^{-1} , we find

$$\frac{\phi_Q}{\delta\phi} \sim \frac{Q}{\partial_{\mathbf{x}} U \partial_{\mathbf{p}}} \sim \frac{1}{24} \frac{\partial_{\mathbf{x}}^2 U}{U p_D^2} = \frac{1}{24} \left(\frac{\hbar k T}{L M_a} \right)^2 = 5.8 \times 10^{-6}. \quad (16)$$

Consistent with the phase noise given in Eq. (7), one should ignore the Q -term in Eq. (9a). However if one uses the AI technique to measure the Newtonian gravitational constant G with an accuracy of several ppm (the level achieved is already 150ppm [5]) then the Q -term should be included. Anticipating innovations capable of reducing the phase

noise to $\phi_{err} \sim 3 \times 10^{-7}$ rad [36], one has to include the Q -term. Consequently, we will include the corrections resulting from this term.

To summarize, there are two types of quantum corrections to the AI phase that are to be considered. The first, ϕ_q , arises from inhomogeneous gravitational field modifications of the Doppler phase associated the recoil the atoms undergo on interacting with the Raman fields. The ratio $\phi_q/\delta\phi$ is of order $\hbar kT/M_a L$. The second, ϕ_Q , arises from quantum corrections to the off-diagonal elements of the Wigner distribution during periods of free evolution. The ratio $\phi_Q/\delta\phi$ is of order $(\hbar kT/M_a L)^2$.

It is possible to increase both $\delta\phi$ and the quantum corrections ϕ_q and ϕ_Q using larger values of the effective wave vector k . Moreover, since $\delta\phi \propto k$, $\phi_q \propto k^2$ and $\phi_Q \propto k^3$ [see Eqs. (3, 4, 16)], the relative weight of the quantum corrections also increases. There are at least four ways to increase k : production of higher order atomic density harmonics in a standing wave field in the Raman-Nath regime [see Eq. (4) in [1]], higher order Bragg scattering [37], multicolor techniques [38], and Raman standing wave techniques [39]. For example, standing wave pulses in the Raman-Nath regime were used to produce the 10th harmonic of the atomic density without excessive loss of signal magnitude and without sub-recoil cooling [40]. A $4\hbar k$ beam splitter was demonstrated using an extension of the Raman standing wave technique [41] and a $51\hbar k$ beam splitter has been produced using higher order Bragg scattering [42]. Recently a high order Bragg scattering atom interferometer was used to determine the fine structure constant with a resolution 0.25ppb [3]. On the theoretical side, it was shown that, with a proper choice of field polarization, Raman standing waves in the Raman-Nath regime can be used to create a $4\hbar k$ beam splitter without increasing the number of separated Raman pulses [39, 43]. To account for such enhancements, our calculations of the AI's phase are carried out for an effective k -vector that is scaled by an integer factor n_k .

This article is arranged as follows. In the next section we derive exact and approximate expressions for the phases $\delta\phi$, ϕ_q , and ϕ_Q . The results of numerical calculations of the phases are given in Sec. III for a stationary test mass and a test mass moving at constant velocity. The calculations enable us to establish the regions of validity of the approximate expressions for the phases.

II. BASIC FORMALISM

The working medium of the atom interferometer consists of a cloud of atoms that are launched with some initial velocity at $t = 0$. The cloud interacts with three Raman pulses that are separated in time. In the time intervals between the pulses, the atoms move under the influence of a gravitational potential $U(\mathbf{x}, t)$. The cloud is assumed to be characterized by a Wigner distribution $f(\mathbf{x}, \mathbf{p})$ at time $t = 0$ and the cloud is assumed to be sufficiently localized such that, at any time, the gravitational field is the same for all atoms in the cloud. In other words, the cloud can be considered as a point insofar as its interactions with both the Earth's and the test mass' gravitational fields. The problem can be broken down into periods of "free evolution" of density matrix elements before the first Raman pulse is applied and for the time intervals between subsequent Raman pulses and into time intervals in which the Raman fields are applied. By "free evolution," we mean evolution in the absence of applied radiation fields, but including the effects produced by $U(\mathbf{x}, t)$. We consider each region separately and then piece together the total response.

We will see that the quantum corrections leading to ϕ_q originate in the recoil the atoms undergo as a result of their interaction with the Raman pulses. Following the interactions this recoil leads to a contribution to the Doppler phase of the off-diagonal density matrix elements ρ_{eg} (g and e are sublevels of the atoms' ground state manifold) in the time intervals between the pulses. In addition, the momentum derivatives of the Doppler phase factors give rise to the Q -term corrections; as such, the Q -term corrections depend only on the free evolution of off-diagonal density matrix elements between the pulses.

A. Density matrix evolution between the Raman pulses

Between the Raman pulses, the Wigner function evolves according to Eqs. (9). When the distance L over which the gravitational potential energy varies significantly is much larger than \hbar divided by the characteristic width Δp of the momentum distribution, *i.e.*

$$\frac{\hbar}{\Delta p L} \ll 1, \quad (17)$$

we can expand Q [Eq. (9b)] in a power series in \hbar to obtain

$$Q \approx -\frac{\hbar^2}{24} \chi'_{ikl}(\mathbf{x}, t) \partial_{\mathbf{p}_i} \partial_{\mathbf{p}_k} \partial_{\mathbf{p}_l}, \quad (18)$$

where

$$\chi'_{ikl}(\mathbf{x}, t) = -\partial_{x_i} \partial_{x_k} \partial_{x_l} U(\mathbf{x}, t). \quad (19)$$

A summation convention implicit in Eq. (18) will be used in all subsequent equations. Repeated indices and symbols appearing on the right-hand-side (rhs) of an equation are to be summed over, unless they also appear on the left-hand side (lhs) of that equation.

We have already shown in Eq. (16) that the Q -term can be considered as a small perturbation, allowing us to write

$$\rho(\mathbf{x}, \mathbf{p}, t) = \rho_0(\mathbf{x}, \mathbf{p}, t) + \rho_Q(\mathbf{x}, \mathbf{p}, t), \quad (20)$$

where $\rho_0(\mathbf{x}, \mathbf{p}, t)$ is the unperturbed density matrix obeying the equation

$$\left\{ \partial_t + \frac{\mathbf{P}}{M_a} \partial_{\mathbf{x}} - \partial_{\mathbf{x}} U(\mathbf{x}, t) \partial_{\mathbf{p}} \right\} \rho_0(\mathbf{x}, \mathbf{p}, t) = 0 \quad (21)$$

and $\rho_Q(\mathbf{x}, \mathbf{p}, t)$ is a perturbation whose evolution is governed by the equation

$$\left\{ \partial_t + \frac{\mathbf{P}}{M_a} \partial_{\mathbf{x}} - \partial_{\mathbf{x}} U(\mathbf{x}, t) \partial_{\mathbf{p}} \right\} \rho_Q(\mathbf{x}, \mathbf{p}, t) = -Q \rho_0(\mathbf{x}, \mathbf{p}, t). \quad (22)$$

The $\rho_0(\mathbf{x}, \mathbf{p}, t)$ term contains the ϕ_q corrections, while the $\rho_Q(\mathbf{x}, \mathbf{p}, t)$ term provides the ϕ_Q corrections.

Equation (21) has been studied in Ref. [17] for the Earth's gravitational field. In this article we obtain a solution of Eq. (21) in the presence of a test mass and solve Eq. (22) to get the contribution to the AI phase arising from the Q -term. We assume that density matrix is known at some preceding time t' and arbitrarily set $\rho_Q(\mathbf{x}, \mathbf{p}, t') = 0$ at this time, such that, at $t = t'$, $\rho_0(\mathbf{x}, \mathbf{p}, t') = \rho(\mathbf{x}, \mathbf{p}, t')$. The solution of the homogeneous Eq. (21) is then given by [17]

$$\rho_0(\mathbf{x}, \mathbf{p}, t) = \rho[\mathbf{X}(\mathbf{x}, \mathbf{p}, t', t), \mathbf{P}(\mathbf{x}, \mathbf{p}, t', t), t'], \quad (23)$$

where $\{\mathbf{X}(\mathbf{x}, \mathbf{p}, t_1, t_2), \mathbf{P}(\mathbf{x}, \mathbf{p}, t_1, t_2)\}$ are atomic classical position and momentum at time t_1 subject to the constraint that the position and momentum are specified by $\{\mathbf{x}, \mathbf{p}\}$ at time t_2 . In other words, in Eq. (23), we look for the values $\{\mathbf{X}(\mathbf{x}, \mathbf{p}, t', t), \mathbf{P}(\mathbf{x}, \mathbf{p}, t', t)\}$ for which $\{\mathbf{X}(t'), \mathbf{P}(t')\}$ will lead to values $\{\mathbf{X}(t), \mathbf{P}(t)\} = \{\mathbf{x}, \mathbf{p}\}$ under the influence of the applied fields.

Turning our attention to Eq. (22), we see that the curly brackets in that equation is a total time derivative, enabling us to write

$$\frac{d\rho_Q(\mathbf{x}, \mathbf{p}, t'')}{dt''} = -Q \rho_0(\mathbf{x}, \mathbf{p}, t'') \quad (24)$$

Integrating this equation from t'' equals t' to t , using the fact that $\rho_Q(\mathbf{x}, \mathbf{p}, t') = 0$, and making use of Eqs. (23), (18), and (19), we find

$$\rho_Q(\mathbf{x}, \mathbf{p}, t) = \frac{\hbar^2}{24} \int_{t'}^t dt'' [\chi'_{ikl}(\boldsymbol{\xi}, t'') \partial_{\pi_i} \partial_{\pi_k} \partial_{\pi_l} \rho_0(\boldsymbol{\xi}, \boldsymbol{\pi}, t'')]_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, t'', t), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, t'', t)}. \quad (25)$$

Using Eq. (23) one more time, we arrive at

$$\rho_Q(\mathbf{x}, \mathbf{p}, t) = \frac{\hbar^2}{24} \int_{t'}^t dt'' [\chi'_{ikl}(\boldsymbol{\xi}, t'') \partial_{\pi_i} \partial_{\pi_k} \partial_{\pi_l} \rho_0(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, t', t''), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, t', t''), t'')] \left\{ \begin{array}{c} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left\{ \begin{array}{c} \mathbf{X}(\mathbf{x}, \mathbf{p}, t'', t) \\ \mathbf{P}(\mathbf{x}, \mathbf{p}, t'', t) \end{array} \right\}. \quad (26)$$

B. Changes in Density Matrix Elements Produced by the Raman Pulses

Consider now a cloud of atoms having two hyperfine sublevels g and e in the ground state manifold. The atoms are prepared in level g at time $t = 0$ and they proceed to interact with a $\frac{\pi}{2} - \pi - \frac{\pi}{2}$ sequence of Raman pulses applied at times

$$\tau = \{\tau_1, \tau_2 = \tau_1 + T, \tau_3 = \tau_1 + 2T\}, \quad (27)$$

where τ_1 is time delay between cloud launch and the first Raman pulse and T is the time delay between pulses. The initial atomic density matrix (8) is given by

$$\rho_{gg}(\mathbf{x}, \mathbf{p}, 0) = f(\mathbf{x}, \mathbf{p}), \quad (28a)$$

$$\rho_{eg}(\mathbf{x}, \mathbf{p}, 0) = \rho_{ee}(\mathbf{x}, \mathbf{p}, 0) = 0, \quad (28b)$$

where $f(\mathbf{x}, \mathbf{p})$ is the Wigner distribution at $t = 0$.

If a $\pi/2$ -pulse applied at time τ_j , the density matrix elements undergo changes given by [17]

$$\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{j+}) = \frac{1}{2} [\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{j-}) + \rho_{gg}(\mathbf{x}, \mathbf{p} - \hbar\mathbf{k}, \tau_{j-})] + \text{Re} \left\{ i \exp \left[-i \left(\mathbf{k} \cdot \mathbf{x} - \delta_{12}^{(j)} \tau_j - \phi_j \right) \right] \rho_{eg} \left(\mathbf{x}, \mathbf{p} - \frac{\hbar\mathbf{k}}{2}, \tau_{j-} \right) \right\}, \quad (29a)$$

$$\rho_{gg}(\mathbf{x}, \mathbf{p}, \tau_{j+}) = \frac{1}{2} [\rho_{ee}(\mathbf{x}, \mathbf{p} + \hbar\mathbf{k}, \tau_{j-}) + \rho_{gg}(\mathbf{x}, \mathbf{p}, \tau_{j-})] - \text{Re} \left\{ i \exp \left[-i \left(\mathbf{k} \cdot \mathbf{x} - \delta_{12}^{(j)} \tau_j - \phi_j \right) \right] \rho_{eg} \left(\mathbf{x}, \mathbf{p} + \frac{\hbar\mathbf{k}}{2}, \tau_{j-} \right) \right\}, \quad (29b)$$

$$\begin{aligned} \rho_{eg}(\mathbf{x}, \mathbf{p}, \tau_{j+}) &= \frac{i}{2} \exp \left[i \left(\mathbf{k} \cdot \mathbf{x} - \delta_{12}^{(j)} \tau_j - \phi_j \right) \right] \left[\rho_{ee} \left(\mathbf{x}, \mathbf{p} + \frac{\hbar\mathbf{k}}{2}, \tau_{j-} \right) - \rho_{gg} \left(\mathbf{x}, \mathbf{p} - \frac{\hbar\mathbf{k}}{2}, \tau_{j-} \right) \right] \\ &+ \frac{1}{2} \left\{ \rho_{eg}(\mathbf{x}, \mathbf{p}, \tau_{j-}) + \exp \left[2i \left(\mathbf{k} \cdot \mathbf{x} - \delta_{12}^{(j)} \tau_j - \phi_j \right) \right] \rho_{ge}(\mathbf{x}, \mathbf{p}, \tau_{j-}) \right\}, \end{aligned} \quad (29c)$$

Similarly, for π -pulse applied at time τ_j ,

$$\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{j+}) = \rho_{gg}(\mathbf{x}, \mathbf{p} - \hbar\mathbf{k}, \tau_{j-}), \quad (30a)$$

$$\rho_{gg}(\mathbf{x}, \mathbf{p}, \tau_{j+}) = \rho_{ee}(\mathbf{x}, \mathbf{p} + \hbar\mathbf{k}, \tau_{j-}), \quad (30b)$$

$$\rho_{eg}(\mathbf{x}, \mathbf{p}, \tau_{j+}) = \exp \left[2i \left(\mathbf{k} \cdot \mathbf{x} - \delta_{12}^{(j)} \tau_j - \phi_j \right) \right] \rho_{ge}(\mathbf{x}, \mathbf{p}, \tau_{j-}), \quad (30c)$$

In these equations, \mathbf{k} is an effective wave vector (assumed to be the same for all the pulses), $\delta_{12}^{(j)}$ is the detuning between the hyperfine transition frequency and the effective frequency of the Raman fields (that is the frequency difference of the two fields used to create the Raman pulse), ϕ_j is the phase difference between traveling components of the Raman field, and $\tau_{j\pm}$ are times just after and before the pulse. We allow pulses to have different detunings and phases $\delta_{12}^{(j)}, \phi_j$ ($j = 1, 2, 3$). It is assumed that the temporal width of the Raman pulse is sufficiently short that all phases related to the detuning and Doppler shifts are effectively frozen during the application of the pulses.

If a $\pi/2$ pulse acts on a ground state atom, it produces a superposition of ground and excited states. If there was a momentum \mathbf{p} associated with the ground state amplitude $a_g(\mathbf{p})$ before the pulse is applied, the excited state amplitude $a_e(\mathbf{p})$ depends on $a_g(\mathbf{p} - \hbar\mathbf{k})$. As a consequence, the off-diagonal density matrix element following the pulse involves the product of state amplitudes evolving with different momenta. It is this difference in momentum that leads to the Q -term correction in periods of free evolution.

C. AI Signal

Our goal is to calculate $\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{3+})$, the excited state atomic density matrix element following the 3rd Raman pulse, since $\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{3+})$ can be related to experimentally measurable quantities. To carry out the calculation, we use Eqs. (23, 26) for the "free evolution" of density matrix elements before the first Raman pulse is applied and for the time intervals between subsequent Raman pulses and use Eqs. (29, 30) for changes in the density matrix elements resulting from the application of the Raman pulses. In these free evolution regions, density matrix elements are affected by the presence of a gravitational potential that ultimately contributes to the phase of the AI signal.

From the time the cloud is launched at $t = 0$ to the time τ_1 that the first Raman pulse is applied, the only non-vanishing density matrix element is $\rho_{gg}(\mathbf{x}, \mathbf{p}, t)$. In the time interval between $t = 0$ and $t = \tau_1$, this density matrix element evolves to

$$\rho_{gg}(\mathbf{x}, \mathbf{p}, \tau_{1-}) = f(\mathbf{X}(\mathbf{x}, \mathbf{p}, 0, \tau_1), \mathbf{P}(\mathbf{x}, \mathbf{p}, 0, \tau_1)). \quad (31)$$

For reasons to be discussed below, corrections from the Q term can be neglected in this time interval. After the first $\pi/2$ -pulse, the density matrix elements change to

$$\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{1+}) = \frac{1}{2} f(\mathbf{X}(\mathbf{x}, \mathbf{p} - \hbar \mathbf{k}, 0, \tau_1), \mathbf{P}(\mathbf{x}, \mathbf{p} - \hbar \mathbf{k}, 0, \tau_1)), \quad (32a)$$

$$\rho_{gg}(\mathbf{x}, \mathbf{p}, \tau_{1+}) = \frac{1}{2} f(\mathbf{X}(\mathbf{x}, \mathbf{p}, 0, \tau_1), \mathbf{P}(\mathbf{x}, \mathbf{p}, 0, \tau_1)), \quad (32b)$$

$$\rho_{eg}(\mathbf{x}, \mathbf{p}, \tau_{1+}) = -\frac{i}{2} \exp\left[i\left(\mathbf{k} \cdot \mathbf{x} - \delta_{12}^{(1)} \tau_1 - \phi_1\right)\right] f\left(\mathbf{X}\left(\mathbf{x}, \mathbf{p} - \frac{\hbar \mathbf{k}}{2}, 0, \tau_1\right), \mathbf{P}\left(\mathbf{x}, \mathbf{p} - \frac{\hbar \mathbf{k}}{2}, 0, \tau_1\right)\right). \quad (32c)$$

One uses these density matrix elements as initial values for the free evolution between the 1st and 2nd pulses of the unperturbed density matrix; that is,

$$\rho_0(\mathbf{x}, \mathbf{p}, \tau_{1+}) = \rho(\mathbf{x}, \mathbf{p}, \tau_{1+}). \quad (33)$$

We now consider the modifications produced by the Q -term in the time interval between the 1st and 2nd pulses. The modifications produced by the Q -term (26) in the atomic coherence before the second pulse acts, ρ_{Qeg} can be calculated from Eqs. (26, 32c, 33) as

$$\rho_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{2-}) = -i \frac{\hbar^2}{48} \int_{\tau_1}^{\tau_2} dt \times \left\{ \chi'_{ikl}(\boldsymbol{\xi}, t) \partial_{\pi_i} \partial_{\pi_k} \partial_{\pi_l} \left[\exp\left[i\left(\mathbf{k} \cdot \mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) - \delta_{12}^{(1)} \tau_1 - \phi_1\right)\right] \times f\left(\begin{array}{c} \mathbf{X}(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) - \frac{\hbar \mathbf{k}}{2}, 0, \tau_1), \\ \mathbf{P}(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) - \frac{\hbar \mathbf{k}}{2}, 0, \tau_1) \end{array}\right) \right] \right\} \left. \begin{array}{c} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left. \begin{array}{c} \mathbf{x} \\ \mathbf{p} \end{array} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_2)}. \quad (34)$$

In Eq. (34), the $\boldsymbol{\pi}$ derivatives lead to two types of terms. The first of these originates from the thermal distribution and is of order

$$\partial_{\pi_i}^{\text{Thermal}} \sim p_0^{-1}, \quad (35)$$

where p_0 is thermal momentum defined in Eq. (11). The second arises from the phase factor $\exp\left[i\left(\mathbf{k} \cdot \mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) - \delta_{12}^{(1)} \tau_1 - \phi_1\right)\right]$ in Eq. (34), evaluated at $t - \tau_1 \sim T$. To estimate this contribution, we "turn off" the gravitational field. In this approximation

$$\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) = \boldsymbol{\xi} - \boldsymbol{\pi}(t - \tau_1)/M_a \quad (36)$$

and the Doppler phase becomes equal to $\mathbf{k} \cdot \boldsymbol{\pi}(t - \tau_1)/M_a$. This phase factor is a rapidly oscillating function of momentum $\boldsymbol{\pi}$ having period of order p_D defined by Eq. (12), from which we find

$$\partial_{\pi_i}^{\text{Doppler}} \sim p_D^{-1}. \quad (37)$$

In the limit that

$$kv_0 T \gg 1, \quad (38)$$

the thermal derivative is smaller than Doppler derivative by the ratio given in Eq. (13) and can be neglected.

When inequality (38) holds, the time separation between pulses T is sufficiently large to insure that the dominant contributions to the Q -term comes from the momentum derivatives of the Doppler phase factor. As we will show, the atomic levels' populations (ρ_{ee} and ρ_{gg}) have no phase factor for $0 < t < \tau_{3-}$; therefore the Q -term corrections arise only from the atomic coherence ρ_{eg} . As a consequence, we can neglect any contribution to the Q -term corrections from atomic state populations. It was for this reason we did not include any Q -term corrections to the Wigner distribution for the time interval $0 < t < \tau_{1-}$. *In the Doppler limit defined by Eq. (38), the AI phase is pretty much independent of the atomic momentum and spatial distributions.*

Calculating the derivatives and retaining those contributions to the derivatives arising from the Doppler phase only,

we arrive at

$$\begin{aligned} \rho_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{2-}) &= -\frac{\hbar^2}{48} k_u k_v k_w \int_{\tau_1}^{\tau_2} dt \\ &\times [\chi'_{ikl}(\boldsymbol{\xi}, t) \partial_{\pi_i} \mathbf{X}_u(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \partial_{\pi_k} \mathbf{X}_v(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \partial_{\pi_l} \mathbf{X}_w(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)] \left\{ \begin{array}{c} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_2)} \\ &\left\{ \begin{array}{c} \exp \left[i \left(\mathbf{k} \cdot \mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) - \delta_{12}^{(1)} \tau_1 - \phi_1 \right) \right] \\ \times f \left(\begin{array}{c} \mathbf{X}(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) - \frac{\hbar \mathbf{k}}{2}, 0, \tau_1), \\ \mathbf{P}(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) - \frac{\hbar \mathbf{k}}{2}, 0, \tau_1) \end{array} \right) \end{array} \right\} \left\{ \begin{array}{c} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_2)}, \end{aligned} \quad (39)$$

where k_u is the u th component of the effective k -vector. In this approximation, the derivative no longer acts on the term inside the curly brackets. Therefore we can apply the multiplication law,

$$\left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}(\mathbf{X}(\mathbf{x}, \mathbf{p}, t', t''), \mathbf{P}(\mathbf{x}, \mathbf{p}, t', t''), t, t') = \left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}(\mathbf{x}, \mathbf{p}, t, t'') \quad (40)$$

to get

$$\left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, t, \tau_2), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, t, \tau_2)} = \left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2). \quad (41)$$

The expression inside the curly brackets of Eq. (39) becomes t -independent and the Q -term just before the second pulse is given by

$$\begin{aligned} \rho_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{2-}) &= -\frac{\hbar^2}{48} \left\{ \exp \left[i \left(\mathbf{k} \cdot \boldsymbol{\xi} - \delta_{12}^{(1)} \tau_1 - \phi_1 \right) \right] f(\boldsymbol{\xi}, \boldsymbol{\pi}) \right\} \left\{ \begin{array}{c} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left\{ \begin{array}{c} \mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2) \\ \mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2) - \hbar \mathbf{k} / 2 \end{array} \right\} \\ &\times k_u k_v k_w \int_{\tau_1}^{\tau_2} dt [\chi'_{ikl}(\boldsymbol{\xi}, t) \partial_{\pi_i} \mathbf{X}_u(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \partial_{\pi_k} \mathbf{X}_v(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \partial_{\pi_l} \mathbf{X}_w(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)] \left\{ \begin{array}{c} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_2)}. \end{aligned} \quad (42)$$

We still need an expression for the time evolution of $\rho_0(\mathbf{x}, \mathbf{p}, t)$ between the first and second pulses. From Eqs. (23, 32, 40), we find

$$\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{2-}) = \frac{1}{2} f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1))_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2) - \hbar \mathbf{k}}, \quad (43a)$$

$$\rho_{gg}(\mathbf{x}, \mathbf{p}, \tau_{2-}) = \frac{1}{2} f(\mathbf{X}(\mathbf{x}, \mathbf{p}, 0, \tau_2), \mathbf{P}(\mathbf{x}, \mathbf{p}, 0, \tau_2)), \quad (43b)$$

$$\begin{aligned} \rho_{0eg}(\mathbf{x}, \mathbf{p}, \tau_{2-}) &= -\frac{i}{2} \left\{ \exp \left[i \left(\mathbf{k} \cdot \boldsymbol{\xi} - \delta_{12}^{(1)} \tau_1 - \phi_1 \right) \right] \right. \\ &\times f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \left. \right\}_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2) - \hbar \mathbf{k} / 2} \end{aligned} \quad (43c)$$

At time τ_2 the π pulse acts which, according to Eqs. (30), transforms these density matrix elements at time τ_{2-} into

$$\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{2+}) = \frac{1}{2} f(\mathbf{X}(\mathbf{x}, \mathbf{p} - \hbar \mathbf{k}, 0, \tau_2), \mathbf{P}(\mathbf{x}, \mathbf{p} - \hbar \mathbf{k}, 0, \tau_2)), \quad (44a)$$

$$\rho_{gg}(\mathbf{x}, \mathbf{p}, \tau_{2+}) = \frac{1}{2} f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1))_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p} + \hbar \mathbf{k}, \tau_1, \tau_2), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p} + \hbar \mathbf{k}, \tau_1, \tau_2) - \hbar \mathbf{k}} \quad (44b)$$

$$\begin{aligned} \rho_{0eg}(\mathbf{x}, \mathbf{p}, \tau_{2+}) &= \frac{i}{2} \left\{ \exp \left\{ i \left[\mathbf{k} \cdot (2\mathbf{x} - \boldsymbol{\xi}) - 2\delta_{12}^{(2)} \tau_2 + \delta_{12}^{(1)} \tau_1 - 2\phi_2 + \phi_1 \right] \right\} \right. \\ &\times f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \left. \right\}_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2) - \hbar \mathbf{k} / 2}, \end{aligned} \quad (44c)$$

$$\begin{aligned} \rho_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{2+}) &= -\frac{\hbar^2}{48} \left\{ \exp \left\{ i \left[\mathbf{k} \cdot (2\mathbf{x} - \boldsymbol{\xi}) - 2\delta_{12}^{(2)} \tau_2 + \delta_{12}^{(1)} \tau_1 - 2\phi_2 + \phi_1 \right] \right\} \right. \\ &\times f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \left. \right\}_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_2) - \hbar \mathbf{k} / 2} \\ &\times k_u k_v k_w \int_{\tau_1}^{\tau_2} dt [\chi'_{ikl}(\boldsymbol{\xi}, t) \\ &\times \partial_{\pi_i} \mathbf{X}_u(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \partial_{\pi_k} \mathbf{X}_v(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \partial_{\pi_l} \mathbf{X}_w(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)] \left\{ \begin{array}{c} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_2)}. \end{aligned} \quad (44d)$$

The next step is to calculate the Q -term corrections in time interval $[\tau_2, \tau_3]$. Each density matrix element in Eqs. (44) produces a Q -term correction. However, the diagonal matrix elements given by Eqs. (44a, 44b) contain no rapidly oscillating phase factors in momentum space allowing us to neglect their Q -term corrections. Moreover, Eq. (44d) is already linear in Q and can produce only higher order corrections that we neglect in this work. As a consequence, we need consider only the Q -term correction produced by the coherence in Eq. (44c), which we denote as ρ'_{Qeg} . From Eq. (26) we find

$$\begin{aligned} \rho'_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{3-}) &= i \frac{\hbar^2}{48} \int_{\tau_2}^{\tau_3} dt \{ \chi'_{ikl}(\boldsymbol{\xi}, t) \partial_{\pi_i} \partial_{\pi_k} \partial_{\pi_l} \\ &\quad \times \exp \{ i [\mathbf{k} \cdot (2\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, t) - \mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)) \\ &\quad - 2\delta_{12}^{(2)}\tau_2 + \delta_{12}^{(1)}\tau_1 - 2\phi_2 + \phi_1] \} \\ &\quad \times f \left(\mathbf{X} \left(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) - \frac{\hbar\mathbf{k}}{2}, 0, \tau_1 \right) \right. \\ &\quad \left. \mathbf{P} \left(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) - \frac{\hbar\mathbf{k}}{2}, 0, \tau_1 \right) \right) \} \left\{ \begin{matrix} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{matrix} \right\} = \left\{ \begin{matrix} \mathbf{X} \\ \mathbf{P} \end{matrix} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_3)} \end{aligned} \quad (45)$$

where we used the multiplication law (40),

$$\left\{ \begin{matrix} \mathbf{X} \\ \mathbf{P} \end{matrix} \right\} (\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, t), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, t), \tau_1, \tau_2) = \left\{ \begin{matrix} \mathbf{X} \\ \mathbf{P} \end{matrix} \right\} (\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t). \quad (46)$$

In Eq. (45) the differentiation over momentum $\boldsymbol{\pi}$ is carried out only for the Doppler phase factors. After differentiation, we apply the multiplication law two more times to the phase factor and distribution f , namely

$$\left\{ \begin{matrix} \mathbf{X} \\ \mathbf{P} \end{matrix} \right\} (\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_i, t) \left\{ \begin{matrix} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{matrix} \right\} = \left\{ \begin{matrix} \mathbf{X} \\ \mathbf{P} \end{matrix} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_3)} = \left\{ \begin{matrix} \mathbf{X} \\ \mathbf{P} \end{matrix} \right\} (\mathbf{x}, \mathbf{p}, \tau_i, \tau_3) \quad (47)$$

for $i = 1, 2$, and find that these terms become t -independent. As a result one gets for Q -term ρ'_{Qeg} before the third pulse,

$$\begin{aligned} \rho'_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{3-}) &= -\frac{\hbar^2}{48} \left\{ \exp \left\{ i \left[\mathbf{k} \cdot (2\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3) - \boldsymbol{\xi}) - 2\delta_{12}^{(2)}\tau_2 + \delta_{12}^{(1)}\tau_1 - 2\phi_2 + \phi_1 \right] \right\} \right. \\ &\quad \times f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \Big|_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3)-\hbar\mathbf{k}/2} \\ &\quad \times k_u k_v k_w \int_{\tau_2}^{\tau_3} dt \left\{ \chi'_{ikl}(\boldsymbol{\xi}, t) \left[\frac{\partial \mathbf{X}_u(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)}{\partial \pi_i} - 2 \frac{\partial \mathbf{X}_u(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, t)}{\partial \pi_i} \right] \right. \\ &\quad \times \left[\frac{\partial \mathbf{X}_v(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)}{\partial \pi_k} - 2 \frac{\partial \mathbf{X}_v(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, t)}{\partial \pi_k} \right] \\ &\quad \left. \left. \times \left[\frac{\partial \mathbf{X}_w(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)}{\partial \pi_l} - 2 \frac{\partial \mathbf{X}_w(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, t)}{\partial \pi_l} \right] \right\} \right\} \left\{ \begin{matrix} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{matrix} \right\} = \left\{ \begin{matrix} \mathbf{X} \\ \mathbf{P} \end{matrix} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_3)}. \end{aligned} \quad (48)$$

The value of $\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{3+})$ will depend both on $\rho'_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{3-})$ and $\rho_0(\mathbf{x}, \mathbf{p}, \tau_{3-})$. In other words, we must also

calculate the time evolution of $\rho_0(\mathbf{x}, \mathbf{p}, t)$ between the second and third pulses. Applying Eq. (23), we find

$$\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{3-}) = \frac{1}{2} f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_2), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_2))_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3)-\hbar\mathbf{k}}; \quad (49a)$$

$$\rho_{gg}(\mathbf{x}, \mathbf{p}, \tau_{3-}) = \frac{1}{2} f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1))_{\substack{\boldsymbol{\xi}=\mathbf{X}(\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3), \mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3)+\hbar\mathbf{k}, \tau_1, \tau_2), \\ \boldsymbol{\pi}=\mathbf{P}(\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3), \mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3)+\hbar\mathbf{k}, \tau_1, \tau_2)-\hbar\mathbf{k}}} \quad (49b)$$

$$\rho_{0eg}(\mathbf{x}, \mathbf{p}, \tau_{3-}) = \frac{i}{2} \left\{ \exp \left\{ i \left[\mathbf{k} \cdot (2\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3) - \boldsymbol{\xi}) - 2\delta_{12}^{(2)}\tau_2 + \delta_{12}^{(1)}\tau_1 - 2\phi_2 + \phi_1 \right] \right\} \right. \\ \left. \times f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \right\}_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3)-\hbar\mathbf{k}/2}, \quad (49c)$$

$$\rho_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{3-}) = -\frac{\hbar^2}{48} \left\{ \exp \left\{ i \left[\mathbf{k} \cdot (2\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3) - \boldsymbol{\xi}) - 2\delta_{12}^{(2)}\tau_2 + \delta_{12}^{(1)}\tau_1 - 2\phi_2 + \phi_1 \right] \right\} \right. \\ \left. \times f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \right\}_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3)-\hbar\mathbf{k}/2} \\ \times \mathbf{k}_u \mathbf{k}_v \mathbf{k}_w \int_{\tau_1}^{\tau_2} dt [\chi'_{ikl}(\boldsymbol{\xi}, t) \partial_{\pi_i} \mathbf{X}_u(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \partial_{\pi_k} \mathbf{X}_v(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \\ \times \partial_{\pi_l} \mathbf{X}_w(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)] \left\{ \begin{array}{l} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left\{ \begin{array}{l} \mathbf{X} \\ \mathbf{P} \end{array} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_3)}. \quad (49d)$$

Combining the different contributions to the off diagonal density matrix element given by Eqs. (49c, 49d, 48) and factoring out a common phase factor, we obtain

$$\rho_{eg}(\mathbf{x}, \mathbf{p}, \tau_{3-}) \approx \rho_{0eg}(\mathbf{x}, \mathbf{p}, \tau_{3-}) + \rho_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{3-}) + \rho'_{Qeg}(\mathbf{x}, \mathbf{p}, \tau_{3-}) \\ = \frac{i}{2} \left\{ \exp \left\{ i \left[\mathbf{k} \cdot (2\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3) - \boldsymbol{\xi}) - 2\delta_{12}^{(2)}\tau_2 + \delta_{12}^{(1)}\tau_1 - 2\phi_2 + \phi_1 \right] \right\} \right. \\ \left. \times f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \right\}_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3)-\hbar\mathbf{k}/2} \left[1 - i\tilde{\phi}_Q(\mathbf{x}, \mathbf{p}) \right] \quad (50a)$$

$$\approx \frac{i}{2} \left\{ \exp \left\{ i \left[\mathbf{k} \cdot (2\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3) - \boldsymbol{\xi}) - \tilde{\phi}_Q(\mathbf{x}, \mathbf{p}) - 2\delta_{12}^{(2)}\tau_2 + \delta_{12}^{(1)}\tau_1 - 2\phi_2 + \phi_1 \right] \right\} \right. \\ \left. \times f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \right\}_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, \tau_3)-\hbar\mathbf{k}/2}, \quad (50b)$$

$$\tilde{\phi}_Q(\mathbf{x}, \mathbf{p}) = -\frac{\hbar^2}{24} k_u k_v k_w \left\{ \int_{\tau_1}^{\tau_2} dt \chi'_{ikl}(\boldsymbol{\xi}', t) \partial_{\pi'_i} \mathbf{X}_u(\boldsymbol{\xi}', \boldsymbol{\pi}', \tau_1, t) \partial_{\pi'_k} \mathbf{X}_v(\boldsymbol{\xi}', \boldsymbol{\pi}', \tau_1, t) \partial_{\pi'_l} \mathbf{X}_w(\boldsymbol{\xi}', \boldsymbol{\pi}', \tau_1, t) \right. \\ \left. + \int_{\tau_2}^{\tau_3} dt \chi'_{ikl}(\boldsymbol{\xi}', t) \left[\frac{\partial \mathbf{X}_u(\boldsymbol{\xi}', \boldsymbol{\pi}', \tau_1, t)}{\partial \pi'_i} - 2 \frac{\partial \mathbf{X}_u(\boldsymbol{\xi}', \boldsymbol{\pi}', \tau_2, t)}{\partial \pi'_i} \right] \left[\frac{\partial \mathbf{X}_v(\boldsymbol{\xi}', \boldsymbol{\pi}', \tau_1, t)}{\partial \pi'_k} - 2 \frac{\partial \mathbf{X}_v(\boldsymbol{\xi}', \boldsymbol{\pi}', \tau_2, t)}{\partial \pi'_k} \right] \right. \\ \left. \times \left[\frac{\partial \mathbf{X}_w(\boldsymbol{\xi}', \boldsymbol{\pi}', \tau_1, t)}{\partial \pi'_l} - 2 \frac{\partial \mathbf{X}_w(\boldsymbol{\xi}', \boldsymbol{\pi}', \tau_2, t)}{\partial \pi'_l} \right] \right\} \left\{ \begin{array}{l} \boldsymbol{\xi}' \\ \boldsymbol{\pi}' \end{array} \right\} = \left\{ \begin{array}{l} \mathbf{X} \\ \mathbf{P} \end{array} \right\}_{(\mathbf{x}, \mathbf{p}, t, \tau_3)}. \quad (50c)$$

Finally we use Eqs. (29a, 49a, 49b, 50) to calculate $\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{3+})$ following the $\pi/2$ pulse at time τ_3 as

$$\rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{3+}) = \frac{1}{4} f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_2), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_2))_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_2, \tau_3)-\hbar\mathbf{k}} \\ + \frac{1}{4} f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \left\{ \begin{array}{l} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left\{ \begin{array}{l} \mathbf{X}(\mathbf{X}(\mathbf{x}, \mathbf{p}-\hbar\mathbf{k}, \tau_2, \tau_3), \mathbf{P}(\mathbf{x}, \mathbf{p}-\hbar\mathbf{k}, \tau_2, \tau_3)+\hbar\mathbf{k}, \tau_1, \tau_2) \\ \mathbf{P}(\mathbf{X}(\mathbf{x}, \mathbf{p}-\hbar\mathbf{k}, \tau_2, \tau_3), \mathbf{P}(\mathbf{x}, \mathbf{p}-\hbar\mathbf{k}, \tau_2, \tau_3)+\hbar\mathbf{k}, \tau_1, \tau_2)-\hbar\mathbf{k} \end{array} \right\} \\ - \frac{1}{2} \left\{ \cos \left[\mathbf{k} \cdot \left(\mathbf{x} - 2\mathbf{X} \left(\mathbf{x}, \mathbf{p} - \frac{\hbar\mathbf{k}}{2}, \tau_2, \tau_3 \right) + \boldsymbol{\xi} \right) + \tilde{\phi}_Q \left(\mathbf{x}, \mathbf{p} - \frac{\hbar\mathbf{k}}{2} \right) \right. \right. \\ \left. \left. - \delta_{12}^{(3)}\tau_3 + 2\delta_{12}^{(2)}\tau_2 - \delta_{12}^{(1)}\tau_1 - \phi_3 + 2\phi_2 - \phi_1 \right] \right. \\ \left. \times f(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)) \right\}_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}-\hbar\mathbf{k}/2, \tau_1, \tau_3), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}-\hbar\mathbf{k}/2, \tau_1, \tau_3)-\hbar\mathbf{k}/2}. \quad (51)$$

This density matrix element can be used to calculate any physically measured observable associated with atoms in state e . For example, one could measure the state e population given as

$$w = \int d\mathbf{x} d\mathbf{p} \rho_{ee}(\mathbf{x}, \mathbf{p}, \tau_{3+}). \quad (52)$$

The first two terms in Eq. (51) are responsible for the background signal. When substituted into Eq. (52), they yield a background contribution equal to 1/2, allowing us to write

$$w = \frac{1}{2} (1 - \tilde{w}), \quad (53)$$

where the interferometric term \tilde{w} is given by

$$\tilde{w} = \int d\mathbf{x}d\mathbf{p} \left\{ \cos \left[\mathbf{k} \cdot \left(\mathbf{x} - 2\mathbf{X} \left(\mathbf{x}, \mathbf{p} - \frac{\hbar\mathbf{k}}{2}, \tau_2, \tau_3 \right) + \boldsymbol{\xi} \right) + \tilde{\phi}_Q \left(\mathbf{x}, \mathbf{p} - \frac{\hbar\mathbf{k}}{2} \right) - \delta_{12}^{(3)}\tau_3 + 2\delta_{12}^{(2)}\tau_2 - \delta_{12}^{(1)}\tau_1 - \phi_3 + 2\phi_2 - \phi_1 \right] \right. \\ \left. \times f \left(\mathbf{X} \left(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1 \right), \mathbf{P} \left(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1 \right) \right) \right\}_{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p} - \hbar\mathbf{k}/2, \tau_1, \tau_3), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p} - \hbar\mathbf{k}/2, \tau_1, \tau_3) - \hbar\mathbf{k}/2}. \quad (54)$$

To carry out the integration we express all position and momenta in terms of the position and momentum variables at time 0, denoted by

$$\{\mathbf{x}', \mathbf{p}'\} = \{\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, 0, \tau_1)\} \quad (55)$$

In terms of these variables,

$$\{\boldsymbol{\xi}, \boldsymbol{\pi}\} = \{\mathbf{X}(\mathbf{x}', \mathbf{p}', \tau_1, 0), \mathbf{P}(\mathbf{x}', \mathbf{p}', \tau_1, 0)\}, \quad (56a)$$

$$\{\mathbf{x}, \mathbf{p}\} = \{\mathbf{X}(\mathbf{X}(\mathbf{x}', \mathbf{p}', \tau_1, 0), \mathbf{P}(\mathbf{x}', \mathbf{p}', \tau_1, 0) + \hbar\mathbf{k}/2, \tau_3, \tau_1), \\ \mathbf{P}(\mathbf{X}(\mathbf{x}', \mathbf{p}', \tau_1, 0), \mathbf{P}(\mathbf{x}', \mathbf{p}', \tau_1, 0) + \hbar\mathbf{k}/2, \tau_3, \tau_1) + \hbar\mathbf{k}/2\}, \quad (56b)$$

$$|\partial\{\mathbf{x}, \mathbf{p}\} / \partial\{\mathbf{x}', \mathbf{p}'\}| = 1, \quad (56c)$$

$$\mathbf{X}(\mathbf{x}, \mathbf{p} - \hbar\mathbf{k}/2, \tau_2, \tau_3) = \mathbf{X}(\mathbf{X}(\mathbf{x}', \mathbf{p}', \tau_1, 0), \mathbf{P}(\mathbf{x}', \mathbf{p}', \tau_1, 0) + \hbar\mathbf{k}/2, \tau_2, \tau_1). \quad (56d)$$

After redefining $\{\mathbf{x}', \mathbf{p}'\} \rightarrow \{\mathbf{x}, \mathbf{p}\}$, one finds

$$\tilde{w} = \int d\mathbf{x}d\mathbf{p} \cos \left[\phi(\mathbf{x}, \mathbf{p}) - \delta_{12}^{(3)}\tau_3 + 2\delta_{12}^{(2)}\tau_2 - \delta_{12}^{(1)}\tau_1 - \phi_3 + 2\phi_2 - \phi_1 \right] f(\mathbf{x}, \mathbf{p}), \quad (57)$$

where the phase $\phi(\mathbf{x}, \mathbf{p})$ of the AI is defined as

$$\phi(\mathbf{x}, \mathbf{p}) = \phi_r(\mathbf{x}, \mathbf{p}) + \phi_Q(\mathbf{x}, \mathbf{p}); \quad (58a)$$

$$\phi_r(\mathbf{x}, \mathbf{p}) = \mathbf{k} \cdot [\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_3, \tau_1) - 2\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, \tau_1) + \boldsymbol{\xi}]_{\{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \hbar\mathbf{k}/2\}}; \quad (58b)$$

$$\phi_Q(\mathbf{x}, \mathbf{p}) = \tilde{\phi}_Q[\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_3, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_3, \tau_1)]_{\{\boldsymbol{\xi}=\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \boldsymbol{\pi}=\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \hbar\mathbf{k}/2\}}, \quad (58c)$$

with $\tilde{\phi}_Q$ given by Eq. (50c).

1. Atom trajectories in the presence of the test mass

To calculate the phases in Eqs. (58) we need expressions for the propagation functions $\{\mathbf{X}(\mathbf{x}, \mathbf{p}, t, t'), \mathbf{P}(\mathbf{x}, \mathbf{p}, t, t')\}$, i.e. atomic position and momentum at time t subject to the initial value $\{\mathbf{x}, \mathbf{p}\}$ at time t' . These functions evolve as

$$\dot{\mathbf{X}}(\mathbf{x}, \mathbf{p}, t, t') = \frac{\mathbf{P}(\mathbf{x}, \mathbf{p}, t, t')}{M_a}, \quad (59a)$$

$$\dot{\mathbf{P}}(\mathbf{x}, \mathbf{p}, t, t') = M_a \{\mathbf{g} + \delta\mathbf{g}[\mathbf{X}(\mathbf{x}, \mathbf{p}, t, t'), t]\}. \quad (59b)$$

We neglect in Eq. (59) the gravity-gradient, centrifugal and Coriolis forces caused by the rotating Earth. When $\delta\mathbf{g}(\mathbf{x}, t)$ is a perturbation, the approximate solutions of Eqs. (59) are [31]

$$\mathbf{X}(\mathbf{x}, \mathbf{p}, t, t') \approx \mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, t') + \delta\mathbf{X}(\mathbf{x}, \mathbf{p}, t, t'), \quad (60a)$$

$$\mathbf{P}(\mathbf{x}, \mathbf{p}, t, t') \approx \mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, t, t') + \delta\mathbf{P}(\mathbf{x}, \mathbf{p}, t, t'), \quad (60b)$$

$$\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, t') = \mathbf{x} + \frac{\mathbf{p}}{M_a}(t - t') + \mathbf{g} \frac{(t - t')^2}{2}, \quad (60c)$$

$$\mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, t, t') = \mathbf{p} + M_a\mathbf{g}(t - t'); \quad (60d)$$

$$\delta\mathbf{X}(\mathbf{x}, \mathbf{p}, t, t') = \int_{t'}^t dt'' (t - t'') \delta\mathbf{g}[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t'', t'), t''] \quad (60e)$$

$$\delta\mathbf{P}(\mathbf{x}, \mathbf{p}, t, t') = M_a \int_{t'}^t dt'' \delta\mathbf{g}[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t'', t'), t'']. \quad (60f)$$

Each of the functions $\{\mathbf{X}^{(0)}, \mathbf{P}^{(0)}, \delta\mathbf{X}, \delta\mathbf{P}\}$ obeys the multiplication law (40)

$$\left\{ \begin{array}{c} \mathbf{X}^{(0)} \\ \mathbf{P}^{(0)} \end{array} \right\} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t', t''), \mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, t', t''), t, t' \right) = \left\{ \begin{array}{c} \mathbf{X}^{(0)} \\ \mathbf{P}^{(0)} \end{array} \right\}(\mathbf{x}, \mathbf{p}, t, t''); \quad (61a)$$

$$\left\{ \begin{array}{c} \delta\mathbf{X} \\ \delta\mathbf{P} \end{array} \right\} (\delta\mathbf{X}(\mathbf{x}, \mathbf{p}, t', t''), \delta\mathbf{P}(\mathbf{x}, \mathbf{p}, t', t''), t, t') = \left\{ \begin{array}{c} \delta\mathbf{X} \\ \delta\mathbf{P} \end{array} \right\}(\mathbf{x}, \mathbf{p}, t, t'). \quad (61b)$$

2. Phases

It remains for us to calculate the phases $\phi_r(\mathbf{x}, \mathbf{p})$ and $\phi_Q(\mathbf{x}, \mathbf{p})$. In the following two subsections, we obtain both exact integral and approximate integral and analytic expressions for these phases. In Sec. III, the exact expressions are evaluated numerically and the range of validity of the approximate expressions is established.

a. $\phi_r(\mathbf{x}, \mathbf{p})$ The phase ϕ_r includes a "classical" part (vanishing in the limit $\hbar \rightarrow 0$), as well as a quantum correction ϕ_q . The contributions to ϕ_r resulting from the Earth's gravitational field and the rotation of the Earth were calculated approximately in [17, 27]. The classical component of these contributions to ϕ_r has been calculated exactly [28]. In this paper we concentrate on the additions to ϕ_r caused by the test mass' field. The "classical" part of this addition has been evaluated in Ref. [24]. Contributions to the phase from the Earth's rotation is neglected.

It is shown in the appendix how approximate expressions for the propagators needed in Eq. (58b) can be obtained from Eqs. (60). It then follows that the phase ϕ_r given in Eq. (58b) can be written as a sum of three terms,

$$\phi_r(\mathbf{x}, \mathbf{p}) = \phi_0(\mathbf{x}, \mathbf{p}) + \delta\phi(\mathbf{x}, \mathbf{p}) + \phi_q(\mathbf{x}, \mathbf{p}); \quad (62a)$$

$$\phi_0(\mathbf{x}, \mathbf{p}) = \mathbf{k} \cdot \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_3, 0) - 2\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2, 0) + \mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) \right] = \mathbf{k} \cdot \mathbf{g}T^2, \quad (62b)$$

$$\delta\phi(\mathbf{x}, \mathbf{p}) = \mathbf{k} \cdot \boldsymbol{\psi} \equiv \mathbf{k} \cdot [\delta\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_3, 0) - 2\delta\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_2, 0) + \delta\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0)], \quad (62c)$$

$$\phi_q(\mathbf{x}, \mathbf{p}) = \mathbf{k} \cdot \boldsymbol{\psi}_q, \quad (62d)$$

$$\begin{aligned} \boldsymbol{\psi}_q = & \int_{\tau_1}^{\tau_3} dt (\tau_3 - t) \left\{ \delta\mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0) + \frac{\hbar\mathbf{k}}{2M_a}(t - \tau_1), t \right] - \delta\mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \right\} \\ & - 2 \int_{\tau_1}^{\tau_2} dt (\tau_2 - t) \left\{ \delta\mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0) + \frac{\hbar\mathbf{k}}{2M_a}(t - \tau_1), t \right] - \delta\mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \right\}. \end{aligned} \quad (62e)$$

The term $\phi_0(\mathbf{x}, \mathbf{p})$ is the classical contribution from the Earth's field, the term $\delta\phi(\mathbf{x}, \mathbf{p})$ is the classical contribution from the test mass' field, and the term $\phi_q(\mathbf{x}, \mathbf{p})$ is the quantum correction to the test mass' field.

To evaluate the classical contribution to the phase given by Eq. (62c), we use Eq. (60e) to arrive at

$$\boldsymbol{\psi} = \tau_3 \mathbf{u}_{20} - \tau_1 \mathbf{u}_{10} + \mathbf{u}_{11} - \mathbf{u}_{21}, \quad (63a)$$

$$\mathbf{u}_{\alpha\beta} = \int_{\tau_\alpha}^{\tau_\alpha+T} dt'' (t'')^\beta \delta\mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t'', 0), t'' \right]. \quad (63b)$$

Equations (62c, 63) have been used in Ref. [24]. With the simple change of variables, $t = \tau_2 + \theta$ for $\mathbf{u}_{2\beta}$ and $t = \tau_1 + \theta$ for $\mathbf{u}_{1\beta}$, we find

$$\boldsymbol{\psi} = \int_0^T d\theta \left\{ (T - \theta) \delta\mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0), \tau_2 + \theta \right] + \theta \delta\mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0), \tau_1 + \theta \right] \right\}. \quad (64)$$

If the test mass moves without rotation and follows a trajectory denoted by $\mathbf{x}_m(t)$, then

$$\delta\mathbf{g}(\mathbf{x}, t) = \delta\mathbf{g}[\mathbf{x} - \mathbf{x}_m(t)] \quad (65)$$

and

$$\boldsymbol{\psi} = \int_0^T d\theta \left\{ (T - \theta) \delta\mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0) - \mathbf{x}_m(\tau_2 + \theta) \right] + \theta \delta\mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0) - \mathbf{x}_m(\tau_1 + \theta) \right] \right\}. \quad (66)$$

This is the exact expression for $\boldsymbol{\psi}$ that is used in Sec. III.

We can arrive at an approximate expression for $\boldsymbol{\psi}$ if we assume that the distance between the atoms and the test mass is sufficiently large to keep only those terms that are linear in the field gradient. In other words, we can evaluate

the field of the test mass at some average displacement \mathbf{x}_C between the test mass and the atoms' trajectory. If we choose [44]

$$\mathbf{x}_C = \frac{1}{T^2} \int_0^T d\theta \left\{ (T - \theta) \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0) - \mathbf{x}_m(\tau_2 + \theta) \right] + \theta \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0) - \mathbf{x}_m(\tau_1 + \theta) \right] \right\}, \quad (67)$$

expand

$$\delta \mathbf{g}_i(\mathbf{x}) \approx \delta \mathbf{g}_i(\mathbf{x}_C) + \underline{\gamma}(\mathbf{x})(\mathbf{x} - \mathbf{x}_C), \quad (68)$$

where $\underline{\gamma}(\mathbf{x})$ is the gravity-gradient tensor having matrix elements

$$\gamma_{ij}(\mathbf{x}) = \frac{\partial \delta \mathbf{g}_i(\mathbf{x})}{\partial x_j}, \quad (69)$$

and substitute the result back into Eq. (66), we find that the term proportional to $\underline{\gamma}(\mathbf{x})$ vanishes (\mathbf{x}_C was chosen to insure this). We then obtain an approximate expression $\delta \phi_a$ for the classical contribution to the phase ϕ_r given by

$$\delta \phi \approx \delta \phi_a = \mathbf{k} \cdot \boldsymbol{\psi}_a, \quad (70a)$$

$$\boldsymbol{\psi}_a = \delta \mathbf{g}(\mathbf{x}_C) T^2. \quad (70b)$$

We now turn our attention to the quantum correction. The vector $\boldsymbol{\psi}_q$ given in Eq. (62d) can be rewritten as

$$\begin{aligned} \boldsymbol{\psi}_q = & \int_{\tau_2}^{\tau_3} dt (\tau_3 - t) \left\{ \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0) + \frac{\hbar \mathbf{k}}{2M_a} (t - \tau_1), t \right] - \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \right\} \\ & + \int_{\tau_1}^{\tau_2} dt (t - \tau_1) \left\{ \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0) + \frac{\hbar \mathbf{k}}{2M_a} (t - \tau_1), t \right] - \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \right\}. \end{aligned} \quad (71)$$

Substituting $t = \tau_2 + \theta$ in the first term of Eq. (71) and $t = \tau_1 + \theta$ in the second term, we obtain

$$\begin{aligned} \boldsymbol{\psi}_q = & \int_0^T d\theta \left\{ (T - \theta) \left[\delta \mathbf{g} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0) + \frac{\hbar \mathbf{k}}{2M_a} (T + \theta), \tau_2 + \theta \right) - \delta \mathbf{g} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0), \tau_2 + \theta \right) \right] \right. \\ & \left. + \theta \left[\delta \mathbf{g} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0) + \frac{\hbar \mathbf{k}}{2M_a} \theta, \tau_1 + \theta \right) - \delta \mathbf{g} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0), \tau_1 + \theta \right) \right] \right\}. \end{aligned} \quad (72)$$

For translational motion, when the gravitational field of the test mass is given by Eq. (65),

$$\begin{aligned} \boldsymbol{\psi}_q = & \int_0^T d\theta \left\{ (T - \theta) \left[\delta \mathbf{g} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0) + \frac{\hbar \mathbf{k}}{2M_a} (T + \theta) - \mathbf{x}_m(\tau_2 + \theta) \right) - \delta \mathbf{g} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0) - \mathbf{x}_m(\tau_2 + \theta) \right) \right] \right. \\ & \left. + \theta \left[\delta \mathbf{g} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0) + \frac{\hbar \mathbf{k}}{2M_a} \theta - \mathbf{x}_m(\tau_1 + \theta) \right) - \delta \mathbf{g} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0) - \mathbf{x}_m(\tau_1 + \theta) \right) \right] \right\}, \end{aligned} \quad (73)$$

This is the exact expression for $\boldsymbol{\psi}_q$ that is used in Sec. III.

There are two approximate expressions we will derive for $\boldsymbol{\psi}$. When the recoil effect is small,

$$\frac{\hbar k}{2M_a} T \ll X^{(0)}(\mathbf{x}, \mathbf{p}, T, 0), \quad (74)$$

we can expand the arguments in Eq. (73) to obtain a first approximation $\phi_q \approx \phi_{qn}$ given by

$$\phi_q \approx \phi_{qn} = \mathbf{k} \cdot \boldsymbol{\psi}_{qn} \quad (75a)$$

$$\boldsymbol{\psi}_{qn} = \int_0^T d\theta \left\{ (T^2 - \theta^2) \underline{\gamma} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0), \tau_2 + \theta \right] + \theta^2 \underline{\gamma} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0), \tau_1 + \theta \right] \right\} \frac{\hbar \mathbf{k}}{2M_a}, \quad (75b)$$

where

$$\gamma_{ij}(\mathbf{x}, t) = \frac{\partial \delta \mathbf{g}_i(\mathbf{x}, t)}{\partial x_j}. \quad (76)$$

For translational motion of the test mass, $\underline{\gamma}(\mathbf{x}, t) = \underline{\gamma}[\mathbf{x} - \mathbf{x}_m(t)]$ and Eq. (75b) reduces to

$$\psi_{qn} = \int_0^T d\theta \left\{ (T^2 - \theta^2) \underline{\gamma} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0) - \mathbf{x}_m(\tau_2 + \theta) \right] + \theta^2 \underline{\gamma} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0) - \mathbf{x}_m(\tau_1 + \theta) \right] \right\} \frac{\hbar \mathbf{k}}{2M_a}. \quad (77)$$

The second approximate expression we obtain for ψ_q is the limit of Eq. (77) when the distance between the atoms and the test mass is sufficiently large to keep only those terms that are linear in the field gradient. If we choose

$$\mathbf{x}_{qC} = \frac{1}{T^3} \int_0^T d\theta \left\{ (T^2 - \theta^2) \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0) - \mathbf{x}_m(\tau_2 + \theta) \right] + \theta^2 \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0) - \mathbf{x}_m(\tau_1 + \theta) \right] \right\}, \quad (78)$$

and expand

$$\underline{\gamma}_{ij}(\mathbf{x}) \approx \underline{\gamma}_{ij}(\mathbf{x}_{qC}) + \chi_{ijl}(\mathbf{x}_{qC})(\mathbf{x} - \mathbf{x}_{qC})_l, \quad (79)$$

where

$$\chi_{ijl}(\mathbf{x}) = \partial_{\mathbf{x}_l} \underline{\gamma}_{ij}(\mathbf{x}) \quad (80)$$

is an element of the gravity curvature tensor, then the contribution from the second term in Eq. (79) vanishes and we find an approximate expression ϕ_{qa} for the phase given by

$$\phi_q(\mathbf{x}, \mathbf{p}) \approx \phi_{qa}(\mathbf{x}, \mathbf{p}) = \mathbf{k} \cdot \psi_{qa}, \quad (81a)$$

$$\psi_{qa} = \underline{\gamma}(\mathbf{x}_{qC}) \frac{\hbar \mathbf{k}}{2M_a} T^3. \quad (81b)$$

b. $\phi_Q(\mathbf{x}, \mathbf{p})$ We now consider Q -term quantum corrections to the phase given by Eqs. (58c, 50c). We first replace $\{\mathbf{x}, \mathbf{p}\}$ by $\{\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_3, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_3, \tau_1)\}$ in Eq. (50c) to obtain

$$\left\{ \begin{array}{c} \boldsymbol{\xi}' \\ \boldsymbol{\pi}' \end{array} \right\} = \left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}(\mathbf{X}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_3, \tau_1), \mathbf{P}(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_3, \tau_1), t, \tau_3) = \left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}(\boldsymbol{\xi}, \boldsymbol{\pi}, t, \tau_1), \quad (82)$$

allowing us to write $\phi_Q(\mathbf{x}, \mathbf{p})$ as

$$\begin{aligned} \phi_Q(\mathbf{x}, \mathbf{p}) = & -\frac{\hbar^2}{24} k_u k_v k_w \left\{ \int_{\tau_1}^{\tau_2} dt \chi'_{ikl}(\boldsymbol{\xi}, t) \partial_{\pi_i} \mathbf{X}_u(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \partial_{\pi_k} \mathbf{X}_v(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \partial_{\pi_l} \mathbf{X}_w(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t) \right. \\ & + \int_{\tau_2}^{\tau_3} dt \chi'_{ikl}(\boldsymbol{\xi}, t) \left[\frac{\partial \mathbf{X}_u(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)}{\partial \pi_i} - 2 \frac{\partial \mathbf{X}_u(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, t)}{\partial \pi_i} \right] \left[\frac{\partial \mathbf{X}_v(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)}{\partial \pi_k} - 2 \frac{\partial \mathbf{X}_v(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, t)}{\partial \pi_k} \right] \\ & \left. \times \left[\frac{\partial \mathbf{X}_w(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_1, t)}{\partial \pi_l} - 2 \frac{\partial \mathbf{X}_w(\boldsymbol{\xi}, \boldsymbol{\pi}, \tau_2, t)}{\partial \pi_l} \right] \right\} \left\{ \begin{array}{c} \boldsymbol{\xi} \\ \boldsymbol{\pi} \end{array} \right\} = \left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}_{(\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \hbar \mathbf{k}/2, t, \tau_1)}. \quad (83) \end{aligned}$$

When atoms move between the Raman pulses under the action of the homogeneous gravitational field \mathbf{g} of the Earth and the inhomogeneous perturbation $\delta \mathbf{g}(\mathbf{x}, t)$ caused by the test mass, the only contribution to $\chi'_{ikl}(\boldsymbol{\xi}, t)$ [defined in Eq. (19)] results from the presence of the test mass,

$$\chi'_{ikl}(\mathbf{x}, t) = M_a \chi_{ikl}(\mathbf{x}, t), \quad (84a)$$

where

$$\chi_{ijl}(\mathbf{x}, t) = \partial_{\mathbf{x}_l} \gamma_{ij}(\mathbf{x}, t). \quad (85)$$

Since we calculate the AI phase to first order in $\delta \mathbf{g}$, it is sufficient to calculate the atom trajectory in Eq. (83) to zeroth order in $\delta \mathbf{g}$, i.e. to set

$$\left\{ \begin{array}{c} \mathbf{X} \\ \mathbf{P} \end{array} \right\}(\mathbf{x}, \mathbf{p}, t, t') = \left\{ \begin{array}{c} \mathbf{X}^{(0)} \\ \mathbf{P}^{(0)} \end{array} \right\}(\mathbf{x}, \mathbf{p}, t, t'), \quad (86)$$

which results in

$$\partial_{p_i} \mathbf{X}_j^{(0)}(\mathbf{x}, \mathbf{p}, t, t') = \frac{\delta_{ij}}{M_a} (t - t'), \quad (87)$$

where δ_{ij} is a Kronecker delta. Moreover, since we are interested in calculating $\phi_Q(\mathbf{x}, \mathbf{p})$ to second order in the recoil momentum $\hbar\mathbf{k}$, we can neglect the contribution of the recoil term in the braces of Eq. (83). We then apply the multiplication law (40) and obtain

$$\phi_Q(\mathbf{x}, \mathbf{p}) = \frac{\hbar^2}{24M_a^2} k_i k_j k_l \left\{ \int_{\tau_1}^{\tau_2} dt \chi_{ijl} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] (t - \tau_1)^3 + \int_{\tau_2}^{\tau_3} dt \chi_{ijl} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] (\tau_3 - t)^3 \right\} \quad (88)$$

As before, we transform the integral to one from 0 to T ,

$$\phi_Q(\mathbf{x}, \mathbf{p}) = \frac{\hbar^2}{24M_a^2} k_i k_j k_l \int_0^T d\theta \left\{ \theta^3 \chi_{ijl} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0), \tau_1 + \theta \right] + (T - \theta)^3 \chi_{ijl} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0), \tau_2 + \theta \right] \right\}. \quad (89)$$

If the test mass moves without rotation, then

$$\chi_{ikl}(\mathbf{x}, t) = \chi_{ikl}[\mathbf{x} - \mathbf{x}_m(t)] \quad (90)$$

and

$$\phi_Q(\mathbf{x}, \mathbf{p}) = \frac{\hbar^2}{24M_a^2} k_i k_j k_l \int_0^T d\theta \left\{ \begin{array}{l} \theta^3 \chi_{ijl} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0) - \mathbf{x}_m(\tau_1 + \theta) \right] \\ + (T - \theta)^3 \chi_{ijl} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0) - \mathbf{x}_m(\tau_2 + \theta) \right] \end{array} \right\}. \quad (91)$$

This is the exact expression for ψ_q that is used in Sec. III.

We can obtain an approximate expression for $\phi_Q(\mathbf{x}, \mathbf{p})$ when the distance between the atoms and the test mass is sufficiently large to keep only those terms that are linear in the field curvature. If we choose

$$\mathbf{x}_{QC} = \frac{2}{T^4} \int_0^T d\theta \left\{ \theta^3 \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1 + \theta, 0) - \mathbf{x}_m(\tau_1 + \theta) \right] + (T - \theta)^3 \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_2 + \theta, 0) - \mathbf{x}_m(\tau_2 + \theta) \right] \right\} \quad (92)$$

and expand

$$\chi_{ijl}(\mathbf{x}) \approx \chi_{ijl}(\mathbf{x}_{QC}) + (\partial_{\mathbf{x}_C})_m \chi_{ijl}(\mathbf{x}_{QC})(\mathbf{x} - \mathbf{x}_{QC})_m, \quad (93)$$

the contribution to the Q-term from the second term in Eq. (93) vanishes and we find an approximate expression ϕ_{Qa} for the phase given by

$$\phi_Q(\mathbf{x}, \mathbf{p}) \approx \phi_{Qa}(\mathbf{x}, \mathbf{p}) = \frac{\hbar^2}{48M_a^2} k_i k_j k_l \chi_{ijl}(\mathbf{x}_{QC}) T^4. \quad (94)$$

III. POINT SOURCE TEST MASS

For a point source test mass M moving along the trajectory $\mathbf{x}_m(t)$, the gravitational field, gravity-gradient tensor, and gravity curvature tensor are given by

$$\delta\mathbf{g}(\mathbf{x}, t) = \delta\mathbf{g}[\mathbf{x} - \mathbf{x}_m(t)]; \quad \delta\mathbf{g}(\mathbf{x}) = -GM \frac{\mathbf{x}}{x^3}, \quad (95a)$$

$$\gamma_{jl}(\mathbf{x}, t) = \gamma_{jl}[\mathbf{x} - \mathbf{x}_m(t)]; \quad \gamma_{jl}(\mathbf{x}) = -GM \left(\frac{\delta_{jl}}{x^3} - 3 \frac{x_j x_l}{x^5} \right), \quad (95b)$$

$$\chi_{ijl}(\mathbf{x}, t) = \chi_{ijl}[\mathbf{x} - \mathbf{x}_m(t)], \quad (95c)$$

$$\chi_{ijl}(\mathbf{x}) = GM \left[\frac{3}{x^5} (\delta_{jl} x_i + \delta_{il} x_j + \delta_{ij} x_l) - 15 \frac{x_i x_j x_l}{x^7} \right], \quad (95d)$$

where $G = 6.67428 \times 10^{-11} \text{ m}^3/\text{kg}\cdot\text{s}^2$ is the Newtonian gravitational constant.

We numerically calculated the classical part of the phase given by Eqs. (62c, 66) (for $\delta\phi$) and the quantum corrections given by Eqs. (62d, 73) (for ϕ_q) and (91) (for ϕ_Q) and determined when these terms become measurable in the presence of phase noise given by Eq. (7). Moreover, we checked the validity of the approximate expressions given in Eqs. (70, 67), (77), (81, 78), and (94, 92). The results vary with the test mass' weight, shape, trajectory, as well as with the operating parameters of the atom interferometer.

The calculations are carried out for a test mass moving with constant velocity \mathbf{v}_m ,

$$\mathbf{x}_m(t) = \mathbf{x}_{m0} + \mathbf{v}_m t, \quad (96)$$

where

$$\mathbf{x}_{m0} = (x_{m0}, y_{m0}, z_{m0}). \quad (97)$$

is the location of the test mass at time $t = 0$.

We assume that, at $t = 0$, the atoms are launched from the origin of the North-East-Down frame in the vertical direction. That is, it is assumed that \mathbf{g} is in the positive z direction, and that the cloud position is given by

$$z(t) = v_0 t + \frac{1}{2} g t^2; \quad (98a)$$

$$x(t) = y(t) = 0, \quad (98b)$$

where $\mathbf{v}_0 = v_0 \mathbf{u}_z$ is the launch velocity, taken to be along the z -axis. In this case one finds from Eqs. (60c, 67, 78, 92) that

$$\mathbf{x}_C = -\mathbf{x}_{m0} + [\mathbf{v}_0 - \mathbf{v}_m] (\tau_1 + T) + \frac{1}{2} \mathbf{g} \left(\tau_1^2 + 2\tau_1 T + \frac{7}{6} T^2 \right), \quad (99a)$$

$$\mathbf{x}_{qC} = -\mathbf{x}_{m0} + [\mathbf{v}_0 - \mathbf{v}_m] \left(\tau_1 + \frac{7}{6} T \right) + \frac{1}{2} \mathbf{g} \left(\tau_1^2 + \frac{7}{3} \tau_1 T + \frac{3}{2} T^2 \right), \quad (99b)$$

$$\mathbf{x}_{QC} = -\mathbf{x}_{m0} + [\mathbf{v}_0 - \mathbf{v}_m] (\tau_1 + T) + \mathbf{g} \left(\frac{\tau_1^2}{2} + \tau_1 T + \frac{8}{15} T^2 \right). \quad (99c)$$

The launch velocity is chosen as

$$\mathbf{v}_0 = -\mathbf{g} (\tau_1 + T), \quad (100)$$

corresponding to a symmetric fountain geometry in which the atomic cloud reaches its highest point at time τ_2 , when the second pulse is applied. Calculations have been performed for a stationary test mass having

$$\mathbf{x}_m^{(1)}(t) = \mathbf{x}_{m0} = (0, y_{m0}, z_{m0}) \quad (101)$$

and a test mass moving with constant velocity $\mathbf{v}_m = (5 \text{ m/s}, 0, 0)$ along the x axis,

$$\mathbf{x}_m^{(2)}(t) = \mathbf{x}_{m0} + \mathbf{v}_m t = (x_{m0}, y_{m0}, z_{m0}) + (5 \text{ m/s}, 0, 0) t. \quad (102)$$

The parameters characterizing the atom interferometer, the test mass, and the Earth's field are summarized in Table I. Since the cloud trajectory and the effective wave vectors are vertical, y_{m0} can be considered as an "impact parameter"

Earth's gravitational field	$\mathbf{g} = \{0, 0, 9.8 \text{ m/s}^2\}$
Multiple- $\hbar k$ beam splitter factor	$n_k = 25$
Effective wave vector	$\mathbf{k} = \{0, 0, -k\}$, $k = 4.0275 \times 10^8 \text{ m}^{-1}$
Time between launch and first Raman pulse	$\tau_1 = 10 \text{ ms}$
Time between Raman pulses	$T = 200 \text{ ms}$
Launch velocity	$\mathbf{v}_0 = -\mathbf{g} (\tau_1 + T)$
Error of atom interferometer phase measurement	$\phi_{err} = 10^{-3} \text{ rad}$
test mass	$M = 50 \text{ kg}$
atomic mass	87

TABLE I: Parameters of the atom interferometer and gravitational sources.

for the test mass relative to the cloud trajectory along the z -axis

Equations (95) can be used either for a point mass or a spherical mass having constant density ρ and radius

$$y_{\min} = \left[\frac{3M}{4\pi\rho} \right]^{1/3}. \quad (103)$$

For the highest density in nature, $\rho = 22600 \text{ kg/m}^3$, corresponding to osmium [45],

$$y_{\min} \approx 0.0808 \text{ m}. \quad (104)$$

For impact parameters $y_{m0} < y_{\min}$ Eqs. (95) are valid only for those values of z_{m0} for which atom trajectory does not intersect the spherical test mass. In the case of stationary sphere, this requirement translates into one in which the distance between the cloud and the sphere is always greater than y_{\min} ; that is, for any $t > 0$,

$$y_{m0}^2 + [z(t) - z_{m0}]^2 > y_{\min}^2. \quad (105)$$

With $z(t)$ given by Eq. (98a) and \mathbf{v}_0 given by Eq. (100), we can show that this inequality is satisfied if

$$-\sqrt{y_{\min}^2 - y_{m0}^2} - \frac{1}{2}g(\tau_1 + T)^2 > z_{m0}, \text{ or } z_{m0} > \sqrt{y_{\min}^2 - y_{m0}^2}. \quad (106)$$

For the moving sphere the range of allowed initial positions (x_{m0}, y_{m0}, z_{m0}) for the center of the sphere is more difficult to calculate. For this reason, in the case 2 we consider only impact parameters larger than sphere's radius, $y_{m0} > y_{\min}$, for which, evidently, any values of (x_{m0}, z_{m0}) are allowed.

For each impact parameter y_{m0} and for the parameters given in Table I, we explore various test mass trajectories and obtain:

1. maximum of the magnitude of the phase $|\delta\phi|_{\max}$ obtained from Eqs. (62c, 66);
2. maximum of the magnitude of the phase difference $|\delta\phi - \delta\phi_a|_{\max}$ obtained from Eqs. (62c, 66; 70, 99a);
3. the maximum of the magnitude of the quantum correction $|\phi_q|_{\max}$ obtained from Eqs. (62d, 73);
4. maximum of the magnitude of the phase difference $|\phi_q - \phi_{qn}|_{\max}$ obtained from Eqs. (62d, 73; 75a, 77);
5. maximum of the magnitude of the phase difference $|\phi_q - \phi_{qa}|_{\max}$ obtained from Eqs. (62d, 73; 81, 99b);
6. maximum of the magnitude of the quantum correction $|\phi_Q|_{\max}$ obtained from Eq. (91);
7. maximum of the magnitude of the phase difference $|\phi_Q - \phi_{Qa}|_{\max}$ obtained from Eqs. (91; 94, 99c).

The results of the calculations are illustrated graphically in Figs. 1, 2. From these figures, one can construct regions where the various approximate expressions for the phases are valid. The impact parameters corresponding to the various regions are given in Table II and the validity conditions for the various approximate expressions for the phases in these regions is listed below.

Region	stationary test mass	test mass moving with constant velocity
1	$y_{m0} < 0.166 \text{ m}$	$y_{m0} < 0.250 \text{ m}$
2	$y_{m0} > 0.292 \text{ m}$	$y_{m0} > 0.249 \text{ m}$
3	$y_{m0} < 0.473 \text{ m}$	$y_{m0} < 0.407 \text{ m}$
4	$y_{m0} > 0.488 \text{ m}$	$y_{m0} > 0.732 \text{ m}$
5	$y_{m0} < 0.525 \text{ m}$	$y_{m0} < 1.34 \text{ m}$
6	$y_{m0} > 1.16 \text{ m}$	$y_{m0} > 1.11 \text{ m}$
7	$y_{m0} > 4.55 \text{ m}$	$y_{m0} > 4.53 \text{ m}$

TABLE II: Locations of regions of validity of the exact and approximate expressions for the stationary and moving test mss. The regions refer to regions 1-7 given in the text.

- Region 1. One should use the exact expression, Eq. (91), for ϕ_Q in this region; only outside this region are the approximate expressions given by Eqs. (94, 99c) valid (see plots g_1 in the figures);
- Region 2. The phase ϕ_Q is negligible in this region (see plots f_1 in the figures);
- Region 3. One should use the exact expressions, Eqs. (62d, 73), for the quantum correction ϕ_q in this region; only outside this region does the approximate expression given by Eqs. (75a, 77) become valid (see plots d_1 in the figures);
- Region 4. One can use the approximate expressions for ϕ_{qa} given by Eqs. (81, 99b) in this region (see plots e_1 in the figures);

- Region 5. One should use the exact expressions, Eqs. (62c, 66), for the classical part of the phase $\delta\phi$; only outside this region does the approximate expression given by Eqs. (70, 99a) become valid (see plots b_1 in the figures);
- Region 6. The phase ϕ_q is negligible in this region (see plots c_1 in the figures);
- Region 7. The phase $\delta\phi$ produced by the test mass falls below the phase noise ϕ_{err} , so the effect of the test mass cannot be measured in this region (see plots a_1 in the figures).

These numerical calculations show that the approximate expressions that were obtained based on assumptions about the approximate homogeneity of the field in Ref. [44] for the classical contributions to the phase or on the approximate homogeneity of the field gradient and curvature for the quantum corrections to the phase become valid only in regions where the phases are smaller by 1-2 orders of magnitude the maximum values for the phases that occur in regions where the field inhomogeneity plays an important role.

By using an effective wave vector that is 25 times larger than the wave vector of a tow-field Raman pulse, we reach a limit where the quantum correction ϕ_q is comparable with the classical part of the phase, while the Q -term is still small. Further increase of the effective wave vector or time interval T between the pulses could bring us to situation when quantum correction dominate over the classical part of the phase.

Appendix A:

In this appendix, we show how we arrive at Eqs. (62) from Eq. (58b) and Eqs (60).

To calculate the phase $\phi_r(\mathbf{x}, \mathbf{p})$, we need an approximate expression for the propagator $\mathbf{X}\left(\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar\mathbf{k}}{2}, \tau_s, \tau_1\right)$. Using Eqs. (60, 61) we find

$$\begin{aligned}
\mathbf{X}\left(\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar\mathbf{k}}{2}, \tau_s, \tau_1\right) &\approx \mathbf{X}^{(0)}\left(\begin{array}{c} \mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \delta\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \\ \mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \delta\mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar\mathbf{k}}{2}, \tau_s, \tau_1 \end{array}\right) \\
&+ \delta\mathbf{X}\left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar\mathbf{k}}{2}, \tau_s, \tau_1\right) \\
&= \mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \delta\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \\
&\frac{1}{M_a}\left[\mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + M_a \int_0^{\tau_1} dt \delta\mathbf{g}\left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t\right] + \frac{\hbar\mathbf{k}}{2}\right](\tau_s - \tau_1) + \frac{1}{2}\mathbf{g}(\tau_s - \tau_1)^2 \\
&+ \int_{\tau_1}^{\tau_s} dt (\tau_s - t) \delta\mathbf{g}\left[\mathbf{X}^{(0)}\left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar\mathbf{k}}{2}, t, \tau_1\right), t\right], \tag{A1}
\end{aligned}$$

From Eqs. (61, 60c), it follows that

$$\begin{aligned}
\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_s, 0) &= \mathbf{X}^{(0)}\left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \tau_s, \tau_1\right) \\
&= \mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{1}{M_a}\mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0)(\tau_s - \tau_1) + \frac{1}{2}\mathbf{g}(\tau_s - \tau_1)^2, \tag{A2}
\end{aligned}$$

allowing us to rewrite

$$\begin{aligned}
\mathbf{X}\left(\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar\mathbf{k}}{2}, \tau_s, \tau_1\right) &\approx \mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_s, 0) + \frac{\hbar\mathbf{k}}{2M_a}(\tau_s - \tau_1) \\
&+ (\tau_s - \tau_1) \int_0^{\tau_1} dt \delta\mathbf{g}\left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t\right] + \int_0^{\tau_1} dt (\tau_1 - t) \delta\mathbf{g}\left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t\right] \\
&+ \int_{\tau_1}^{\tau_s} dt (\tau_s - t) \delta\mathbf{g}\left[\mathbf{X}^{(0)}\left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar\mathbf{k}}{2}, t, \tau_1\right), t\right]. \tag{A3}
\end{aligned}$$




FIG. 1: Stationary source. Plot a_1 , maximum of the phase magnitude given in Eqs. (62c, 66) versus the impact parameter y_{m0} ; Plot a_2 , the same as plot a_1 but for $y_{m0} < y_{min}$; Plot a_3 , z -dependence of the exact (black curve) and approximate (red curve) phases at $\{x_{m0}, y_{m0}\} = \{0, 0\}$, where the phase achieves its maximum magnitude, Plot a_4 , the same as plot a_3 but at $\{x_{m0}, y_{m0}\} = \{0, 4.55\}$ m, where phase first becomes smaller than ϕ_{err} . Plots $\{b_1, b_2\}$, the same as plot $\{a_1, a_2\}$ but for the maximum magnitude of the difference between exact and approximate phases given in Eqs. (62c, 66) and (70, 99a); Plot b_3 , the same as plot a_3 , but at the point $\{x_{m0}, y_{m0}\} = \{0, 0.525\}$ m, where the magnitude of difference shown on plot b_1 first becomes smaller than ϕ_{err} ; Plot b_4 , difference between black and red curves in plot b_3 . Plots $\{c_1 - c_4\}$, the same as plots $\{a_1 - a_4\}$ but for the magnitude of the quantum correction given in Eqs. (62d, 73); the values of $\{x_{m0}, y_{m0}\}$ are $\{0, 0\}$ for c_3 and $\{0, 1.16\}$ m for c_4 ; exact quantum correction ϕ_q , approximations ϕ_{qn} and ϕ_{qa} are shown in black, red and blue, respectively. Plots $\{d_1 - d_4\}$, the same as plots $\{b_1 - b_4\}$ but for the maximum magnitude of the difference between exact and approximate quantum corrections given in Eqs. (62d, 73) and (75a, 77); $\{x_{m0}, y_{m0}\} = \{0, 0.473\}$ m in plot d_3 where the magnitude of difference shown on plot d_1 first becomes smaller than ϕ_{err} . Plots $\{e_1 - e_4\}$, the same as plots $\{d_1 - d_4\}$ but for the maximum magnitude of the difference between exact and approximate quantum corrections given in Eqs. (62d, 73) and (81, 99b); $\{x_{m0}, y_{m0}\} = \{0, 0.488\}$ m in plot e_3 . Plots $\{f_1 - f_4\}$, the same as plots $\{a_1 - a_4\}$ but for the magnitude of the Q -term given in Eq. (91), values of $\{x_{m0}, y_{m0}\}$ are $\{0, 0\}$ for f_3 and $\{x_{m0}, y_{m0}\} = \{0, 0.292\}$ m for f_4 . Plots $\{g_1 - g_4\}$, the same as plots $\{b_1 - b_4\}$ but for the maximum magnitude of the difference between exact and approximate Q -terms given in Eqs. (91) and (94, 99c); $\{x_{m0}, y_{m0}\} = \{0, 0.166\}$ m in plot g_3 , where the magnitude of the difference shown on plot g_1 first becomes smaller than ϕ_{err} .




FIG. 2: Test mass moving with constant velocity 5 m/s. Plots α_1 are the same as plots α_1 in Fig. 1, plots α_2, α_3 are the same as α_3, α_4 in Fig. 1, where $\alpha = a - g$. Values of $\{x_{m0}, y_{m0}\}$ are $\{-1.05 \text{ m}, y_{\min}\}$ for plots a_2, f_2 , $\{-1.05 \text{ m}, 4.53 \text{ m}\}$ for plot a_3 , $\{-1.05, 1.34\}$ for plots b_2 , $\{-1.13 \text{ m}, y_{\min}\}$ for plot c_2 , $\{-1.22 \text{ m}, 1.11 \text{ m}\}$ for plot c_3 , $\{-1.33 \text{ m}, 0.407 \text{ m}\}$ for plot d_2 , $\{-1.21 \text{ m}, 0.732 \text{ m}\}$ for plot e_2 , $\{-1.05 \text{ m}, 0.249 \text{ m}\}$ for plot f_3 , and $\{-1.05 \text{ m}, 0.250 \text{ m}\}$ for plot g_2 .

Using the relationship

$$\begin{aligned} \mathbf{X}^{(0)} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar \mathbf{k}}{2}, t, \tau_1 \right) &= \frac{\hbar \mathbf{k}}{2M_a} (t - \tau_1) \\ &+ \mathbf{X}^{(0)} \left(\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_1, 0), t, \tau_1 \right) \\ &= \mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0) + \frac{\hbar \mathbf{k}}{2M_a} (t - \tau_1), \end{aligned} \quad (\text{A4})$$

we can write

$$\begin{aligned} \mathbf{X} \left(\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar \mathbf{k}}{2}, \tau_s, \tau_1 \right) &\approx \mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_s, 0) + \frac{\hbar \mathbf{k}}{2M_a} (\tau_s - \tau_1) \\ &+ (\tau_s - \tau_1) \int_0^{\tau_1} dt \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] + \int_0^{\tau_1} dt (\tau_1 - t) \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \\ &+ \int_{\tau_1}^{\tau_s} dt (\tau_s - t) \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \\ &+ \int_{\tau_1}^{\tau_s} dt (\tau_s - t) \left\{ \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0) + \frac{\hbar \mathbf{k}}{2M_a} (t - \tau_1), t \right] - \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \right\}. \end{aligned} \quad (\text{A5})$$

The propagator (60e) can be expressed as

$$\begin{aligned} \delta \mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_s, 0) &\equiv \int_0^{\tau_s} dt (\tau_s - t) \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \\ &= \int_{\tau_1}^{\tau_s} dt (\tau_s - t) \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] + \int_0^{\tau_1} dt (\tau_s - t) \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \\ &= \int_{\tau_1}^{\tau_s} dt (\tau_s - t) \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] + \int_0^{\tau_1} dt (\tau_1 - t) \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \\ &+ (\tau_s - \tau_1) \int_0^{\tau_1} dt \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right], \end{aligned} \quad (\text{A6})$$

which coincides with the sum of the 3rd, 4th and 5th terms on the right-hand-side (rhs) of Eq. (A5) and reduces to

$$\begin{aligned} \mathbf{X} \left(\mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_1, 0), \mathbf{P}(\mathbf{x}, \mathbf{p}, \tau_1, 0) + \frac{\hbar \mathbf{k}}{2}, \tau_s, \tau_1 \right) &\approx \mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, \tau_s, 0) + \frac{\hbar \mathbf{k}}{2M_a} (\tau_s - \tau_1) + \delta \mathbf{X}(\mathbf{x}, \mathbf{p}, \tau_s, 0) \\ &+ \int_{\tau_1}^{\tau_s} dt (\tau_s - t) \left\{ \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0) + \frac{\hbar \mathbf{k}}{2M_a} (t - \tau_1), t \right] - \delta \mathbf{g} \left[\mathbf{X}^{(0)}(\mathbf{x}, \mathbf{p}, t, 0), t \right] \right\}. \end{aligned} \quad (\text{A7})$$

The first term on the right hand side of this equation is responsible for the phase produced by the Earth's gravitational field. The second term corresponds to the recoil correction to the first term, but this contribution vanishes when Eq. (A7) is substituted into Eq. (58b); there is no quantum correction in an homogeneous field. The third term is responsible for the classical part of the phase produced by the test mass while the fourth term is the recoil correction to the third term.

Substituting this result in the brackets of Eq. (58b) for the 1st ($s = 3$) and 2nd ($s = 2$) terms, we arrive at Eqs. (62).

Acknowledgments

The authors would like to thank Drs. B. Young, M. Matthews, T. Loftus, V. Sonnad and A. Zorn for helpful discussions. This work was performed under the auspices of the Department of Homeland Security, Defense Nuclear Detection Office, Exploratory Research, CFP11-100-RTA-06-FP007, Contract HSHQDC-11-X-00550 by Lawrence Livermore National Laboratory, and the Defense Threat Reduction Agency by AOSense, Inc., under Contract HDTRA1-13-C-0047. Lawrence Livermore National Laboratory is operated by Lawrence Livermore National Security, LLC, for the U.S. Department of Energy, National Nuclear Security Administration under Contract DE-AC52-07NA27344

Author Contributions

S. Libby and B. Dubetsky considered the recoil term and carried out numerical calculations of the classical and recoil contributions to the phase. M. Shverdin and B. Dubetsky derived approximate expressions for the phases and carried out numerical calculations for multi- $\hbar k$ momenta transfer. P. Berman and B. Dubetsky derived and interpreted the various expressions for the phases and, together with M. Shverdin, wrote the manuscript.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

-
- [1] Dubetsky, B.; Kazantsev, A. P.; Chebotayev, V. P.; Yakovlev, V. P. Interference of atoms and formation of atomic spatial arrays in light fields. *Pis'ma Zh. Eksp. Teor. Fiz.* **1984**, *39*, 531-533 [*JETP Lett.* **1984**, *39*, 649-651].
- [2] Weiss, D. S.; Young, B. C.; Chu, S. Precision Measurement of the Photon Recoil of an Atom Using Atomic Interferometry. *Phys. Rev. Lett.* **1993**, *70*, 2706-2709.
- [3] Estey, B.; Yu, Ch.; Müller, H.; Kuan, P.-C.; Lan, S.-Y. High-resolution atom interferometers with suppressed diffraction phases. <http://arxiv.org/abs/1410.8486>.
- [4] Fixler, J. B.; Foster, G. T.; McGuirk, J. M.; Kasevich, M. A. Atom Interferometer Measurement of the Newtonian Constant of Gravity. *Science* **2007**, *315*, 74-77.
- [5] Rosi, G.; Sorrentino, F.; Cacciapuoti, L.; Prevedelli, M.; Tino, G. M. Precision measurement of the Newtonian gravitational constant using cold atoms. *Nature* **2014**, *510*, 518-521.
- [6] Kasevich, M.; Chu, S. Atomic Interferometry Using Stimulated Raman Transitions. *Phys. Rev. Lett.* **1991**, *67*, 181-184.
- [7] Cahn, S. B.; Kumarakrishnan, A.; Shim, U.; Sleator, T.; Berman, P. R.; Dubetsky, B. Time-Domain de Broglie Wave Interferometry. *Phys. Rev. Lett.* **1997**, *79*, 784-787.
- [8] Peters, A.; Chung, K. Y.; Chu, S. Measurement of gravitational acceleration by dropping atoms. *Nature* **1999**, *400*, 849-852.
- [9] Mok, C.; Barrett, B.; Carew, A.; Berthiaume, R.; Beattie, S.; Kumarakrishnan, A. Demonstration of improved sensitivity of echo interferometers to gravitational acceleration. *Phys. Rev. A* **2013**, *88*, 023614.
- [10] Snadden, M. J.; McGuirk, J. M.; Bouyer, P.; Haritos, K. G.; Kasevich, M. A. Measurement of the Earth's Gravity Gradient with an Atom Interferometer-Based Gravity Gradiometer. *Phys. Rev. Lett.* **1998**, *81*, 971-974.
- [11] McGuirk, J. M.; Foster, G. T.; Fixler, J. B.; Snadden, M. J.; Kasevich, M. A. Sensitive absolute-gravity gradiometry using atom interferometry. *Phys. Rev. A* **2002**, *65*, 033608.
- [12] Rosi, G.; Cacciapuoti, L.; Sorrentino, F.; Menchetti, M.; Prevedelli, M.; Tino, G. M. Measurement of the Gravity-Field Curvature by Atom Interferometry. *Phys. Rev. Lett.* **2015**, *114*, 013001.
- [13] Riehle, F.; Kisters, Th.; Witte, A.; Helmcke, J.; Borde, Ch. J. Optical Ramsey Spectroscopy in a Rotating Frame: Sagnac Effect in a Matter-Wave Interferometer. *Phys. Rev. Lett.* **1991**, *67*, 177-180.
- [14] Lenef, A.; Hammond, T. D.; Smith, E. T.; Chapman, M. S.; Rubenstein, R. A.; Pritchard, D. E. Rotation Sensing with an Atom Interferometer. *Phys. Rev. Lett.* **1997**, *78*, 760-763.
- [15] Gustavson, T. L.; Bouyer, P.; Kasevich, M. A. Precision Rotation Measurements with an Atom Interferometer Gyroscope. *Phys. Rev. Lett.* **1997**, *78*, 2046-2049.
- [16] Canuel, B.; Leduc, F.; Holleville, D.; Gauguier, A.; Fils, J.; Virdis, A.; Clairon, A.; Dimarcq, N.; Borde, Ch. J.; Landragin, A. Six-Axis Inertial Sensor Using Cold-Atom Interferometry. *Phys. Rev. Lett.* **2006**, *97*, 010402.
- [17] Dubetsky, B.; Kasevich, M. A. Atom interferometer as a selective sensor of rotation or gravity. *Phys. Rev. A* **2006**, *74*, 023615.
- [18] Barrett, B.; Geiger, R.; Dutta, I.; Meunier, M.; Canuel, B.; Gauguier, A.; Bouyer, P.; Landragin, A. The Sagnac effect: 20 years of development in matter-wave interferometry. *C. R. Physique* **2014**, *15*, 875-883.
- [19] Biedermann, G. W.; Wu, X.; Deslauriers, L.; Roy, S.; Mahadeswaraswamy, C.; Kasevich, M. A. Testing Gravity with Cold-Atom Interferometers. <http://arxiv.org/abs/1412.3210>.
- [20] Hamilton, P.; Jaffe, M.; Haslinger, P.; Simmons, Q.; Müller, H. Atom-Interferometry Constraints on Dark Energy. <http://arxiv.org/abs/1502.03888>
- [21] Borde, C. J.; Howard, J.-C.; Karasiewicz, A. Relativistic phase shifts for Dirac particles interacting with weak gravitational fields in matter-wave interferometers. *Lect. Notes Phys.* **2001**, *562*, 403-438.
- [22] Dimopoulos, S.; Graham, P. W.; Hogan, J. M.; Kasevich, M. A. General relativistic effects in atom interferometry. *Phys. Rev. D* **2008**, *78*, 042003.
- [23] Wicht, A.; Lämmerzahl, C.; Lorek, D.; Dittus, H. Rovibrational quantum interferometers and gravitational waves. *Phys. Rev. A* **2008**, *78*, 013610.
- [24] Dubetsky, B. Optimization and error model for atom interferometry technique to measure Newtonian gravitational constant. <http://arxiv.org/abs/1407.7287>.
- [25] Zorn, A.; Sonnad, V.; Libby, S. B.; Dubetsky, B.; and Shverdin, M, A Semi-Classical, Closed Form Expression for the Phase Response of a Vertical Symmetric Atomic Fountain, to be submitted to *Phys. Rev. A*, 2016.
- [26] Higher order terms in the expansion of the potential have also been considered in Ref. [36].
- [27] Bongs, K.; Launay, R.; Kasevich, M. A. High-order inertial phase shifts for time-domain atom interferometers. *Appl. Phys. B* **2006**, *84*, 599-602.
- [28] Kasevich, M. A.; Dubetsky, B. Kinematic Sensors Employing Atom Interferometer Phases. US Patent 2006/0249666 a1.
- [29] Dickerson, S. M.; Hogan, J. M.; Sugarbaker, A.; Johnson, D. M. S.; Kasevich, M. A. Multiaxis Inertial Sensing with Long-Time Point Source Atom Interferometry. *Phys. Rev. Lett.* **2013**, *111*, 083001.
- [30] Beausoleil, R. G.; Hansch, T. W. Ultrahigh-resolution two-photon optical Ramsey spectroscopy of an atomic fountain. *Phys. Rev. A* **1986**, *33*, 1661-1670.

- [31] Dubetsky, B. Gradiometer response on parallelepiped test mass. Unpublished. **2008**.
 [32] For a test mass density ρ , the field δg is of order $G\rho y_{\min}$. For inequality (2) to be satisfied, we must have

$$\rho \ll \frac{M_E}{y_{\min} R_E^2} \approx 1.5 \times 10^{12} \text{ kg/m}^3,$$

where M_E and R_E are Earth's mass and radius, respectively, and we put $y_{\min} \sim 0.1\text{m}$. Since the right-hand-side (rhs) of this inequality is 8 orders of magnitude larger than maximum atomic density existing on Earth, assumption (2) is reasonable.

- [33] Dubetsky, B.; Berman, P. R. Ground-state Ramsey fringes. *Phys. Rev. A* **1997**, *56*, R1091-R1094.
 [34] Kol'chenko, A. P.; Rautian, S. G.; Sokolovskii, R. I. Interaction of an atom with a strong electromagnetic field with the recoil effect taken into consideration. *Zh. Eksp. Teor. Fiz.* **1968**, *55*, 1864-1873 [*JETP* **1969**, *28*, 986-990].
 [35] Giese, E.; Zeller, W.; Kleinert, S.; Meister, M.; Tamma, V.; Roura, A.; Schleich, W. P. The interface of gravity and quantum mechanics illuminated by Wigner phase space Atom Interferometry. In *Proceedings of the International School of Physics "Enrico Fermi"*; Tino, G. N.; Kasevich, M. Eds.; IOS Press; **2014**; *188*, pp. 171 - 236.
 [36] Hogan, J. M. Testing Gravity with Atom Interferometry. www-conf.slac.stanford.edu/ssi/2011/Hogan-080311.pdf, Talk in *39th SLAC Summer Institute*; 2011.
 [37] Giltner, D. M.; McGowan, R. W.; Lee, S. A. Theoretical and experimental study of the Bragg scattering of atoms from a standing light wave. *Phys. Rev. A* **1995**, *52*, 3966-3972.
 [38] Berman, P. R.; Dubetsky, B.; Cohen, J. L. High-resolution amplitude and phase gratings in atom optics. *Phys. Rev. A* **1998**, *58*, 4801-4810.
 [39] Dubetsky, B.; Berman, P. R. $\lambda/4$, $\lambda/8$, and Higher Order Atom Gratings via Raman Transitions. *Laser Physics* **2002**, *12*, 1161-1170.
 [40] Turlapov, A.; Tonyushkin, A.; Sleator, T. Talbot-Lau effect for atomic de Broglie waves manipulated with light. *Phys. Rev. A* **2005**, *71*, 043612.
 [41] Lévêque, T.; Gauguier, A.; Michaud, F.; Pereira Dos Santos, F.; Landragin, A. Enhancing the Area of a Raman Atom Interferometer Using a Versatile Double-Diffraction Technique. *Phys. Rev. Lett.* **2009**, *103*, 080405.
 [42] Chiow, S.; Kovachy, T.; Chien, H.; Kasevich, M. A. 102hk Large Area Atom Interferometers. *Phys. Rev. Lett.* **2011**, *107*, 130403.
 [43] Dubetsky, B.; Berman, P. R. $\lambda/8$ -period optical potentials. *Phys. Rev. A* **2002**, *66*, 045402.
 [44] Zorn, A. Unpublished. **2011**.
 [45] Chemical elements listed by density. <http://www.lenntech.com/periodic-chart-elements/density.htm>