

# Pairing Nambu-Goldstone modes within nuclear density functional theory

Nobuo Hinohara<sup>1,2</sup> and Witold Nazarewicz<sup>3,4</sup>

<sup>1</sup>*Center for Computational Sciences, University of Tsukuba, Tsukuba, 305-8577, Japan*

<sup>2</sup>*FRIB Laboratory, Michigan State University, East Lansing, Michigan 48824, USA*

<sup>3</sup>*Department of Physics and Astronomy and FRIB Laboratory,  
Michigan State University, East Lansing, Michigan 48824, USA*

<sup>4</sup>*Institute of Theoretical Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland*

We show that the Nambu-Goldstone formalism of the broken gauge symmetry in the presence of the  $T = 1$  pairing condensate offers a quantitative description of the binding energy differences of open-shell superfluid nuclei. We conclude that the pairing rotational moments of inertia are excellent pairing indicators, which are free from ambiguities attributed to odd-mass systems. We offer a new, unified interpretation of the binding-energy differences traditionally viewed in the shell model picture as signatures of the valence nucleon properties. We present the first systematic analysis of the off-diagonal pairing rotational moments of inertia, and demonstrate the mixing of the neutron and proton pairing rotational modes in the ground states of even-even nuclei. Finally, we discuss the importance of mass measurements of neutron-rich nuclei for constraining the pairing energy density functional.

PACS numbers: 21.60.Jz, 21.10.Dr, 14.80.Va, 74.20.-z

*Introduction* — Spontaneous symmetry breaking explains the collective properties of atomic nuclei and provides straightforward physical interpretation of experimental observables associated with collective modes. In atomic nuclei, the Nambu-Goldstone (NG) mode [1, 2] connects two frames of reference: the intrinsic frame where the symmetry is broken and the NG mode is generated, and the laboratory frame where the symmetry is strictly conserved. The excitation of the NG mode can be observed in the laboratory system as a sequence of quantum states originating from a single symmetry-broken intrinsic state. One of the typical examples of the spontaneous symmetry breaking in atomic nuclei is the nuclear deformation due to the rotational symmetry breaking, as a consequence of the attractive particle-hole correlations [3–6]. Rotational bands can be viewed as NG modes in the body-fixed frame.

Nucleonic pairing is another common phenomenon in atomic nuclei associated with spontaneous symmetry breaking. Ground states of most nuclei can be well described as pair condensates, in which the particle number symmetry is broken. Superconducting nuclear states result in a NG mode called the pairing rotation, which is seen experimentally through ground-state sequences of even-even nuclei [7–11]. The topic of pairing rotations continues to generate much excitement, especially in the context of neutron-rich nuclei [12–16].

Nuclear density functional theory (DFT) is currently the only available microscopic many-body theory that is applicable to the whole nuclear chart. One of the reasons for its success is the flexibility of the formalism to naturally incorporate the spontaneous symmetry breaking mechanism. The form of the nuclear energy density functional (EDF) is constrained by symmetry considerations; popular Skyrme EDFs are built from density-

bilinear terms both in the particle-hole and pairing channels [17, 18]. Considerably less is known about the pairing EDF, primarily because of the lack of the experimental observables that can inform us about the detailed structure of the pairing EDF.

The coupling constants in the pairing EDF are usually fitted so that the theoretical pairing gaps in even-even nuclei reproduce the experimental odd-even mass differences, and such a strategy has been adopted in the recent optimization work [19–21]. Although the pairing gap is a fundamental order parameter for the superfluid phase, its relationship to the experimental odd-even mass difference is indirect. Moreover there exist multiple definitions of theoretical pairing gaps and there are various prescriptions for extracting the odd-even mass difference from experiment [22–24]. To avoid ambiguities, it would be best to calculate the odd-even mass difference directly from the theory. Unfortunately, this involves additional uncertainties pertaining to the definition of the ground state of an odd- $A$  nucleus [25]. Moreover, since ground-state configurations of odd- $A$  nuclei internally break the time-reversal symmetry, poorly-known time-odd terms of the EDF must be considered. Although some of the time-odd functionals are constrained through the local gauge invariance of the EDF [26], the optimization of the unconstrained time-odd coupling constants has barely started [27]. Consequently, the precision of the nuclear EDF for odd- $A$  systems is not as good as that for even-even systems. It is thus desirable to constrain the pairing EDF based on experimental data involving even-even systems only.

*Objectives* — In this Letter we assess the performance of nuclear DFT for pairing rotational bands in even-even nuclei, both semi-magic and doubly-open shell systems. We study pairing rotational moments of inertia and as-

sess their validity as indicators of nucleonic pairing. We check the sensitivity of pairing rotations in neutron-rich nuclei on the density dependence of the pairing functional.

*Definitions* — The pairing rotational picture is based on a single intrinsic “deformed” one-body field in a gauge space. The ground-state energy of a system with  $N/2$  fermionic pairs can be expanded up to the second order in the particle number with respect to a reference system with particle number  $N_0$  [8–10, 28],

$$E(N) = E(N_0) + \lambda_n(N_0)\Delta N + \frac{\Delta N^2}{2\mathcal{J}(N_0)}, \quad (1)$$

where  $\Delta N = N - N_0$ ,  $\lambda_n(N_0) = dE/dN|_{N=N_0}$  is the chemical potential, and the second order term is the pairing rotational energy with the moment of inertia  $\mathcal{J}(N_0)^{-1} = d^2E/dN^2|_{N=N_0}$ . In the case of a two-fermion system, Eq. (1) can be generalized by considering two coupled pairing rotational modes. In particular, when both neutrons and protons exhibit pair condensate, there exist two NG eigenmodes being linear combinations of the neutron and proton pairing rotations [29]. The corresponding rotational energy can be written as

$$E_{\text{rot}}^{\text{pair}} = \sum_{\tau, \tau' = n, p} \frac{\Delta N_\tau \Delta N_{\tau'}}{2\mathcal{J}_{\tau\tau'}}, \quad (2)$$

where  $N_n = N$ ,  $N_p = Z$ ,  $\Delta N_n = N - N_0$ ,  $\Delta N_p = Z - Z_0$ , and the tensor

$$\mathcal{J}_{\tau\tau'} = \left. \frac{\partial N_\tau}{\partial \lambda_{\tau'}} \right|_{\Delta N'_\tau = 0} = \left[ \frac{\partial^2 E}{\partial N_\tau \partial N_{\tau'}} \right]^{-1} \Big|_{\Delta N_\tau = \Delta N_{\tau'} = 0} \quad (3)$$

is the pairing rotational moment of inertia. The tensor  $\mathcal{J}_{\tau\tau'}$  is very sensitive to pairing correlations. Since it is related to the second derivative of the total energy with respect to particle number, the corresponding Thouless-Valatin (TV) inertia for the NG mode can be readily derived by means of the self-consistent quasiparticle random-phase approximation (QRPA) [30, 31].

In the region of particle numbers where static pairing dominates,  $\mathcal{J}_{\tau\tau'}$  can be extracted from experimental two-nucleon separation energies  $S_{2n}$  and  $S_{2p}$ . For instance, by taking  $\lambda_n(N, Z) = -\frac{1}{4}[S_{2n}(N+2, Z) + S_{2n}(N, Z)]$  the moments of inertia can be written as:

$$\mathcal{J}_{nn}^{-1}(N, Z) = \frac{1}{4} [S_{2n}(N, Z) - S_{2n}(N+2, Z)], \quad (4)$$

$$\mathcal{J}_{np}^{-1}(N, Z) = \frac{1}{4} [S_{2n}(N+2, Z+2) - S_{2n}(N+2, Z)]. \quad (5)$$

(The analogous expressions for  $\lambda_p$  and  $\mathcal{J}_{pp}$  are given in terms of  $S_{2p}$ .)

*Method* — To compute the TV moments of inertia for pairing rotations we employ the linear response formalism of nuclear DFT in the finite amplitude method

(FAM) [32] variant. The FAM allows one to handle all the two-quasiparticle states on the QRPA level with a smaller computational cost than that of the traditional matrix formulation of the QRPA. The TV moment of inertia is given by a response function of the particle number operator at zero frequency. In this study, we follow the FAM formulation of Ref. [29] for NG modes.

The computations were performed with the FAM code [33, 34] using the DFT solver HFBTHO [35] in a single-particle basis consisting of 20 harmonic oscillator shells. We employed the recently developed EDF UNEDF1-HFB [36] that was optimized at the full Hartree-Fock-Bogoliubov (HFB) level. For the pairing energy density we use the density-dependent ansatz [37]

$$\tilde{\chi}_\tau(\mathbf{r}) = \frac{1}{2}V_0^\tau \left[ 1 - \eta \frac{\rho_0(\mathbf{r})}{\rho_c} \right] |\tilde{\rho}_\tau(\mathbf{r})|^2, \quad (6)$$

where  $\tilde{\rho}$  is the pairing density,  $\rho_0$  is the isoscalar density,  $\rho_c = 0.16 \text{ fm}^{-3}$ ,  $V_0^\tau$  is the strength, and  $\eta$  is the parameter that controls the density dependence of the pairing interaction.

In UNEDF1-HFB, the mixed-type pairing ( $\eta = 0.5$ ) is employed. To analyze the sensitivity of results on the density dependence of pairing functional, we also studied the volume-type ( $\eta = 0$ ) and surface-type ( $\eta = 1$ ) pairing with the strengths adjusted to reproduce the average neutron pairing gap in  $^{120}\text{Sn}$  and average proton pairing gap in  $^{92}\text{Mo}$  assuming the default pairing energy window of 60 MeV. These nuclei were chosen because the average pairing gaps computed with UNEDF1-HFB are close to the experimental values. The resulting pairing strengths are  $V_0^n = -146.07 \text{ MeV fm}^3$  and  $V_0^p = -161.72 \text{ MeV fm}^3$  for the volume pairing, and  $V_0^n = -474.32 \text{ MeV fm}^3$  and  $V_0^p = -551.37 \text{ MeV fm}^3$  for the surface pairing.




FIG. 1. (Color online). Neutron pairing rotational energy measured from a reference state in  $^{116}\text{Sn}$ . The parabolic expression (1) with  $\lambda_n$  and  $\mathcal{J}_{nn}$  evaluated for the reference nucleus is shown as a solid line. The HFB (squares) and experimental (circles) values have been extracted from binding energies according to  $E_{\text{rot}}^{\text{pair}} = E(N) - E(66) - \lambda_n(66)(N - 66)$ .

*Results* — We start with the classic case of neutron

pairing rotations in a semi-magic chain of Sn isotopes [10]. The theoretical values of the chemical potential and the TV inertia have been computed for the reference nucleus  $^{116}\text{Sn}$  ( $N_0 = 66$ ). As seen in Fig. 1, the harmonic approximation (1) works very well in this case; indeed, the TV pairing inertia agrees with experiment even when  $N$  is far from  $N_0$ . This shows that a single intrinsic pairing field of  $^{116}\text{Sn}$  explains the binding energy behavior in terms of the dynamics of the NG mode.

In general, the higher order corrections in  $\Delta N$  are not negligible; in analogy with the angular momentum alignment within a rotational band, a change of the intrinsic structure with neutron number is expected along a pairing rotational band. This is seen in Fig. 1 through the deviation of the HFB values (or experiment) from the parabolic behavior. To account for the changes of the intrinsic pairing field, we carry out systematic FAM+HFB calculations for chains of semi-magic nuclei. Figure 2 displays associated chemical potentials and pairing rotational moments of inertia.

The pairing rotational moments of inertia for Sn and Pb isotopes behave fairly smoothly, and the pairing rotational picture holds in the medium-mass Ca isotopes. In general, we see a remarkably good agreement between TV moments of inertia with experiment. The exceptions are weakly-paired systems around the magic numbers for which a transition to the pairing vibrational picture takes place. In such cases, e.g., for  $^{130}\text{Sn}$  and  $^{42,46,50}\text{Ca}$ , the experimental indicator (4) involves nuclei for which our HFB calculations predict vanishing pairing. The finite-difference approximation of the second-order derivative is questionable there.

For the doubly closed shell nuclei, the theoretical pairing rotational inertia is zero as the NG mode is absent due to the vanishing static pairing. Moreover, the expression (4) for the experimental inertia  $\mathcal{J}_{\tau\tau}$  is proportional to the inverse of the so-called two-nucleon shell gap indicator  $\delta_{2\tau}$  [39]. This latter quantity has been attributed to the size of the magic gap. As it was already noted in Ref. [40], the validity of  $\delta_{2\tau}$  as a signature of a shell closure is lost in regions where the structure of nuclear ground states is rapidly changing. Based on our results for semi-magic nuclei shown in Fig. 2, we can make an even stronger statement: outside shell closures, the two-nucleon shell gap indicator  $\delta_{2\tau}$  has nothing or little to do with the distribution of single-particle energies; it is primarily governed by pairing correlations and serves as a good indicator of the gauge symmetry breaking.

We now study the proton pairing by investigating pairing rotation in the  $N = 50$  isotones. As shown in Fig. 2(d), the proton pairing moments of inertia are smaller than the neutron ones in the similar mass region, and the agreement with experiment is excellent. In the figure, we also plot the Belyaev moment of inertia [41], which does not include the effect of the residual correlations at the QRPA level. As discussed in Ref. [29],

the difference between TV and Belyaev proton inertia can be attributed to the Coulomb-induced QRPA correlations. Here we recall that the proton pairing strength required to provide good agreement with experimental odd-even mass differences is significantly larger than the neutron strength,  $V_0^p/V_0^n \approx 1.1$ , and this is consistent with the results of the global survey [24]. The large effect of Coulomb correlations on  $\mathcal{J}_{pp}$ , manifesting itself through the difference between Belyaev and TV proton pairing rotational inertia, confirms the conclusion of Ref. [42] that the Coulomb substantially suppresses proton pairing.

The density and momentum dependence of the pairing functional are not well known because standard observables probing pairing channel, such as odd-even mass staggering or moments of inertia of deformed nuclei, show weak sensitivity to details. In this context, pairing rotational inertia of single-shell-closed nuclei can serve as a good indicator of the pairing interaction. The results of calculations for semi-magic nuclei in Fig. 2, based on pairing fitted to experimental odd-even mass differences, are fairly similar for volume-, mixed-, and surface-pairing variants, except for very neutron-rich nuclei where the surface pairing gives appreciably lower values of  $\mathcal{J}_{nn}$ . Of particular interest is the behavior of pairing rotational inertia in the very neutron-rich Ca isotopes beyond  $^{56}\text{Ca}$ , where the pairing functional of volume type yields 1.5–2 times larger value of  $\mathcal{J}_{nn}$  than the mixed pairing interaction. Mass measurements of even-even Ca isotopes beyond  $N = 36$  will be useful to better constrain the density dependence of the pairing EDF.

Finally we discuss the doubly-open-shell nuclei. When both neutrons and protons show pair condensate, there exist two NG modes whose eigenmodes are linear combinations of the neutron and proton pairing rotations [29]. In Fig. 3, we show the full pairing rotational moment of inertia tensor for the Er isotopes and  $N = 100$  isotones. Both examples are representative of well deformed, open-shell nuclei with static neutron and proton pairing. Our calculations give an excellent agreement with experiment for the full pairing rotational inertia tensor. In general, the sensitivity to the density dependence of pairing interaction is fairly weak except for very neutron-rich (proton-deficient) nuclei.

The off-diagonal moment of inertia  $\mathcal{J}_{np}$  shows quantitative agreement with the experimental data. We emphasize that the example shown in Fig. 3 represents the first systematic calculation of the off-diagonal inertia for the two-dimensional pairing rotation, which was seen as a tilted energy kernel in the gauge space in Ref. [43]. The agreement with experiment confirms that the two pairing-rotational NG modes are indeed mixed through the residual interaction in QRPA. Another interesting aspect of  $\mathcal{J}_{np}$  is that the inverse of this quantity is formally equivalent – up to a trivial shift ( $Z \rightarrow Z+2, N \rightarrow N+2$ ) – to the mass indicator  $-\delta V_{pn}$  [44], often referred to as, and




FIG. 2. (Color online). Chemical potential (top panels) and pairing rotational moment of inertia (bottom panels) for Sn (a), Pb (b), and Ca (c) isotopes, and  $N = 50$  isotones (d). Spherical UNEDF1-HFB solutions with mixed (squares), volume (circles), and surface (triangles) pairing are compared to experimental data from Ref. [38]. For  $N = 50$  isotones, Belyaev moments of inertia are also shown by lines without symbols.




FIG. 3. (Color online). Pairing rotational moments of inertia  $J_{nn}$  (top),  $J_{np}$  (middle), and  $J_{pp}$  (bottom) for open-shell Er isotopes (left) and  $N = 100$  isotones (right) exhibiting static neutron and proton pairing.

interpreted in terms of, the empirical proton-neutron interaction energy. Indeed, in the extreme shell model picture,  $\delta V_{pn}$  represents the net interaction of the last two valence neutrons with the last two valence protons [45–48]. While the large-scale superfluid DFT calculations of  $\delta V_{pn}$  generally match the experimental data on the double binding-energy difference (5) [49], the direct interpretation of this quantity in terms of the valence proton-neutron interaction is under debate [50]. As pointed out in Ref. [49], while the value of  $\delta V_{pn}$  averaged over many states (shells) approximates probes the bulk symmetry energy term of the EDF, the *local* behavior of  $\delta V_{pn}$  carries important information about shell effects and many-body correlations. The relation (5) between  $J_{np}$  and  $\delta V_{pn}$

sheds a new light on the interpretation of this quantity in doubly open-shell nuclei in terms of the simultaneous spontaneous breaking of the neutron *and* proton gauge symmetries of the  $T = 1$  pairing. In this respect, we would question the findings of Ref. [47] that the pairing energy plays a relatively minor role in understanding of  $\delta V_{pn}$ .

*Conclusions* — We show that the  $T = 1$  pairing rotational moments of inertia of semi-magic and doubly-open-shell nuclei can be described qualitatively within the NG formalism of the broken gauge symmetry. Since the experimental mass difference relation representing the pairing inertia tensor is solely based on binding energies of even-even nuclei, it is an excellent indicator of nuclear pairing properties. In many respects,  $\mathcal{J}_{\tau\tau'}$  is superior to other quantities commonly used to inform us about the magnitude of pairing correlations, such as odd-even mass differences, which involve properties of odd-mass systems that depend on poorly known time-odd fields impacting individual orbits blocked by an odd nucleon. Furthermore, we demonstrate that the pairing rotational inertia tensor can be directly expressed in terms of the binding energies differences  $\delta_{2n}$ ,  $\delta_{2p}$ , and  $\delta V_{pn}$  - all traditionally regarded as signatures of the valence nucleon properties in the shell model picture. We now propose a unified interpretation of these quantities in terms of the gauge symmetry breaking associated with the collective  $T = 1$  pairing phases. Of course, for nuclei close to shell or subshell closures, with weak pairing correlations, the traditional single-particle interpretation is expected to be more appropriate.

We present the first systematic analysis of the off-diagonal pairing rotational moments of inertia  $\mathcal{J}_{np}$ , and demonstrate the mixing of the neutron and proton pairing rotational modes in the ground states of open-shell even-even nuclei. Our analysis of isotopic and isotonic chains indicates that the pairing rotational moments of inertia of neutron-rich nuclei can be used to constrain the

pairing functional of nuclear DFT, including its density dependence. In this context, mass measurements of very neutron-rich isotopes are extremely desirable. Theoretically, clarifying the role of the missing neutron-proton contribution of the  $T = 1$  pairing functional to  $\mathcal{J}_{np}$  within the isospin invariant EDF [17, 51, 52] will be an exciting subject for future investigations.

This work is supported by the U.S. Department of Energy, Office of Sciences, Office of Nuclear Physics under Awards No. de-sc0013365 (Michigan State University) and No. de-sc0008511 (NUCLEI SciDAC-3). Numerical calculation was performed at the COMA (PACS-IX) System at the Center for Computational Sciences, University of Tsukuba.

- 
- [1] Y. Nambu, Phys. Rev. **117**, 648 (1960).  
 [2] J. Goldstone, Il Nuovo Cimento **19**, 154 (1961).  
 [3] A. Bohr and B. R. Mottelson, *Nuclear Structure, Vol. II* (W. A. Benjamin, Reading, MA, 1975).  
 [4] P.-G. Reinhard and E. W. Otten, Nucl. Phys. A **420**, 173 (1984).  
 [5] W. Nazarewicz, Nucl. Phys. A **574**, 27 (1994).  
 [6] S. Frauendorf, Rev. Mod. Phys. **73**, 463 (2001).  
 [7] D. R. Bes and R. A. Broglia, Nucl. Phys. **80**, 289 (1966).  
 [8] B. F. Bayman, D. R. Bes, and R. A. Broglia, Phys. Rev. Lett. **23**, 1299 (1969).  
 [9] D. R. Bes and R. A. Broglia, Varenna Lectures **69**, 55 (1977).  
 [10] D. M. Brink and R. A. Broglia, *Nuclear Superfluidity, Pairing in Finite Systems* (Cambridge University Press, Cambridge, UK, 2005).  
 [11] R. A. Broglia, J. Terasaki, and N. Giovanardi, Phys. Rep. **335**, 1 (2000).  
 [12] W. von Oertzen and A. Vitturi, Rep. Prog. Phys. **64**, 1247 (2001).  
 [13] H. Shimoyama and M. Matsuo, Phys. Rev. C **84**, 044317 (2011).  
 [14] R. M. Clark, A. O. Macchiavelli, L. Fortunato, and R. Krücken, Phys. Rev. Lett. **96**, 032501 (2006).  
 [15] G. Potel, F. Barranco, F. Marini, A. Idini, E. Vigezzi, and R. A. Broglia, Phys. Rev. Lett. **107**, 092501 (2011), **108**, 069904 (2012).  
 [16] G. Potel, A. Idini, F. Barranco, E. Vigezzi, and R. A. Broglia, Phys. Rev. C **87**, 054321 (2013).  
 [17] E. Perlińska, S. G. Rohoziński, J. Dobaczewski, and W. Nazarewicz, Phys. Rev. C **69**, 014316 (2004).  
 [18] S. G. Rohoziński, J. Dobaczewski, and W. Nazarewicz, Phys. Rev. C **81**, 014313 (2010).  
 [19] M. Kortelainen, T. Lesinski, J. Moré, W. Nazarewicz, J. Sarich, N. Schunck, M. V. Stoitsov, and S. Wild, Phys. Rev. C **82**, 024313 (2010).  
 [20] M. Kortelainen, J. McDonnell, W. Nazarewicz, P.-G. Reinhard, J. Sarich, N. Schunck, M. V. Stoitsov, and S. M. Wild, Phys. Rev. C **85**, 024304 (2012).  
 [21] M. Kortelainen, J. McDonnell, W. Nazarewicz, E. Olsen, P.-G. Reinhard, J. Sarich, N. Schunck, S. M. Wild, D. Davesne, J. Erler, and A. Pastore, Phys. Rev. C **89**, 054314 (2014).  
 [22] W. Satuła, J. Dobaczewski, and W. Nazarewicz, Phys. Rev. Lett. **81**, 3599 (1998).  
 [23] T. Duguet, P. Bonche, P.-H. Heenen, and J. Meyer, Phys. Rev. C **65**, 014311 (2001).  
 [24] G. F. Bertsch, C. A. Bertulani, W. Nazarewicz, N. Schunck, and M. V. Stoitsov, Phys. Rev. C **79**, 034306 (2009).  
 [25] N. Schunck, J. Dobaczewski, J. McDonnell, J. Moré, W. Nazarewicz, J. Sarich, and M. V. Stoitsov, Phys. Rev. C **81**, 024316 (2010).  
 [26] J. Dobaczewski and J. Dudek, Phys. Rev. C **52**, 1827 (1995).  
 [27] M. T. Mustonen and J. Engel, Phys. Rev. C, in press, arXiv:1510.02136.  
 [28] R. Beck, M. Kleber, and H. Schmidt, Z. Phys. **250**, 155 (1972).  
 [29] N. Hinohara, Phys. Rev. C **92**, 034321 (2015).  
 [30] D. J. Thouless and J. G. Valatin, Nucl. Phys. **31**, 211 (1962).  
 [31] P. Ring and P. Schuck, *The Nuclear Many-Body Problem* (Springer-Verlag, Berlin, 2000).  
 [32] T. Nakatsukasa, T. Inakura, and K. Yabana, Phys. Rev. C **76**, 024318 (2007).  
 [33] M. Stoitsov, M. Kortelainen, T. Nakatsukasa, C. Losa, and W. Nazarewicz, Phys. Rev. C **84**, 041305 (2011).  
 [34] N. Hinohara, M. Kortelainen, W. Nazarewicz, and E. Olsen, Phys. Rev. C **91**, 044323 (2015).  
 [35] M. V. Stoitsov, N. Schunck, M. Kortelainen, N. Michel, H. Nam, E. Olsen, J. Sarich, and S. Wild, Comput. Phys. Commun. **184**, 1592 (2013).  
 [36] N. Schunck, J. D. McDonnell, J. Sarich, S. M. Wild, and D. Higdon, J. Phys. G **42**, 034024 (2015).  
 [37] J. Dobaczewski, W. Nazarewicz, and M. V. Stoitsov, Eur. Phys. J. A **15**, 21 (2002).  
 [38] G. Audi, M. Wang, A. H. Wapstra, F. G. Kondev, M. MacCormick, X. Xu, and B. Pfeiffer, Chin. Phys. C **36**, 1287 (2012).  
 [39] D. Lunney, J. M. Pearson, and C. Thibault, Rev. Mod. Phys. **75**, 1021 (2003).  
 [40] M. Bender, T. Cornelius, G. A. Lalazissis, J. A. Maruhn, W. Nazarewicz, and P.-G. Reinhard, Eur. Phys. J. A **14**, 23 (2002).  
 [41] S. T. Beliaev, Nucl. Phys. **24**, 322 (1961).  
 [42] M. Anguiano, J. L. Egido, and L. M. Robledo, Nucl. Phys. A **683**, 227 (2001).  
 [43] X. B. Wang, J. Dobaczewski, M. Kortelainen, L. F. Yu, and M. V. Stoitsov, Phys. Rev. C **90**, 014312 (2014).  
 [44] J.-Y. Zhang, R. F. Casten, and D. S. Brenner, Phys. Lett. B **227**, 1 (1989).  
 [45] R. B. Cakirli and R. F. Casten, Phys. Rev. Lett. **96**, 132501 (2006).  
 [46] Y. Oktem, R. B. Cakirli, R. F. Casten, R. J. Casperson, and D. S. Brenner, Phys. Rev. C **74**, 027304 (2006).  
 [47] C. Qi, Phys. Lett. B **717**, 436 (2012).  
 [48] D. Bonatsos, S. Karampagia, R. B. Cakirli, R. F. Casten, K. Blaum, and L. A. Susam, Phys. Rev. C **88**, 054309 (2013).  
 [49] M. Stoitsov, R. B. Cakirli, R. F. Casten, W. Nazarewicz, and W. Satuła, Phys. Rev. Lett. **98**, 132502 (2007).  
 [50] M. Bender and P.-H. Heenen, Phys. Rev. C **83**, 064319 (2011).  
 [51] K. Sato, J. Dobaczewski, T. Nakatsukasa, and W. Satuła, Phys. Rev. C **88**, 061301 (2013).  
 [52] J. A. Sheikh, N. Hinohara, J. Dobaczewski, T. Nakat-

sukasa, W. Nazarewicz, and K. Sato, Phys. Rev. C **89**, 054317 (2014).