

Implications of the diphoton excess on Left-Right models and gauge unification

Frank F. Deppisch,^{1,*} Chandan Hati,^{2,3,†} Sudhanwa Patra,^{4,‡} Prativa Pritimita,^{4,§} and Utpal Sarkar^{2,¶}

¹*Department of Physics and Astronomy, University College London, London WC1E 6BT, United Kingdom*

²*Physical Research Laboratory, Navrangpura, Ahmedabad 380 009, India*

³*Indian Institute of Technology Gandhinagar, Chandkheda, Ahmedabad 382 424, India*

⁴*Center of Excellence in Theoretical and Mathematical Sciences, Siksha 'O' Anusandhan University, Bhubaneswar-751030, India*

The recent diphoton excess signal at an invariant mass of 750 GeV can be interpreted in the framework of left-right symmetric models with additional scalar singlets and vector-like fermions. We propose a minimal scenario for such a purpose. Extending the LRSM framework to include these new vector-like fermionic fields, on the other hand, results in interesting phenomenological implications for the LRSM fermion masses and mixing. Furthermore, existence of such vector-like fermions can also have interesting implications for baryogenesis and the dark matter sector. The introduction of a real bi-triplet scalar which contains a potential DM candidate will allow the gauge couplings to unify at $\approx 10^{17.5}$ GeV.

I. INTRODUCTION

The CMS and ATLAS collaborations have recently reported a roughly 3σ excess in the diphoton channel at an invariant mass of about 750 GeV in the first 3 fb^{-1} of collected data from Run 2 of the LHC at 13 TeV [1, 2]. The Landau-Yang theorem forbids the possibility of a massive spin one resonance decaying to $\gamma\gamma$. The leading interpretations of the excess within the context of new physics scenarios therefore consist of postulating a fundamental spin zero or spin two particle with mass of about 750 GeV. However no enhancements have been seen in the dijet, $t\bar{t}$, diboson or dilepton channels posing a clear challenge to the possible interpretations of this excess. The absence of a peaked $\gamma\gamma$ angular distribution in the observed events towards the beam direction disfavors [3] the spin two hypothesis and the spin zero resonance interpretation seems more favourable from a theoretical point of view.

A large number of interpretations of the diphoton signal in terms of physics beyond the Standard Model have been proposed in the literature [4–119]. One of the possibilities that has been largely explored in the literature is a scalar or pseudo-scalar resonance produced through gluon-gluon fusion and decaying to $\gamma\gamma$ via loop diagrams with circulating fermions or bosons. A new resonance coupling with the Standard Model (SM) t quark or W^\pm can give rise to such loop diagrams, however, they will be highly suppressed at the large $\gamma\gamma$ invariant masses and the dominant decay channel would have to be $t\bar{t}$ or W^+W^- . Hence the observation of the $\gamma\gamma$ resonance at 750 GeV (much greater than the electroweak symmetry breaking scale) hints towards the existence of vector-like fermions around that mass scale. Given that both the ATLAS and CMS collaborations have suggested signal

events consistent with each other at a tempting 3σ statistical significance level, hinting towards a new physics scenario, it is important to explore the possible model framework that can naturally accommodate such vector-like fermions.

From a theoretical stand point, a framework that can explain the diphoton excess while being consistent with other searches for new physics are particularly intriguing. To this end, one must mention the results reported by the CMS Collaboration in the first run of LHC for the right-handed gauge boson W_R search at $\sqrt{s} = 8$ TeV and 19.7 fb^{-1} of integrated luminosity [120]. A 2.8σ local excess was reported in the $eejj$ channel in the energy range $1.8 \text{ TeV} < m_{eejj} < 2.2 \text{ TeV}$, hinting at a right handed gauge counterpart of the SM $SU(2)_L$ broken around the TeV scale. The Left-Right Symmetric Model (LRSM) framework with $g_R \neq g_L$ can explain such signal with the possibility of being embedded into a ultraviolet complete higher gauge group [121–123]. It is thus an interesting exercise to explore the possibility of naturally accommodating the $\gamma\gamma$ excess also in such a framework.

In this paper, we explore the possibility of extending the standard LRSM framework with vector-like fermions and singlet scalars which can explain the diphoton signal. Adding such new vector-like fermionic fields, on the other hand, results in interesting phenomenological implications for the LRSM fermion masses and mixing. Moreover, existence of such vector-like fermions can have interesting implications for baryogenesis and the potential dark matter sector. In gauged flavor groups with left-right symmetry [124] or quark-lepton symmetric models [125], vector-like fermions are naturally accommodated while in LRSMs originating from D-brane or heterotic string compactifications also often include vector-like fermions [126, 127]. We propose a minimal LRSM that hosts such vector-like fermions and which can explain the diphoton signal. We also explore the possible fermion masses and mixing phenomenology and the implications of these vector-like particles in baryogenesis and the dark matter sector.

The plan of rest of this paper is as follows. In section II, we discuss the LRSM accommodating new vector-like

*Electronic address: f.deppisch@ucl.ac.uk

†Electronic address: chandan@prl.res.in

‡Electronic address: sudha.astro@gmail.com

§Electronic address: pratibha.pritimita@gmail.com

¶Electronic address: utpal@prl.res.in

fermions and the implications on masses and mixing of the fermions. In section III, we discuss the general aspects of the diphoton signal in the context of a scalar resonance decaying to $\gamma\gamma$ through circulating vector-like fermions in the loop. In section IV, we explore the possibility of obtaining a gauge coupling unification including the new vector like fields. In section V, we discuss the implications of the vector-like fermions for baryogenesis and the dark matter sector. In section VI, we summarize and make concluding remarks.

II. LEFT-RIGHT SYMMETRIC MODEL FRAMEWORK

The left-right symmetric extension of the SM has the basic gauge group given by

$$\mathcal{G}_{L,R} \equiv SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}, \quad (1)$$

where $B-L$ is the difference between baryon and lepton number. The electric charge is related to the third component of isospin in the $SU(2)_{L,R}$ gauge groups and the $B-L$ charge as

$$Q = T_{3L} + T_{3R} + 1/2(B-L). \quad (2)$$

The quarks and leptons transform under the LRSM gauge group as

$$q_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix} \equiv [2, 1, \frac{1}{3}, 3], \quad q_R = \begin{pmatrix} u_R \\ d_R \end{pmatrix} \equiv [1, 2, \frac{1}{3}, 3],$$

$$\ell_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \equiv [2, 1, -1, 1], \quad \ell_R = \begin{pmatrix} \nu_R \\ e_R \end{pmatrix} \equiv [1, 2, -1, 1],$$

where the gauge group representations are written in the form $[SU(2)_L, SU(2)_R, B-L, SU(3)_C]$.

Originally the left-right symmetric extension of the SM [128–133], was introduced to give a natural explanation for parity violation seen in radioactive beta decay and to consistently address the light neutrino masses through introduction of right-handed neutrinos. The latter form doublets with the right handed charged fermions under the $SU(2)_R$ gauge group. If $SU(2)_R$ breaks at around the TeV scale, LRSMs offer a rich interplay between high energy collider signals and low energy processes such as neutrinoless double beta decay and lepton flavor violation [134]. The principal prediction of this scenario is a TeV scale right-handed gauge boson W_R . The CMS and ATLAS collaborations had reported several excesses at around 2 TeV in Run 1 of the LHC, pointing towards such a possibility. From the first results of Run 2, no di-jet and diboson excesses have been reported (more data is required to exclude the diboson excesses reported in run 1), the relatively “cleaner” $eejj$ channel signal is still present hinting at a 2 TeV W_R . In light of the diphoton excess it is important to revisit the LRSM framework to explore the possibility of accommodating such signal and the possible implications.

As already mentioned in the introduction and discussed in section III, the 750 GeV diphoton excess can be

explained through the resonant production and decay of a scalar or pseudoscalar particle. To this end, we propose a simple left-right symmetric model with a scalar singlet S and vector-like fermions added to the minimal particle content of left-right symmetric models¹.

We extend the standard LRSM framework to include isosinglet vector-like copies of LRSM fermions. This kind of a vector-like fermion spectrum is very naturally embedded in gauged flavour groups with left-right symmetry [124] or quark-lepton symmetric models [125]. The field content of this model and the relevant transformations under the LRSM gauge group are shown in Tab. I.

Field	$SU(2)_L$	$SU(2)_R$	$B-L$	$SU(3)_C$
q_L	2	1	1/3	3
q_R	1	2	1/3	3
ℓ_L	2	1	-1	1
ℓ_R	1	2	-1	1
$U_{L,R}$	1	1	4/3	3
$D_{L,R}$	1	1	-2/3	3
$E_{L,R}$	1	1	-2	1
$N_{L,R}$	1	1	0	1
H_L	2	1	-1	0
H_R	1	2	-1	0
S	1	1	0	0

TABLE I: LRSM representations of extended field content.

The relevant Yukawa part of the Lagrangian is given by

$$\mathcal{L} = - \sum_X (\lambda_{SXX} S \bar{X} X + M_X \bar{X} X) - (\lambda_U^L \tilde{H}_L \bar{q}_L U_R + \lambda_U^R \tilde{H}_R \bar{q}_R U_L + \lambda_D^L H_L \bar{q}_L D_R + \lambda_D^R H_R \bar{q}_R D_L + \lambda_E^L H_L \bar{\ell}_L E_R + \lambda_E^R H_R \bar{\ell}_R E_L + \lambda_N^L \tilde{H}_L \bar{\ell}_L N_R + \lambda_N^R \tilde{H}_R \bar{\ell}_R N_L + \text{h.c.}), \quad (3)$$

where the summation is over $X = U, D, E, N$ and we suppress flavour and colour indices on the fields and couplings. $\tilde{H}_{L,R}$ denotes $\tau_2 H_{L,R}^*$, where τ_2 is the usual second Pauli matrix.

The vacuum expectation values (VEVs) of the Higgs doublets $H_R(1, 2, -1)$ and $H_L(2, 1, -1)$ break the LRSM gauge group to the SM gauge group and the SM gauge group to $U(1)_{\text{EM}}$ respectively, with an ambiguity of parity breaking, which can either be broken at the TeV scale or at a much higher scale M_P . In the latter case, the Yukawa couplings can be different for right-type and left-type Yukawa terms because of the renormalization group running below M_P , $\lambda_X^R \neq \lambda_X^L$. Hence, we distinguish the left and right handed couplings explicitly

¹ We assume the resonance to be a new singlet scalar and it can easily be generalized to a pseudoscalar case.

with the subscripts L and R . We use the VEV normalizations $\langle H_L \rangle = (0, v_L)^T$ and $\langle H_R \rangle = (0, v_R)^T$ with $v_L = 175$ GeV and v_R constrained by searches for the heavy right-handed W_R boson at colliders and at low energies, $v_R \gtrsim 1 - 3$ TeV (depending on the right-handed gauge coupling). Due to the absence of a bidoublet Higgs scalar, normal Dirac mass terms for the SM fermions are absent and the charged fermion mass matrices assume a seesaw structure. However, if one does not want to depend on a “universal” seesaw structure, a Higgs bidoublet Φ can be introduced along with $H_{L,R}$.

After symmetry breaking, the mass matrices for the fermions are given by

$$M_{uU} = \begin{pmatrix} 0 & \lambda_U^L v_L \\ \lambda_U^R v_R & M_U \end{pmatrix}, M_{dD} = \begin{pmatrix} 0 & \lambda_D^L v_L \\ \lambda_D^R v_R & M_D \end{pmatrix},$$

$$M_{eE} = \begin{pmatrix} 0 & \lambda_E^L v_L \\ \lambda_E^R v_R & M_E \end{pmatrix}, M_{\nu N} = \begin{pmatrix} 0 & \lambda_N^L v_L \\ \lambda_N^R v_R & M_N \end{pmatrix}, \quad (4)$$

The mass eigenstates can be found by rotating the mass matrices via left and right orthogonal transformations $O^{L,R}$ (we assume all parameters to be real). For example, the up quark diagonalization yields $O_U^{LT} \cdot M_{uU} \cdot O_U^R = \text{diag}(\hat{m}_u, \hat{M}_U)$. Up to leading order in $\lambda_U^L v_L$, the resulting up-quark masses are

$$\hat{M}_U \approx \sqrt{M_U^2 + (\lambda_U^R v_R)^2}, \quad \hat{m}_u \approx \frac{(\lambda_U^L v_L)(\lambda_U^R v_R)}{\hat{M}_U}, \quad (5)$$

and the mixing angles $\theta_U^{L,R}$ parametrizing $O_U^{L,R}$,

$$\tan(2\theta_U^L) \approx \frac{2(\lambda_U^L v_L)M_U}{M_U^2 + (\lambda_U^R v_R)^2}, \quad \tan(2\theta_U^R) \approx \frac{2(\lambda_U^R v_R)M_U}{M_U^2 - (\lambda_U^R v_R)^2}. \quad (6)$$

The other fermion masses and mixings are given analogously. For an order of magnitude estimate one may approximate the phenomenologically interesting regime with the limit $\lambda_U^R v_R \rightarrow M_U$ in which case the mixing angles approach $\theta_U^L \rightarrow \hat{m}_u/\hat{M}_U$ and $\theta_U^R \rightarrow \pi/4$. This means that θ_U^L is negligible for all fermions but the top quark and its vector partner [90].

We here neglect the flavour structure of the Yukawa couplings $\lambda_X^{L,R}$ and λ_{SXX} which will determine the observed quark and leptonic mixing. The hierarchy of SM fermion masses can be generated by either a hierarchy in the Yukawa couplings or in the masses of the of the vector like fermions.

As described above, the light neutrino masses are of Dirac-type as well, analogously given by

$$\hat{m}_\nu = \frac{\lambda_N^L \lambda_N^R v_L v_R}{M_N}, \quad (7)$$

It is natural to assume that $M_N \gg v_R$, as the vector like N is a singlet under the model gauge group. In this case, the scenario predicts naturally light Dirac neutrinos [124].

III. DIPHOTON SIGNAL FROM A SCALAR RESONANCE

One may attempt to interpret the diphoton excess at as the resonant production of the singlet scalar S with mass $M_S = 750$ GeV. Considering the possible production mechanisms for the resonance at 750 GeV it is interesting to note that the CMS and ATLAS did not report a signal in the $\sim 20\text{fb}^{-1}$ data at 8 TeV in Run 1. One possible interpretation of this can be that the resonance at 750 GeV is produced through a mechanism with a steeper energy dependence. Excluding the possibility of an associated production of this resonance, the most favourable mechanism is gluon-gluon fusion which we here also consider as the dominant production mechanism. Subsequently, the scalar with mass 750 GeV decays to two photons via a loop as well. The cross section can be expressed as

$$\sigma(pp \rightarrow \gamma\gamma) = \frac{C_{gg}}{M_{Ss}} \Gamma_{gg} \text{Br}_{\gamma\gamma}, \quad (8)$$

with the proton centre of mass energy \sqrt{s} and the parton distribution integral $C_{gg} = 174$ at $\sqrt{s} = 8$ TeV and $C_{gg} = 2137$ at $\sqrt{s} = 13$ TeV [14]. One can obtain a best fit guess of the cross section by reconstructing the likelihood, assumed to be Gaussian, from the 95% C.L. expected and observed limits in an experimental search. For the diphoton excess, we use a best fit cross section value of 7 fb found by combining the 95% CL ranges from ATLAS and CMS at 13 TeV and 8 TeV for a resonance mass of 750 GeV [14].

Apart from the necessary decay modes of the scalar S i.e. $S \rightarrow gg$ and $S \rightarrow \gamma\gamma$, S may also decay to other particles; due to the necessary SM invariance and the fact that $M_S > m_Z$, $S \rightarrow \gamma\gamma$ necessitates the decays $S \rightarrow \gamma Z$ and ZZ which are suppressed by $2 \tan^2 \theta_W \approx 0.6$ and $\tan^4 \theta_W \approx 0.1$ relative to $\Gamma(S \rightarrow \gamma\gamma)$ [14]. Furthermore, S in our model may also decay to SM fermions due to mixing with the heavy vector-like fermions. As described above, the mixing is only sizeable for the top and its vector partner. The total width is thereby given by $\Gamma_S \approx \Gamma_{gg} + 1.7 \times \Gamma_{\gamma\gamma} + \Gamma_{t\bar{t}}$.

We would like to stress that while the diphoton resonance undoubtedly is the motivation behind this work, the purpose of this paper is to construct a consistent LRSM framework that can naturally accommodate vector-like fermions taking the diphoton signal as a hint and explore the consequent phenomenology. One can find similar interpretations of the diphoton excess in [90, 118, 119] in models with a singlet scalar accompanied by vector-like fermions.

Production of a scalar resonance in gluon fusion via a loop of vector-like quarks and subsequent decay of scalar resonance to $\gamma\gamma$ via a loop of vector-like quarks and leptons. There are contributions to $\Gamma(S \rightarrow \gamma\gamma)$ from quark-like vector fermion $\psi_Q = U, D$ and lepton-like vector fermion $\psi_L = E$ propagating inside the loop. Apart from quark-like vector fermion contributing to the production of scalar through gluon fusion, there could be another

top-quark mediated diagram via mixing with SM Higgs boson.

In the LRSM framework discussed in section II, the vector-like degrees of freedom contribute to the loop leading to $S \rightarrow gg$ and $S \rightarrow \gamma\gamma$. The partial decay widths are given by [17]

$$\begin{aligned}\Gamma_{\gamma\gamma} &= \frac{\alpha^2 M_S^3}{256\pi^3} \left| \sum_X \frac{N_X^C Q_X^2 \lambda'_{SXX}}{M_X} \mathcal{A} \left(\frac{m_S^2}{4M_X^2} \right) \right|^2, \\ \Gamma_{gg} &= K \frac{\alpha_s^2 M_S^3}{128\pi^3} \left| \sum_X \frac{\lambda'_{SXX}}{M_X} \mathcal{A} \left(\frac{m_S^2}{4M_X^2} \right) \right|^2.\end{aligned}\quad (9)$$

Here, the sums in $\Gamma_{\gamma\gamma}$ and Γ_{gg} are over all and coloured fermion species and flavours, respectively. N_X^C is the number of color degrees of freedom of a species, i.e. 1 for leptonic vector-like fermions and 3 for quark-like fermions. Similarly, Q_X is the electric charge of the species. The effective coupling of S to a fermion species is $\lambda'_{SXX} = \lambda_{SXX} (O_X^R)_{1X} (O_X^L)_{1X}$, i.e. the coupling λ_{SXX} dressed with the corresponding left and right mixing matrix element. The value of the parameters $\alpha \approx 1/127$, $\alpha_s \approx 0.1$ and $K \approx 1.7$ [17] in which $A(x)$ is a loop function defined by

$$A(x) = \frac{2}{x^2} [x + (x-1)f(x)], \quad (10)$$

with

$$f(x) = \begin{cases} \arcsin^2 \sqrt{x} & x \leq 1 \\ -\frac{1}{4} \left[\ln \left(\frac{1+\sqrt{1-x}}{1-\sqrt{1-x}} \right) - i\pi \right]^2 & x > 1. \end{cases} \quad (11)$$

In addition, the decay of S to a pair of fermions (here only relevant for the top) is given by

$$\Gamma_{f\bar{f}} = \frac{N_f^C \lambda_{Xff}^2 M_S}{16\pi} \left(1 - \frac{4M_f^2}{M_S^2} \right)^{2/3}. \quad (12)$$

In order to arrive at an estimate for the diphoton production cross section, we assume that the vector fermion masses and couplings to S are degenerate (M_X, λ_{SXX}), except for the top partner (M_T, λ_{STT}). In the limit of large vector fermion masses $M_X \gtrsim M_S/2$, we arrive at the approximation for the partial widths,

$$\begin{aligned}\frac{\Gamma_{gg}}{M_S} &\approx 1.3 \times 10^{-4} \left(\frac{\lambda_{SXX} \cdot \text{TeV}}{M_X} \right)^2, \\ \frac{\Gamma_{\gamma\gamma}}{M_S} &\approx 3.4 \times 10^{-7} \left(\frac{\lambda_{SXX} \text{TeV}}{M_X} \right)^2, \\ \frac{\Gamma_{t\bar{t}}}{M_S} &\approx 1.3 \times 10^{-3} \left(\frac{\lambda_{TXX} \text{TeV}}{M_T} \right)^2.\end{aligned}\quad (13)$$

As discussed in [14] in a model-independent fashion, the diphoton excess can be explained for $10^{-6} \lesssim \Gamma_{gg}/M_S \lesssim 2 \times 10^{-3}$ (the upper limit is due to the limit from dijet searches) and $\Gamma_{\gamma\gamma}/M_S \approx 10^{-6}$, as long as gg and $\gamma\gamma$ are the only decay modes of S . In order to achieve this, the

top partner T needs to have a significantly weaker coupling or heavier mass than the rest of the vector fermions. Assuming the decay width to $t\bar{t}$ contributes negligibly to the total width, the diphoton cross section, Eq. (8) is

$$\sigma(pp \rightarrow \gamma\gamma) \approx 1.7 \text{ fb} \cdot \left(\frac{\lambda_{SXX} \cdot \text{TeV}}{M_X} \right)^2, \quad (14)$$

The experimentally suggested cross section $\sigma(pp \rightarrow \gamma\gamma) \approx 7 \text{ fb}$ can be achieved with $M_X/\lambda_{SXX} \approx 0.5 \text{ TeV}$ (and $\Gamma_{gg}/M_S \approx 5 \times 10^{-4}$ satisfying the dijet limit). In such a scenario, the total width of S is of the order $\Gamma_S \approx 0.5 \text{ GeV}$, i.e. much smaller than the 45 GeV suggested by ATLAS if interpreted as a single particle resonance. $\Gamma_{\gamma\gamma}/\Gamma_{gg}$ can also be independently boosted by introducing a hierarchy with leptonic partners lighter than the quark partners. While certainly marginal and requiring a specific structure among the vector fermions, this demonstrates that the diphoton excess, apart from the broad width seen by ATLAS, can be accommodated in our model.

IV. GAUGE COUPLING UNIFICATION

In the previous section, we have discussed how the inclusion of new vector-like fermions in LRSM can aptly explain the diphoton excess traced around 750 GeV at the LHC. Interestingly this framework can also be embedded in a non-SUSY grand unified theory like $SO(10)$ having left-right symmetry as its only intermediate symmetry breaking step with the breaking chain given as follows

$$SO(10) \xrightarrow{\langle \Sigma \rangle} \mathcal{G}_{2213P} \xrightarrow{\langle H_R \rangle} \mathcal{G}_{213} \xrightarrow{\langle H_L \rangle} \mathcal{G}_{13}. \quad (15)$$

The $SO(10)$ group breaks down to left-right symmetric group $\mathcal{G}_{2213P} \equiv SU(2)_L \times SU(2)_R \times U(1)_{B-L} \times SU(3) \times \mathcal{P}$ via a non-zero VEV of $\Sigma \subset 210_H$. Here, \mathcal{P} is defined as the discrete left-right symmetry, a generalized parity symmetry or charge-conjugation symmetry. The vital step is to break the left-right gauge symmetry and this is attained with the help of the right-handed Higgs doublet H_R . Finally, the SM gauge group is spontaneously broken by its left-handed counterpart H_L . As described above we add another scalar singlet S in order to explain the diphoton signal though it is not contributing to the renormalization group (RG) evolution of the gauge couplings.

In addition to the particle content described in Tab. I, we include a bi-triplet $\eta \equiv (3, 3, 0, 1)$ under \mathcal{G}_{2213P} to achieve successful gauge unification. This can be confirmed by using the relevant RG equation for the gauge couplings g_i ,

$$\mu \frac{\partial g_i}{\partial \mu} = \frac{b_i}{16\pi^2} g_i^3, \quad (16)$$




FIG. 1: Gauge coupling running in the considered model accommodating the diphoton excess, demonstrating successful gauge unification at the scale $M_{\text{GUT}} = 10^{17.55}$ GeV with an intermediate left-right symmetry breaking scale at 10 TeV.

where the one-loop beta-coefficients b_i are given by

$$b_i = -\frac{11}{3}\mathcal{C}_2(G) + \frac{2}{3} \sum_{R_f} T(R_f) \prod_{j \neq i} d_j(R_f) + \frac{1}{3} \sum_{R_s} T(R_s) \prod_{j \neq i} d_j(R_s). \quad (17)$$

Here, $\mathcal{C}_2(G)$ is the quadratic Casimir operator for gauge bosons in their adjoint representation,

$$\mathcal{C}_2(G) \equiv \begin{cases} N & \text{if } SU(N), \\ 0 & \text{if } U(1). \end{cases} \quad (18)$$

$T(R_f)$ and $T(R_s)$ are the traces of the irreducible representation $R_{f,s}$ for a given fermion and scalar, respectively,

$$T(R_{f,s}) \equiv \begin{cases} 1/2 & \text{if } R_{f,s} \text{ is fundamental,} \\ N & \text{if } R_{f,s} \text{ is adjoint,} \\ 0 & \text{if } U(1). \end{cases} \quad (19)$$

and $d(R_{f,s})$ is the dimension of a given representation $R_{f,s}$ under all $SU(N)$ gauge groups except the i -th gauge group under consideration. An additional factor of 1/2 should be multiplied in the case of a real Higgs representation. Using the above particle content, the beta-coefficients are found to be $b_{2L} = -19/6$, $b_Y = 41/10$, $b_{3C} = -7$ from the SM to the LR breaking scale and $b_{2L} = b_{2R} = -13/6$, $b_{BL} = 59/6$, $b_{3C} = -17/3$ from the LR breaking scale to the GUT scale. The resulting running of the gauge couplings is displayed in Fig. 1 with the breaking scales

$$M_{\text{GUT}} = 10^{17.55} \text{ GeV}, \quad M_{\text{LR}} = 10 \text{ TeV}. \quad (20)$$

V. IMPLICATIONS FOR BARYOGENESIS AND DARK MATTER

The vector-like fermions added to the spectrum of the LRSM framework can have very profound implications

for a baryogenesis mechanism such as leptogenesis and the dark matter sector. While the proposal of high scale leptogenesis via singlet heavy Majorana neutrinos (or a heavy Higgs triplet) decay added to the SM is beyond the reach of the present and near future collider experiments, the LRSM scenario provides a window of opportunity for low TeV scale leptogenesis testable at the LHC. However, the observation of a 2 TeV W_R boson at the LHC, through confirmation of the 2.8σ signal of two leptons and two jets reported by the CMS collaboration, would rule out the possibility of high scale as well as TeV scale resonant leptogenesis with the standard LRSM fields due to the unavoidable fast gauge mediated $B - L$ violating interactions [135–142].

On the other hand, the new vector-like fermions added to the LRSM to accommodate the diphoton excess can open a whole new world of possibilities. A particularly interesting possibility is the realization of baryogenesis and dark matter annihilation through a vector-like portal first explored in [143]. As an example, consider the following additional terms in the Lagrangian,

$$\mathcal{L} \supset -(\lambda_{XU} X \bar{u}_R U_L + \text{h.c.}) - m_X^2 X^\dagger X - \lambda_X (X^\dagger X)^2 - \lambda_{HX} H^\dagger H X^\dagger X, \quad (21)$$

where X is a singlet complex dark matter scalar field charged under some exotic global $U(1)_\chi$ symmetry, under which only the vector-like quarks and dark matter fields transform non-trivially. Thus, the introduction of a vector-like quark can connect the dark matter and the SM quarks, which can be readily used to make a connection between the baryon asymmetry and dark matter, as pointed out in [143]. In the case of the LRSM, X has to be an inert doublet scalar field. In rest of this section we focus on sketching the simpler case of the extended SM which can be expanded to the LRSM case by replacing the singlets with appropriate doublet representations. However, in the case of the LRSM some subtleties are present and we comment on them towards the end of this section. On the other hand, this idea can easily be generalized to accommodate a down-type quark portal or charged lepton portal (corresponding to a leptogenesis scenario of baryogenesis).

The basic idea behind the vector-like portal is to generate an asymmetry in the vector-like sector through baryogenesis, which then subsequently gets transferred to the SM baryons and the dark matter sector through the renormalizable couplings in Eq. (21). In addition one can introduce a scalar field Y with the couplings

$$\mathcal{L} \supset -(\lambda_{\nu Y} Y \nu_R \nu_R + \text{h.c.}) - m_Y^2 Y^\dagger Y - \lambda_Y (Y^\dagger Y)^2 - \lambda_{XY} X^\dagger X Y^\dagger Y, \quad (22)$$

which allows the annihilation of a pair of X into Y fields. The latter can subsequently decay into two singlet right handed neutrinos ensuring the asymmetric nature of the dark matter X relic density for a large enough annihilation cross section. Now turning to the question of how to generate the primordial asymmetry in the vector-like sector which defines the final dark matter asymmetry and

baryon asymmetry, let us further add two types of heavy diquarks with the couplings

$$\mathcal{L} \supset \lambda_{\Delta U_L} \Delta_u U_L U_L + \lambda_{\Delta U_R} \Delta_u U_R U_R + \lambda_{\Delta d} \Delta_d d_R d_R + \lambda_\chi \Delta_u \Delta_d \Delta_d \chi + \text{h.c.}, \quad (23)$$

where $\Delta_u : (\bar{6}, 1, -4/3)$, $\Delta_d : (\bar{6}, 1, 2/3)$ and the field χ breaks the local $U(1)_\chi$ symmetry under which X and U have non-trivial charges denoted by $q_\chi(U)$ and $q_\chi(X)$. For the SM fields this charge is simply $B-L$, which right away gives $q_\chi(\Delta_d) = -2/3$. The rest of the charges are determined in terms of the free charge $q_\chi(U)$,

$$q_\chi(\Delta_u) = -2q_\chi(U), \quad q_\chi(\chi) = 2q_\chi(U) + 4/3, \\ q_\chi(X) = 1/3 - q_\chi(U). \quad (24)$$

In order too forbid the dangerous proton decay induced by the operators $\mathcal{O} = X^2, S^2, X^2 S^2, X^4, S^4$ [144], one needs to satisfy the condition

$$q_\chi(\mathcal{O}) \neq n(2q_\chi(U) + 4/3), \quad \text{where } n = 0, \pm 1, \pm 2, \dots \quad (25)$$

From Eq. (23) it follows that after χ acquires a VEV to break the $U(1)_\chi$ symmetry, Δ_u has the decay modes

$$\Delta_u \rightarrow \Delta_d^* \Delta_d^*, \quad \Delta_u \rightarrow \bar{U} \bar{U}, \quad (26)$$

and a CP asymmetry (between the above modes and their conjugate modes) can be obtained by interference of the tree level diagrams with one loop self energy diagrams with two generations of Δ_u . Finally, the asymmetry generated in the vector-like quarks gets transferred to the dark matter asymmetry and baryon asymmetry via the λ_{XU} term in Eq. (21). This mechanism gives a ratio between the dark matter relic density and the baryon asymmetry given by

$$\frac{\Omega_{DM}/m_X}{\Omega_B/M_p} = \frac{79}{28}, \quad (27)$$

and in this model a dark matter mass $m_X \sim 2$ GeV. A typical prediction of this model is neutron-antineutron oscillations induced by the up-type and two down-type diquarks through the mixing of vector-like up-type quarks with the usual up quarks. However, such oscillations will be suppressed by the mixing.

One can similarly construct a leptogenesis model involving vector-like charged leptons. In case of the LRSM a generalization of the above scheme is straightforward;

however, the lepton number violating gauge scattering processes involving a low scale W_R can rapidly wash out any primordial asymmetry generated above the mass scale of W_R . In fact, some of these gauge processes can continue to significantly reduce the generation of lepton asymmetry below the mass scale of W_R , thus the vector-like quark portals seem to be more promising in this case. Other alternatives include mechanisms like neutron-antineutron oscillation or some alternative LRSM scheme such as the Alternative Left-Right Symmetric Model [145] where the dangerous gauge scatterings can be avoided by means of special gauge quantum number assignments of a heavy neutrino [146]. Also note that, in general, one can utilize the singlet neutral vector-like lepton as a dark matter candidate by ensuring the stability against decay into usual LRSM fermions.

VI. CONCLUSIONS

We have considered a unified framework to explain the recent diphoton excess reported by ATLAS and CMS around 750 GeV. The addition of vector-like fermions and a singlet scalar S to LRSM but without a scalar bidoublet explains the fermion masses and mixing via a universal seesaw mechanism. The diphoton signal with $\sigma(pp \rightarrow S \rightarrow \gamma\gamma) \approx 4 - 12$ fb can be explained in this model with TeV scale vector fermions. The broad width suggested by the ATLAS excess cannot be understood, though.

We successfully embed this model within an $SO(10)$ GUT framework by introducing a real bi-triplet scalar. This additional scalar, which contains a potential DM candidate, allows the gauge couplings to unify at the scale $10^{17.5}$ GeV. We also discuss further possibilities in this class of LRSM models with vector-like fermions for mechanisms of baryogenesis and DM.

ACKNOWLEDGMENTS

FFD would like to thank Suchita Kulkarni for useful discussions. PP is grateful to the Department of Science and Technology (DST), Govt. of India INSPIRE Fellowship DST/INSPIRE Fellowship/2014/IF140299. The work of SP is partly supported by DST, India under the financial grant SB/S2/HEP-011/2013. The work US is supported partly by the JC Bose National Fellowship grant under DST, India.

-
- [1] **CMS Collaboration**, J. Olsen, “*CMS physics results from Run 2 presented on Dec. 15th. 2015 Search for new physics in high mass diphoton events in proton-proton collisions at 13TeV*,”
[2] **ATLAS Collaboration**, M. Kado, “*ATLAS physics results from Run 2 presented on Dec. 15th 2015 ATLAS-CONF-2015-081*,”

- [3] J. Ellis and D. S. Hwang, “*Does the ‘Higgs’ have Spin Zero?*,” JHEP **09** (2012) 071, [arXiv:1202.6660](#).
[4] S. Di Chiara, L. Marzola, and M. Raidal, “*First interpretation of the 750 GeV di-photon resonance at the LHC*,” [arXiv:1512.04939](#).
[5] A. Pilaftsis, “*Diphoton Signatures from Heavy Axion Decays at LHC*,” [arXiv:1512.04931](#).

- [6] S. Knapen, T. Melia, M. Papucci, and K. Zurek, “*Rays of light from the LHC*,” arXiv:1512.04928.
- [7] M. Backovic, A. Mariotti, and D. Redigolo, “*Di-photon excess illuminates Dark Matter*,” arXiv:1512.04917.
- [8] E. Molinaro, F. Sannino, and N. Vignaroli, “*Strong dynamics or axion origin of the diphoton excess*,” arXiv:1512.05334.
- [9] R. S. Gupta, S. Jger, Y. Kats, G. Perez, and E. Stamou, “*Interpreting a 750 GeV Diphoton Resonance*,” arXiv:1512.05332.
- [10] J. Ellis, S. A. R. Ellis, J. Quevillon, V. Sanz, and T. You, “*On the Interpretation of a Possible ~ 750 GeV Particle Decaying into $\gamma\gamma$* ,” arXiv:1512.05327.
- [11] T. Higaki, K. S. Jeong, N. Kitajima, and F. Takahashi, “*The QCD Axion from Aligned Axions and Diphoton Excess*,” arXiv:1512.05295.
- [12] Y. Mambrini, G. Arcadi, and A. Djouadi, “*The LHC diphoton resonance and dark matter*,” arXiv:1512.04913.
- [13] D. Buttazzo, A. Greljo, and D. Marzocca, “*Knocking on New Physics’ door with a Scalar Resonance*,” arXiv:1512.04929.
- [14] R. Franceschini, G. F. Giudice, J. F. Kamenik, M. McCullough, A. Pomarol, R. Rattazzi, M. Redi, F. Riva, A. Strumia, and R. Torre, “*What is the gamma gamma resonance at 750 GeV?*,” arXiv:1512.04933.
- [15] A. Angelescu, A. Djouadi, and G. Moreau, “*Scenarios for interpretations of the LHC diphoton excess: two Higgs doublets and vector-like quarks and leptons*,” arXiv:1512.04921.
- [16] B. Bellazzini, R. Franceschini, F. Sala, and J. Serra, “*Goldstones in Diphotons*,” arXiv:1512.05330.
- [17] S. D. McDermott, P. Meade, and H. Ramani, “*Singlet Scalar Resonances and the Diphoton Excess*,” arXiv:1512.05326.
- [18] M. Low, A. Tesi, and L.-T. Wang, “*A pseudoscalar decaying to photon pairs in the early LHC run 2 data*,” arXiv:1512.05328.
- [19] C. Petersson and R. Torre, “*The 750 GeV diphoton excess from the goldstino superpartner*,” arXiv:1512.05333.
- [20] Q.-H. Cao, Y. Liu, K.-P. Xie, B. Yan, and D.-M. Zhang, “*A Boost Test of Anomalous Diphoton Resonance at the LHC*,” arXiv:1512.05542.
- [21] A. Kobakhidze, F. Wang, L. Wu, J. M. Yang, and M. Zhang, “*LHC diphoton excess explained as a heavy scalar in top-seesaw model*,” arXiv:1512.05585.
- [22] P. Agrawal, J. Fan, B. Heidenreich, M. Reece, and M. Strassler, “*Experimental Considerations Motivated by the Diphoton Excess at the LHC*,” arXiv:1512.05775.
- [23] W. Chao, R. Huo, and J.-H. Yu, “*The Minimal Scalar-Stealth Top Interpretation of the Diphoton Excess*,” arXiv:1512.05738.
- [24] S. Fichtel, G. von Gersdorff, and C. Royon, “*Scattering Light by Light at 750 GeV at the LHC*,” arXiv:1512.05751.
- [25] D. Curtin and C. B. Verhaaren, “*Quirky Explanations for the Diphoton Excess*,” arXiv:1512.05753.
- [26] C. Csaki, J. Hubisz, and J. Terning, “*The Minimal Model of a Diphoton Resonance: Production without Gluon Couplings*,” arXiv:1512.05776.
- [27] D. Aloni, K. Blum, A. Dery, A. Efrati, and Y. Nir, “*On a possible large width 750 GeV diphoton resonance at ATLAS and CMS*,” arXiv:1512.05778.
- [28] S. V. Demidov and D. S. Gorbunov, “*On goldstino interpretation of the diphoton excess*,” arXiv:1512.05723.
- [29] J. M. No, V. Sanz, and J. Setford, “*See-Saw Composite Higgses at the LHC: Linking Naturalness to the 750 GeV Di-Photon Resonance*,” arXiv:1512.05700.
- [30] Y. Bai, J. Berger, and R. Lu, “*A 750 GeV Dark Pion: Cousin of a Dark G-parity-odd WIMP*,” arXiv:1512.05779.
- [31] S. Matsuzaki and K. Yamawaki, “*750 GeV Diphoton Signal from One-Family Walking Technipion*,” arXiv:1512.05564.
- [32] B. Dutta, Y. Gao, T. Ghosh, I. Gogoladze, and T. Li, “*Interpretation of the diphoton excess at CMS and ATLAS*,” arXiv:1512.05439.
- [33] D. Becirevic, E. Bertuzzo, O. Sumensari, and R. Z. Funchal, “*Can the new resonance at LHC be a CP-Odd Higgs boson?*,” arXiv:1512.05623.
- [34] P. Cox, A. D. Medina, T. S. Ray, and A. Spray, “*Diphoton Excess at 750 GeV from a Radion in the Bulk-Higgs Scenario*,” arXiv:1512.05618.
- [35] R. Martinez, F. Ochoa, and C. F. Sierra, “*Diphoton decay for a 750 GeV scalar dark matter*,” arXiv:1512.05617.
- [36] L. Bian, N. Chen, D. Liu, and J. Shu, “*A hidden confining world on the 750 GeV diphoton excess*,” arXiv:1512.05759.
- [37] J. Chakraborty, A. Choudhury, P. Ghosh, S. Mondal, and T. Srivastava, “*Di-photon resonance around 750 GeV: shedding light on the theory underneath*,” arXiv:1512.05767.
- [38] A. Ahmed, B. M. Dillon, B. Grzadkowski, J. F. Gunion, and Y. Jiang, “*Higgs-radion interpretation of 750 GeV di-photon excess at the LHC*,” arXiv:1512.05771.
- [39] A. Falkowski, O. Slone, and T. Volansky, “*Phenomenology of a 750 GeV Singlet*,” arXiv:1512.05777.
- [40] Q.-H. Cao, C. B. Jackson, W.-Y. Keung, I. Low, and J. Shu, “*The Higgs Mechanism and Loop-induced Decays of a Scalar into Two Z Bosons*,” Phys. Rev. **D81** (2010) 015010, arXiv:0911.3398.
- [41] I. Low, J. Lykken, and G. Shaughnessy, “*Have We Observed the Higgs (Imposter)?*,” Phys. Rev. **D86** (2012) 093012, arXiv:1207.1093.
- [42] W. Chao, “*Symmetries Behind the 750 GeV Diphoton Excess*,” arXiv:1512.06297.
- [43] L. M. Carpenter, R. Colburn, and J. Goodman, “*Supersoft SUSY Models and the 750 GeV Diphoton Excess, Beyond Effective Operators*,” arXiv:1512.06107.
- [44] E. Megias, O. Pujolas, and M. Quiros, “*On dilatons and the LHC diphoton excess*,” arXiv:1512.06106.
- [45] A. Alves, A. G. Dias, and K. Sinha, “*The 750 GeV S-cion: Where else should we look for it?*,” arXiv:1512.06091.
- [46] E. Gabrielli, K. Kannike, B. Mele, M. Raidal, C. Spethmann, and H. Veerme, “*A SUSY Inspired Simplified Model for the 750 GeV Diphoton Excess*,” arXiv:1512.05961.
- [47] J. S. Kim, J. Reuter, K. Rolbiecki, and R. R. de Austri, “*A resonance without resonance: scrutinizing the diphoton excess at 750 GeV*,” arXiv:1512.06083.
- [48] R. Benbrik, C.-H. Chen, and T. Nomura, “*Higgs*

- singlet as a diphoton resonance in a vector-like quark model,” arXiv:1512.06028.
- [49] G. M. Pelaggi, A. Strumia, and E. Vigiani, “Trinification can explain the di-photon and di-boson LHC anomalies,” arXiv:1512.07225.
- [50] A. E. C. Hernandez and I. Nisandzic, “LHC diphoton 750 GeV resonance as an indication of $SU(3)_c \times SU(3)_L \times U(1)_X$ gauge symmetry,” arXiv:1512.07165.
- [51] J. de Blas, J. Santiago, and R. Vega-Morales, “New vector bosons and the diphoton excess,” arXiv:1512.07229.
- [52] P. S. B. Dev and D. Teresi, “Asymmetric Dark Matter in the Sun and the Diphoton Excess at the LHC,” arXiv:1512.07243.
- [53] S. M. Boucenna, S. Morisi, and A. Vicente, “The LHC diphoton resonance from gauge symmetry,” arXiv:1512.06878.
- [54] K. Kulkarni, “Extension of ν MSM model and possible explanations of recent astronomical and collider observations,” arXiv:1512.06836.
- [55] M. Chala, M. Duerr, F. Kahlhoefer, and K. Schmidt-Hoberg, “Tricking Landau-Yang: How to obtain the diphoton excess from a vector resonance,” arXiv:1512.06833.
- [56] M. Bauer and M. Neubert, “Flavor Anomalies, the Diphoton Excess and a Dark Matter Candidate,” arXiv:1512.06828.
- [57] J. M. Cline and Z. Liu, “LHC diphotons from electroweakly pair-produced composite pseudoscalars,” arXiv:1512.06827.
- [58] L. Berthier, J. M. Cline, W. Shepherd, and M. Trott, “Effective interpretations of a diphoton excess,” arXiv:1512.06799.
- [59] J. S. Kim, K. Rolbiecki, and R. R. de Austri, “Model-independent combination of diphoton constraints at 750 GeV,” arXiv:1512.06797.
- [60] X.-J. Bi, Q.-F. Xiang, P.-F. Yin, and Z.-H. Yu, “The 750 GeV diphoton excess at the LHC and dark matter constraints,” arXiv:1512.06787.
- [61] J. J. Heckman, “750 GeV Diphotons from a $D3$ -brane,” arXiv:1512.06773.
- [62] F. P. Huang, C. S. Li, Z. L. Liu, and Y. Wang, “750 GeV Diphoton Excess from Cascade Decay,” arXiv:1512.06732.
- [63] J. Cao, C. Han, L. Shang, W. Su, J. M. Yang, and Y. Zhang, “Interpreting the 750 GeV diphoton excess by the singlet extension of the Manohar-Wise Model,” arXiv:1512.06728.
- [64] F. Wang, L. Wu, J. M. Yang, and M. Zhang, “750 GeV Diphoton Resonance, 125 GeV Higgs and Muon $g-2$ Anomaly in Deflected Anomaly Mediation SUSY Breaking Scenario,” arXiv:1512.06715.
- [65] O. Antipin, M. Mojaza, and F. Sannino, “A natural Coleman-Weinberg theory explains the diphoton excess,” arXiv:1512.06708.
- [66] X.-F. Han and L. Wang, “Implication of the 750 GeV diphoton resonance on two-Higgs-doublet model and its extensions with Higgs field,” arXiv:1512.06587.
- [67] R. Ding, L. Huang, T. Li, and B. Zhu, “Interpreting 750 GeV Diphoton Excess with R -parity Violation Supersymmetry,” arXiv:1512.06560.
- [68] I. Chakraborty and A. Kundu, “Diphoton excess at 750 GeV: Singlet scalars confront naturalness,” arXiv:1512.06508.
- [69] D. Barducci, A. Goudelis, S. Kulkarni, and D. Sengupta, “One jet to rule them all: monojet constraints and invisible decays of a 750 GeV diphoton resonance,” arXiv:1512.06842.
- [70] W. S. Cho, D. Kim, K. Kong, S. H. Lim, K. T. Matchev, J.-C. Park, and M. Park, “The 750 GeV Diphoton Excess May Not Imply a 750 GeV Resonance,” arXiv:1512.06824.
- [71] T.-F. Feng, X.-Q. Li, H.-B. Zhang, and S.-M. Zhao, “The LHC 750 GeV diphoton excess in supersymmetry with gauged baryon and lepton numbers,” arXiv:1512.06696.
- [72] D. Bardhan, D. Bhatia, A. Chakraborty, U. Maitra, S. Raychaudhuri, and T. Samui, “Radion Candidate for the LHC Diphoton Resonance,” arXiv:1512.06674.
- [73] H. Han, S. Wang, and S. Zheng, “Scalar Explanation of Diphoton Excess at LHC,” arXiv:1512.06562.
- [74] M. Dhuria and G. Goswami, “Perturbativity, vacuum stability and inflation in the light of 750 GeV diphoton excess,” arXiv:1512.06782.
- [75] S. Chang, “A Simple $U(1)$ Gauge Theory Explanation of the Diphoton Excess,” arXiv:1512.06426.
- [76] C. Han, H. M. Lee, M. Park, and V. Sanz, “The diphoton resonance as a gravity mediator of dark matter,” arXiv:1512.06376.
- [77] M. T. Arun and P. Saha, “Gravitons in multiply warped scenarios - at 750 GeV and beyond,” arXiv:1512.06335.
- [78] G. Li, Y.-n. Mao, Y.-L. Tang, C. Zhang, Y. Zhou, and S.-h. Zhu, “A Loop-philic Pseudoscalar,” arXiv:1512.08255.
- [79] H. Han, S. Wang, and S. Zheng, “Dark Matter Theories in the Light of Diphoton Excess,” arXiv:1512.07992.
- [80] J. A. Casas, J. R. Espinosa, and J. M. Moreno, “The 750 GeV Diphoton Excess as a First Light on Supersymmetry Breaking,” arXiv:1512.07895.
- [81] J. Zhang and S. Zhou, “Electroweak Vacuum Stability and Diphoton Excess at 750 GeV,” arXiv:1512.07889.
- [82] J. Liu, X.-P. Wang, and W. Xue, “LHC diphoton excess from colorful resonances,” arXiv:1512.07885.
- [83] K. Das and S. K. Rai, “The 750 GeV Diphoton excess in a $U(1)$ hidden symmetry model,” arXiv:1512.07789.
- [84] H. Davoudiasl and C. Zhang, “A 750 GeV Messenger of Dark Conformal Symmetry Breaking,” arXiv:1512.07672.
- [85] M. Cveti, J. Halverson, and P. Langacker, “String Consistency, Heavy Exotics, and the 750 GeV Diphoton Excess at the LHC,” arXiv:1512.07622.
- [86] M. Badziak, “Interpreting the 750 GeV diphoton excess in minimal extensions of Two-Higgs-Doublet models,” arXiv:1512.07497.
- [87] K. M. Patel and P. Sharma, “Interpreting 750 GeV diphoton excess in $SU(5)$ grand unified theory,” arXiv:1512.07468.
- [88] S. Moretti and K. Yagyu, “The 750 GeV diphoton excess and its explanation in 2-Higgs Doublet Models with a real inert scalar multiplet,” arXiv:1512.07462.
- [89] J. Gu and Z. Liu, “Running after Diphoton,” arXiv:1512.07624.
- [90] P. S. B. Dev, R. N. Mohapatra, and Y. Zhang, “Quark Seesaw Vectorlike Fermions and Diphoton Excess,” arXiv:1512.08507.
- [91] N. Bizot, S. Davidson, M. Frigerio, and J. L. Kneur,

- “Two Higgs doublets to explain the excesses $pp \rightarrow \gamma\gamma(750 \text{ GeV})$ and $h \rightarrow \tau^\pm \mu^\mp$,” [arXiv:1512.08508](#).
- [92] L. A. Anchordoqui, I. Antoniadis, H. Goldberg, X. Huang, D. Lust, and T. R. Taylor, “750 GeV diphotons from closed string states,” [arXiv:1512.08502](#).
- [93] F. Goertz, J. F. Kamenik, A. Katz, and M. Nardecchia, “Indirect Constraints on the Scalar Di-Photon Resonance at the LHC,” [arXiv:1512.08500](#).
- [94] X.-J. Bi, R. Ding, Y. Fan, L. Huang, C. Li, T. Li, S. Raza, X.-C. Wang, and B. Zhu, “A Promising Interpretation of Diphoton Resonance at 750 GeV,” [arXiv:1512.08497](#).
- [95] W. Chao, “Neutrino Catalyzed Diphoton Excess,” [arXiv:1512.08484](#).
- [96] J. Gao, H. Zhang, and H. X. Zhu, “Diphoton excess at 750 GeV: gluon-gluon fusion or quark-antiquark annihilation?,” [arXiv:1512.08478](#).
- [97] J. E. Kim, “Is an axizilla possible for di-photon resonance?,” [arXiv:1512.08467](#).
- [98] Q.-H. Cao, Y. Liu, K.-P. Xie, B. Yan, and D.-M. Zhang, “The Diphoton Excess, Low Energy Theorem and the 331 Model,” [arXiv:1512.08441](#).
- [99] C. Cai, Z.-H. Yu, and H.-H. Zhang, “The 750 GeV diphoton resonance as a singlet scalar in an extra dimensional model,” [arXiv:1512.08440](#).
- [100] F. Wang, W. Wang, L. Wu, J. M. Yang, and M. Zhang, “Interpreting 750 GeV Diphoton Resonance in the NMSSM with Vector-like Particles,” [arXiv:1512.08434](#).
- [101] J. Cao, F. Wang, and Y. Zhang, “Interpreting The 750 GeV Diphoton Excess Within TopFlavor Seesaw Model,” [arXiv:1512.08392](#).
- [102] H. An, C. Cheung, and Y. Zhang, “Broad Diphotons from Narrow States,” [arXiv:1512.08378](#).
- [103] Y.-L. Tang and S.-h. Zhu, “NMSSM extended with vector-like particles and the diphoton excess on the LHC,” [arXiv:1512.08323](#).
- [104] M. Son and A. Urbano, “A new scalar resonance at 750 GeV: Towards a proof of concept in favor of strongly interacting theories,” [arXiv:1512.08307](#).
- [105] J.-C. Park and S. C. Park, “Indirect signature of dark matter with the diphoton resonance at 750 GeV,” [arXiv:1512.08117](#).
- [106] A. Salvio and A. Mazumdar, “Higgs Stability and the 750 GeV Diphoton Excess,” [arXiv:1512.08184](#).
- [107] D. Chway, R. Dermek, T. H. Jung, and H. D. Kim, “Glue to light signal of a new particle,” [arXiv:1512.08221](#).
- [108] E. Ma, “Diphoton Revelation of the Utilitarian Supersymmetric Standard Model,” [arXiv:1512.09159](#).
- [109] K. Kaneta, S. Kang, and H.-S. Lee, “Diphoton excess at the LHC Run 2 and its implications for a new heavy gauge boson,” [arXiv:1512.09129](#).
- [110] A. E. C. Hernandez, “The 750 GeV diphoton resonance can cause the SM fermion mass and mixing pattern,” [arXiv:1512.09092](#).
- [111] I. Low and J. Lykken, “Implications of Gauge Invariance on a Heavy Diphoton Resonance,” [arXiv:1512.09089](#).
- [112] P. V. Dong and N. T. K. Ngan, “Phenomenology of the simple 3-3-1 model with inert scalars,” [arXiv:1512.09073](#).
- [113] S. Kanemura, N. Machida, S. Odori, and T. Shindou, “Diphoton excess at 750 GeV in an extended scalar sector,” [arXiv:1512.09053](#).
- [114] S. Kanemura, K. Nishiwaki, H. Okada, Y. Orikasa, S. C. Park, and R. Watanabe, “LHC 750 GeV Diphoton excess in a radiative seesaw model,” [arXiv:1512.09048](#).
- [115] S. K. Kang and J. Song, “Top-phobic heavy Higgs boson as the 750 GeV diphoton resonance,” [arXiv:1512.08963](#).
- [116] C.-W. Chiang, M. Ibe, and T. T. Yanagida, “Revisiting Scalar Quark Hidden Sector in Light of 750-GeV Diphoton Resonance,” [arXiv:1512.08895](#).
- [117] Y. Hamada, T. Noumi, S. Sun, and G. Shiu, “An $O(750)$ GeV Resonance and Inflation,” [arXiv:1512.08984](#).
- [118] Q.-H. Cao, S.-L. Chen, and P.-H. Gu, “Strong CP Problem, Neutrino Masses and the 750 GeV Diphoton Resonance,” [arXiv:1512.07541](#).
- [119] A. Dasgupta, M. Mitra, and D. Borah, “Minimal Left-Right Symmetry Confronted with the 750 GeV Di-photon Excess at LHC,” [arXiv:1512.09202](#).
- [120] CMS, V. Khachatryan et al., “Search for heavy neutrinos and W bosons with right-handed couplings in proton-proton collisions at $\sqrt{s} = 8 \text{ TeV}$,” *Eur. Phys. J. C* **74** (2014) no. 11, 3149, [arXiv:1407.3683](#).
- [121] F. F. Deppisch, T. E. Gonzalo, S. Patra, N. Sahu, and U. Sarkar, “Signal of Right-Handed Charged Gauge Bosons at the LHC?,” *Phys. Rev.* **D90** (2014) no. 5, 053014, [arXiv:1407.5384](#).
- [122] F. F. Deppisch, T. E. Gonzalo, S. Patra, N. Sahu, and U. Sarkar, “Double beta decay, lepton flavor violation, and collider signatures of left-right symmetric models with spontaneous D-parity breaking,” *Phys. Rev.* **D91** (2015) no. 1, 015018, [arXiv:1410.6427](#).
- [123] P. S. Bhupal Dev and R. N. Mohapatra, “Unified explanation of the $e\bar{e}j\bar{j}$, diboson and dijet resonances at the LHC,” *Phys. Rev. Lett.* **115** (2015) no. 18, 181803, [arXiv:1508.02277](#).
- [124] D. Guadagnoli, R. N. Mohapatra, and I. Sung, “Gauged Flavor Group with Left-Right Symmetry,” *JHEP* **04** (2011) 093, [arXiv:1103.4170](#).
- [125] G. C. Joshi and R. R. Volkas, “Extended weak isospin and fermion masses in a unified model,” *Phys. Rev.* **D45** (1992) 1711–1719.
- [126] G. Aldazabal, L. E. Ibanez, and F. Quevedo, “A D^- brane alternative to the MSSM,” *JHEP* **02** (2000) 015, [arXiv:hep-ph/0001083](#).
- [127] J. P. Conlon, A. Maharana, and F. Quevedo, “Towards Realistic String Vacua,” *JHEP* **05** (2009) 109, [arXiv:0810.5660](#).
- [128] J. C. Pati and A. Salam, “Lepton Number as the Fourth Color,” *Phys.Rev.* **D10** (1974) 275–289.
- [129] R. Mohapatra and J. C. Pati, “A Natural Left-Right Symmetry,” *Phys.Rev.* **D11** (1975) 2558.
- [130] G. Senjanović and R. N. Mohapatra, “Exact Left-Right Symmetry and Spontaneous Violation of Parity,” *Phys.Rev.* **D12** (1975) 1502.
- [131] G. Senjanović, “Spontaneous Breakdown of Parity in a Class of Gauge Theories,” *Nucl.Phys.* **B153** (1979) 334–364.
- [132] R. N. Mohapatra and G. Senjanović, “Neutrino Mass and Spontaneous Parity Violation,” *Phys.Rev.Lett.* **44** (1980) 912.
- [133] R. N. Mohapatra and G. Senjanovic, “Neutrino Masses

- and Mixings in Gauge Models with Spontaneous Parity Violation,* Phys. Rev. **D23** (1981) 165.
- [134] S. Patra, “*Neutrinoless double beta decay process in left-right symmetric models without scalar bidoublet,*” Phys. Rev. **D87** (2013) no. 1, 015002, [arXiv:1212.0612](#).
- [135] E. Ma, S. Sarkar, and U. Sarkar, “*Scale of $SU(2)(R)$ symmetry breaking and leptogenesis,*” Phys. Lett. **B458** (1999) 73–78, [arXiv:hep-ph/9812276](#).
- [136] J.-M. Frere, T. Hambye, and G. Vertongen, “*Is leptogenesis falsifiable at LHC?,*” JHEP **01** (2009) 051, [arXiv:0806.0841](#).
- [137] F. F. Deppisch, J. Harz, M. Hirsch, W.-C. Huang, and H. Pas, “*Falsifying High-Scale Baryogenesis with Neutrinoless Double Beta Decay and Lepton Flavor Violation,*” Phys. Rev. **D92** (2015) no. 3, 036005, [arXiv:1503.04825](#).
- [138] F. F. Deppisch, J. Harz, and M. Hirsch, “*Falsifying High-Scale Leptogenesis at the LHC,*” Phys. Rev. Lett. **112** (2014) 221601, [arXiv:1312.4447](#).
- [139] P. S. Bhupal Dev, C.-H. Lee, and R. N. Mohapatra, “*Leptogenesis Constraints on the Mass of Right-handed Gauge Bosons,*” Phys. Rev. **D90** (2014) no. 9, 095012, [arXiv:1408.2820](#).
- [140] M. Dhuria, C. Hati, R. Rangarajan, and U. Sarkar, “*The $eejj$ Excess Signal at the LHC and Constraints on Leptogenesis,*” JCAP **1509** (2015) no. 09, 035, [arXiv:1502.01695](#).
- [141] M. Dhuria, C. Hati, R. Rangarajan, and U. Sarkar, “*Falsifying leptogenesis for a TeV scale W_R^\pm at the LHC,*” Phys. Rev. **D92** (2015) no. 3, 031701, [arXiv:1503.07198](#).
- [142] P. S. Bhupal Dev, C.-H. Lee, and R. N. Mohapatra, “*TeV Scale Lepton Number Violation and Baryogenesis,*” J. Phys. Conf. Ser. **631** (2015) no. 1, 012007, [arXiv:1503.04970](#).
- [143] P. Fileviez Perez and M. B. Wise, “*Baryon Asymmetry and Dark Matter Through the Vector-Like Portal,*” JHEP **05** (2013) 094, [arXiv:1303.1452](#).
- [144] J. M. Arnold, B. Fornal, and M. B. Wise, “*Simplified models with baryon number violation but no proton decay,*” Phys. Rev. **D87** (2013) 075004, [arXiv:1212.4556](#).
- [145] E. Ma, “*Particle Dichotomy and Left-Right Decomposition of $E(6)$ Superstring Models,*” Phys. Rev. **D36** (1987) 274.
- [146] M. Dhuria, C. Hati, R. Rangarajan, and U. Sarkar, “*Explaining the CMS $eejj$ and e missing p_T jj excess and leptogenesis in superstring inspired E_6 models,*” Phys. Rev. **D91** (2015) no. 5, 055010, [arXiv:1501.04815](#).